



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 02:11 am BST

PDB ID : 3AZE  
Title : Crystal Structure of Human Nucleosome Core Particle Containing H3K64Q mutation  
Authors : Iwasaki, W.; Tachiwana, H.; Kawaguchi, K.; Shibata, T.; Kagawa, W.; Kurumizaka, H.  
Deposited on : 2011-05-25  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	<div> <div>34%</div> <div>32%</div> <div>•</div> <div>30%</div> </div>
1	E	139	<div> <div>40%</div> <div>28%</div> <div>•</div> <div>29%</div> </div>
2	B	106	<div> <div>23%</div> <div>47%</div> <div>•</div> <div>27%</div> </div>
2	F	106	<div> <div>37%</div> <div>42%</div> <div>•</div> <div>20%</div> </div>
3	C	133	<div> <div>42%</div> <div>35%</div> <div>•</div> <div>19%</div> </div>
3	G	133	<div> <div>37%</div> <div>35%</div> <div>6%</div> <div>22%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	129	<div><div></div><div>41%29%•27%</div></div>
4	H	129	<div><div></div><div>27%40%•29%</div></div>
5	I	146	<div><div></div><div>10%7%93%</div></div>
5	J	146	<div><div></div><div>8%8%90%••</div></div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	97	Total	C	N	O	S	0	0	0
			801	504	155	138	4			
1	E	99	Total	C	N	O	S	0	0	0
			816	513	158	141	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP P68431
A	-2	SER	-	EXPRESSION TAG	UNP P68431
A	-1	HIS	-	EXPRESSION TAG	UNP P68431
A	64	GLN	LYS	ENGINEERED MUTATION	UNP P68431
E	-3	GLY	-	EXPRESSION TAG	UNP P68431
E	-2	SER	-	EXPRESSION TAG	UNP P68431
E	-1	HIS	-	EXPRESSION TAG	UNP P68431
E	64	GLN	LYS	ENGINEERED MUTATION	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	77	Total	C	N	O	S	0	0	0
			614	389	119	105	1			
2	F	85	Total	C	N	O	S	0	0	0
			683	430	136	116	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	EXPRESSION TAG	UNP P62805
B	-2	SER	-	EXPRESSION TAG	UNP P62805
B	-1	HIS	-	EXPRESSION TAG	UNP P62805
F	-3	GLY	-	EXPRESSION TAG	UNP P62805

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	-2	SER	-	EXPRESSION TAG	UNP P62805
F	-1	HIS	-	EXPRESSION TAG	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	108	Total	C	N	O	0	0	0
			835	526	165	144			
3	G	104	Total	C	N	O	0	0	0
			805	508	157	140			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	EXPRESSION TAG	UNP P04908
C	-2	SER	-	EXPRESSION TAG	UNP P04908
C	-1	HIS	-	EXPRESSION TAG	UNP P04908
G	-3	GLY	-	EXPRESSION TAG	UNP P04908
G	-2	SER	-	EXPRESSION TAG	UNP P04908
G	-1	HIS	-	EXPRESSION TAG	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	94	Total	C	N	O	S	0	0	0
			736	462	134	138	2			
4	H	92	Total	C	N	O	S	0	0	0
			719	453	129	135	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	EXPRESSION TAG	UNP P06899
D	-2	SER	-	EXPRESSION TAG	UNP P06899
D	-1	HIS	-	EXPRESSION TAG	UNP P06899
H	-3	GLY	-	EXPRESSION TAG	UNP P06899
H	-2	SER	-	EXPRESSION TAG	UNP P06899
H	-1	HIS	-	EXPRESSION TAG	UNP P06899

- Molecule 5 is a DNA chain called 146-MER DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	146	Total	C	N	O	P	0	0	0
			2990	1431	540	874	145			
5	J	144	Total	C	N	O	P	0	0	0
			2949	1411	533	862	143			

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	1	Total	Cl	0	0
			1	1		
6	A	1	Total	Cl	0	0
			1	1		
6	D	1	Total	Cl	0	0
			1	1		
6	E	1	Total	Cl	0	0
			1	1		

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	J	4	Total	Mn	0	0
			4	4		
7	I	5	Total	Mn	0	0
			5	5		
7	D	1	Total	Mn	0	0
			1	1		



- Molecule 1: Histone H3.1






- Molecule 3: Histone H2A type 1-B/E





- Molecule 3: Histone H2A type 1-B/E




- Molecule 4: Histone H2B type 1-J



- Molecule 4: Histone H2B type 1-J




- Molecule 5: 146-MER DNA







● Molecule 5: 146-MER DNA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.14Å 109.34Å 175.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.08 – 3.00 38.08 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.08-3.00) 99.8 (38.08-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.60 (at 3.01Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.244 , 0.302 0.245 , 0.303	Depositor DCC
$R_{free}$ test set	2099 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	65.0	Xtriage
Anisotropy	0.547	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 58.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.049 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	11962	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/813	0.61	0/1091
1	E	0.43	0/828	0.64	0/1110
2	B	0.38	0/621	0.64	0/832
2	F	0.43	0/691	0.68	0/923
3	C	0.38	0/845	0.65	0/1139
3	G	0.37	0/815	0.63	0/1100
4	D	0.42	0/747	0.64	0/1004
4	H	0.40	0/730	0.66	0/982
5	I	0.35	0/3354	0.74	0/5175
5	J	0.34	0/3308	0.75	0/5104
All	All	0.37	0/12752	0.70	0/18460

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
5	J	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	57	TYR	Sidechain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
5	J	214	DG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	834	92	0
1	E	816	0	851	79	0
2	B	614	0	656	84	0
2	F	683	0	729	63	0
3	C	835	0	897	63	0
3	G	805	0	861	71	0
4	D	736	0	758	52	0
4	H	719	0	740	64	0
5	I	2990	0	1652	281	0
5	J	2949	0	1629	278	0
6	A	1	0	0	1	0
6	D	1	0	0	0	0
6	E	1	0	0	1	0
6	G	1	0	0	0	0
7	D	1	0	0	0	0
7	I	5	0	0	0	0
7	J	4	0	0	0	0
All	All	11962	0	9607	939	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

The worst 5 of 939 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:132:DC:H2"	5:I:133:DA:H5'	1.20	1.20
5:I:101:DC:H2"	5:I:102:DA:H5"	1.19	1.15
5:I:138:DG:H2"	5:I:139:DA:H5"	1.17	1.14
5:J:158:DC:H2"	5:J:159:DC:H5"	1.27	1.13
5:J:252:DT:H2"	5:J:253:DC:H5"	1.26	1.13

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/139 (68%)	82 (86%)	11 (12%)	2 (2%)	7	33
1	E	97/139 (70%)	89 (92%)	7 (7%)	1 (1%)	15	53
2	B	75/106 (71%)	67 (89%)	7 (9%)	1 (1%)	12	45
2	F	83/106 (78%)	75 (90%)	8 (10%)	0	100	100
3	C	106/133 (80%)	90 (85%)	15 (14%)	1 (1%)	17	55
3	G	102/133 (77%)	82 (80%)	17 (17%)	3 (3%)	4	24
4	D	92/129 (71%)	82 (89%)	9 (10%)	1 (1%)	14	50
4	H	90/129 (70%)	75 (83%)	11 (12%)	4 (4%)	2	15
All	All	740/1014 (73%)	642 (87%)	85 (12%)	13 (2%)	8	37

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	86	SER
4	H	55	SER
4	H	104	GLY
1	A	86	SER
1	A	117	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/113 (75%)	81 (95%)	4 (5%)	26	63
1	E	86/113 (76%)	81 (94%)	5 (6%)	20	55
2	B	63/81 (78%)	60 (95%)	3 (5%)	25	62
2	F	70/81 (86%)	67 (96%)	3 (4%)	29	66
3	C	85/102 (83%)	79 (93%)	6 (7%)	14	46
3	G	83/102 (81%)	74 (89%)	9 (11%)	6	26
4	D	80/107 (75%)	75 (94%)	5 (6%)	18	51
4	H	78/107 (73%)	73 (94%)	5 (6%)	17	51
All	All	630/806 (78%)	590 (94%)	40 (6%)	18	51

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	59	GLU
1	E	134	ARG
4	H	34	LYS
1	E	76	GLN
2	F	19	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	49	HIS
1	E	68	GLN
4	H	49	HIS
3	C	38	ASN
3	G	73	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	97/139 (69%)	-0.23	0 100 100	41, 63, 86, 114	0
1	E	99/139 (71%)	-0.27	0 100 100	30, 46, 73, 104	0
2	B	77/106 (72%)	-0.22	0 100 100	35, 60, 81, 89	0
2	F	85/106 (80%)	-0.22	1 (1%) 79 54	29, 43, 62, 102	0
3	C	108/133 (81%)	-0.29	0 100 100	35, 52, 86, 112	0
3	G	104/133 (78%)	-0.29	0 100 100	39, 58, 87, 96	0
4	D	94/129 (72%)	-0.28	0 100 100	30, 55, 83, 106	0
4	H	92/129 (71%)	-0.26	0 100 100	41, 58, 84, 102	0
5	I	146/146 (100%)	0.57	14 (9%) 8 2	43, 126, 174, 195	0
5	J	144/146 (98%)	0.51	12 (8%) 11 3	64, 124, 171, 186	0
All	All	1046/1306 (80%)	-0.04	27 (2%) 56 27	29, 62, 152, 195	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	I	146	DT	8.4
5	J	161	DG	4.1
5	I	55	DA	3.6
5	J	169	DT	3.6
5	J	241	DA	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	MN	I	1004	1/1	0.63	0.09	113,113,113,113	0
7	MN	J	1001	1/1	0.73	0.23	138,138,138,138	0
7	MN	I	1003	1/1	0.75	0.16	106,106,106,106	0
7	MN	I	1001	1/1	0.79	0.14	125,125,125,125	0
7	MN	I	1002	1/1	0.82	0.26	137,137,137,137	0
7	MN	J	1004	1/1	0.89	0.19	98,98,98,98	0
7	MN	J	1002	1/1	0.91	0.22	114,114,114,114	0
7	MN	I	1005	1/1	0.91	0.27	142,142,142,142	0
6	CL	D	202	1/1	0.93	0.13	58,58,58,58	0
7	MN	J	1003	1/1	0.93	0.08	93,93,93,93	0
7	MN	D	201	1/1	0.94	0.32	60,60,60,60	0
6	CL	A	1001	1/1	0.94	0.12	60,60,60,60	0
6	CL	G	1001	1/1	0.97	0.10	59,59,59,59	0
6	CL	E	1001	1/1	0.97	0.21	47,47,47,47	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.