



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 24, 2021 – 01:09 pm GMT

PDB ID : 7AZK  
Title : DNA polymerase sliding clamp from Escherichia coli with peptide 35 bound  
Authors : Monsarrat, C.; Compain, G.; Andre, C.; Martiel, I.; Engilberge, S.; Olieric, V.; Wolff, P.; Brillet, K.; Landolfo, M.; Silva da Veiga, C.; Wagner, J.; Guichard, G.; Burnouf, D.Y.  
Deposited on : 2020-11-16  
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4 (270009), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	<div> <div>4%</div> <div>84%</div> <div>12%</div> <div>5%</div> </div>
1	B	386	<div> <div>8%</div> <div>87%</div> <div>7%</div> <div>5%</div> </div>
1	C	386	<div> <div>6%</div> <div>87%</div> <div>7%</div> <div>5%</div> </div>
1	D	386	<div> <div>5%</div> <div>85%</div> <div>9%</div> <div>6%</div> </div>
2	H	6	<div> <div>83%</div> <div>17%</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain	
2	I	6	<div><div></div></div>	33%
2	J	6	<div><div></div></div>	33%
2	K	6	<div><div></div></div>	33%

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	4	0
			2887	1811	510	547	19			
1	B	365	Total	C	N	O	S	0	5	0
			2844	1792	496	536	20			
1	C	365	Total	C	N	O	S	0	2	0
			2851	1791	503	538	19			
1	D	364	Total	C	N	O	S	0	3	0
			2850	1793	501	537	19			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A073FMV0
A	-18	GLY	-	expression tag	UNP A0A073FMV0
A	-17	SER	-	expression tag	UNP A0A073FMV0
A	-16	SER	-	expression tag	UNP A0A073FMV0
A	-15	HIS	-	expression tag	UNP A0A073FMV0
A	-14	HIS	-	expression tag	UNP A0A073FMV0
A	-13	HIS	-	expression tag	UNP A0A073FMV0
A	-12	HIS	-	expression tag	UNP A0A073FMV0
A	-11	HIS	-	expression tag	UNP A0A073FMV0
A	-10	HIS	-	expression tag	UNP A0A073FMV0
A	-9	SER	-	expression tag	UNP A0A073FMV0
A	-8	SER	-	expression tag	UNP A0A073FMV0
A	-7	GLY	-	expression tag	UNP A0A073FMV0
A	-6	LEU	-	expression tag	UNP A0A073FMV0
A	-5	VAL	-	expression tag	UNP A0A073FMV0
A	-4	PRO	-	expression tag	UNP A0A073FMV0
A	-3	ARG	-	expression tag	UNP A0A073FMV0
A	-2	GLY	-	expression tag	UNP A0A073FMV0
A	-1	SER	-	expression tag	UNP A0A073FMV0
A	0	HIS	-	expression tag	UNP A0A073FMV0
B	-19	MET	-	initiating methionine	UNP A0A073FMV0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP A0A073FMV0
B	-17	SER	-	expression tag	UNP A0A073FMV0
B	-16	SER	-	expression tag	UNP A0A073FMV0
B	-15	HIS	-	expression tag	UNP A0A073FMV0
B	-14	HIS	-	expression tag	UNP A0A073FMV0
B	-13	HIS	-	expression tag	UNP A0A073FMV0
B	-12	HIS	-	expression tag	UNP A0A073FMV0
B	-11	HIS	-	expression tag	UNP A0A073FMV0
B	-10	HIS	-	expression tag	UNP A0A073FMV0
B	-9	SER	-	expression tag	UNP A0A073FMV0
B	-8	SER	-	expression tag	UNP A0A073FMV0
B	-7	GLY	-	expression tag	UNP A0A073FMV0
B	-6	LEU	-	expression tag	UNP A0A073FMV0
B	-5	VAL	-	expression tag	UNP A0A073FMV0
B	-4	PRO	-	expression tag	UNP A0A073FMV0
B	-3	ARG	-	expression tag	UNP A0A073FMV0
B	-2	GLY	-	expression tag	UNP A0A073FMV0
B	-1	SER	-	expression tag	UNP A0A073FMV0
B	0	HIS	-	expression tag	UNP A0A073FMV0
C	-19	MET	-	initiating methionine	UNP A0A073FMV0
C	-18	GLY	-	expression tag	UNP A0A073FMV0
C	-17	SER	-	expression tag	UNP A0A073FMV0
C	-16	SER	-	expression tag	UNP A0A073FMV0
C	-15	HIS	-	expression tag	UNP A0A073FMV0
C	-14	HIS	-	expression tag	UNP A0A073FMV0
C	-13	HIS	-	expression tag	UNP A0A073FMV0
C	-12	HIS	-	expression tag	UNP A0A073FMV0
C	-11	HIS	-	expression tag	UNP A0A073FMV0
C	-10	HIS	-	expression tag	UNP A0A073FMV0
C	-9	SER	-	expression tag	UNP A0A073FMV0
C	-8	SER	-	expression tag	UNP A0A073FMV0
C	-7	GLY	-	expression tag	UNP A0A073FMV0
C	-6	LEU	-	expression tag	UNP A0A073FMV0
C	-5	VAL	-	expression tag	UNP A0A073FMV0
C	-4	PRO	-	expression tag	UNP A0A073FMV0
C	-3	ARG	-	expression tag	UNP A0A073FMV0
C	-2	GLY	-	expression tag	UNP A0A073FMV0
C	-1	SER	-	expression tag	UNP A0A073FMV0
C	0	HIS	-	expression tag	UNP A0A073FMV0
D	-19	MET	-	initiating methionine	UNP A0A073FMV0
D	-18	GLY	-	expression tag	UNP A0A073FMV0
D	-17	SER	-	expression tag	UNP A0A073FMV0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP A0A073FMV0
D	-15	HIS	-	expression tag	UNP A0A073FMV0
D	-14	HIS	-	expression tag	UNP A0A073FMV0
D	-13	HIS	-	expression tag	UNP A0A073FMV0
D	-12	HIS	-	expression tag	UNP A0A073FMV0
D	-11	HIS	-	expression tag	UNP A0A073FMV0
D	-10	HIS	-	expression tag	UNP A0A073FMV0
D	-9	SER	-	expression tag	UNP A0A073FMV0
D	-8	SER	-	expression tag	UNP A0A073FMV0
D	-7	GLY	-	expression tag	UNP A0A073FMV0
D	-6	LEU	-	expression tag	UNP A0A073FMV0
D	-5	VAL	-	expression tag	UNP A0A073FMV0
D	-4	PRO	-	expression tag	UNP A0A073FMV0
D	-3	ARG	-	expression tag	UNP A0A073FMV0
D	-2	GLY	-	expression tag	UNP A0A073FMV0
D	-1	SER	-	expression tag	UNP A0A073FMV0
D	0	HIS	-	expression tag	UNP A0A073FMV0

- Molecule 2 is a protein called Peptide 35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	6	Total	C	Cl	N	O	0	0	0
			54	36	2	6	10			
2	I	6	Total	C	Cl	N	O	0	0	0
			54	36	2	6	10			
2	J	6	Total	C	Cl	N	O	0	0	0
			54	36	2	6	10			
2	K	6	Total	C	Cl	N	O	0	0	0
			54	36	2	6	10			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		
4	C	1	Total	K	0	0
			1	1		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C<sub>10</sub>H<sub>22</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			16	10	6		
5	A	1	Total	C	O	0	0
			10	6	4		
5	B	1	Total	C	O	0	0
			11	7	4		
5	B	1	Total	C	O	0	0
			13	8	5		
5	B	1	Total	C	O	0	0
			16	10	6		
5	C	1	Total	C	O	0	0
			5	3	2		
5	C	1	Total	C	O	0	0
			11	7	4		
5	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		
6	B	1	Total	Cl	0	0
			1	1		
6	D	1	Total	Cl	0	0
			1	1		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			7	4	3		
7	D	1	Total	C	O	0	0
			7	4	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	296	Total	O	0	10
			306	306		
8	H	12	Total	O	0	0
			12	12		
8	B	203	Total	O	0	2
			205	205		
8	I	8	Total	O	0	0
			8	8		
8	C	249	Total	O	0	3
			252	252		
8	J	13	Total	O	0	0
			13	13		
8	D	208	Total	O	0	5
			213	213		
8	K	8	Total	O	0	1
			9	9		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta sliding clamp





- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp



- Molecule 2: Peptide 35



- Molecule 2: Peptide 35



- Molecule 2: Peptide 35



- Molecule 2: Peptide 35



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.81Å 135.83Å 87.54Å 90.00° 106.15° 90.00°	Depositor
Resolution (Å)	84.09 – 2.05 84.09 – 2.05	Depositor EDS
% Data completeness (in resolution range)	78.3 (84.09-2.05) 78.4 (84.09-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.210 , 0.241 0.210 , 0.240	Depositor DCC
$R_{free}$ test set	4327 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.9	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12789	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CL, ACE, ZCL, SOQ, 1PE, K, GOL, ALC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/2942	0.51	0/3983
1	B	0.27	0/2906	0.51	1/3933 (0.0%)
1	C	0.27	0/2907	0.51	0/3931
1	D	0.27	0/2908	0.51	0/3933
2	H	0.20	0/16	0.26	0/19
2	I	0.22	0/16	0.32	0/19
2	J	0.22	0/16	0.24	0/19
2	K	0.22	0/16	0.27	0/19
All	All	0.27	0/11727	0.51	1/15856 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1
2	I	0	2
2	J	0	2
2	K	0	2
All	All	0	7

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	98	LEU	CB-CG-CD1	-7.34	98.52	111.00

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	371	LEU	Mainchain
2	I	368	GLN	Mainchain
2	I	371	LEU	Mainchain
2	J	369	ALC	Mainchain
2	J	370	SOQ	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2887	0	2884	34	0
1	B	2844	0	2853	18	0
1	C	2851	0	2870	21	0
1	D	2850	0	2875	22	0
2	H	54	0	42	0	0
2	I	54	0	42	0	0
2	J	54	0	42	0	0
2	K	54	0	42	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	26	0	35	3	0
5	B	40	0	52	1	0
5	C	16	0	18	0	0
5	D	10	0	13	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	C	7	0	10	2	0
7	D	7	0	10	0	0
8	A	306	0	0	10	0
8	B	205	0	0	1	0
8	C	252	0	0	3	0
8	D	213	0	0	3	0
8	H	12	0	0	0	0
8	I	8	0	0	0	0
8	J	13	0	0	0	0
8	K	9	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	12789	0	11804	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ARG:NH1	8:A:501:HOH:O	2.00	0.94
1:B:50:GLU:OE1	8:B:501:HOH:O	1.97	0.82
1:B:184:ILE:HD11	1:B:188:LEU:HD11	1.64	0.80
1:C:276:GLU:OE1	8:C:601:HOH:O	2.00	0.78
1:A:301:GLU:OE1	8:A:502:HOH:O	2.07	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/386 (96%)	363 (98%)	6 (2%)	1 (0%)	41	31
1	B	366/386 (95%)	360 (98%)	6 (2%)	0	100	100
1	C	363/386 (94%)	357 (98%)	5 (1%)	1 (0%)	41	31
1	D	363/386 (94%)	354 (98%)	7 (2%)	2 (1%)	25	15
2	H	2/6 (33%)	2 (100%)	0	0	100	100
2	I	2/6 (33%)	2 (100%)	0	0	100	100
2	J	2/6 (33%)	2 (100%)	0	0	100	100
2	K	2/6 (33%)	2 (100%)	0	0	100	100
All	All	1470/1568 (94%)	1442 (98%)	24 (2%)	4 (0%)	41	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	GLY
1	C	39	ASP
1	D	208	ASP
1	D	210	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/330 (96%)	316 (100%)	1 (0%)	92	93
1	B	311/330 (94%)	311 (100%)	0	100	100
1	C	314/330 (95%)	312 (99%)	2 (1%)	86	87
1	D	314/330 (95%)	311 (99%)	3 (1%)	76	75
2	H	2/2 (100%)	2 (100%)	0	100	100
2	I	2/2 (100%)	2 (100%)	0	100	100
2	J	2/2 (100%)	2 (100%)	0	100	100
2	K	2/2 (100%)	2 (100%)	0	100	100
All	All	1264/1328 (95%)	1258 (100%)	6 (0%)	88	89

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	198	LYS
1	D	285	VAL
1	D	295	ASN
1	C	204	MET
1	A	15	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 5 are monoatomic - leaving 12 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	368/386 (95%)	0.62	16 (4%) 35 38	17, 27, 49, 108	0
1	B	365/386 (94%)	0.65	32 (8%) 10 10	18, 34, 65, 114	0
1	C	365/386 (94%)	0.67	25 (6%) 17 18	18, 32, 58, 127	0
1	D	364/386 (94%)	0.67	20 (5%) 25 27	20, 35, 64, 97	0
2	H	2/6 (33%)	0.21	0 100 100	29, 29, 29, 30	0
2	I	2/6 (33%)	0.76	0 100 100	27, 27, 27, 37	0
2	J	2/6 (33%)	-0.18	0 100 100	26, 26, 26, 29	0
2	K	2/6 (33%)	-0.25	0 100 100	26, 26, 26, 35	0
All	All	1470/1568 (93%)	0.65	93 (6%) 20 21	17, 32, 61, 127	0

The worst 5 of 93 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	209	GLY	12.6
1	A	119	LEU	9.7
1	D	21	LEU	5.6
1	C	211	ASP	4.7
1	D	73	ARG	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZCL	K	372	14/14	0.64	0.20	59,60,62,63	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZCL	I	372	14/14	0.67	0.21	59,62,64,64	0
2	ZCL	H	372	14/14	0.87	0.16	33,34,36,36	0
2	ALC	I	369	11/12	0.88	0.14	22,23,25,27	0
2	ZCL	J	372	14/14	0.89	0.13	34,35,37,37	0
2	ALC	K	369	11/12	0.89	0.14	27,28,28,29	0
2	ALC	H	369	11/12	0.92	0.13	26,27,29,29	0
2	ALC	J	369	11/12	0.94	0.13	25,25,27,28	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	PEG	D	401	7/7	0.44	0.28	64,64,65,65	0
3	GOL	B	401	6/6	0.46	0.23	66,66,67,67	0
5	1PE	B	403	13/16	0.58	0.33	81,82,86,87	0
5	1PE	D	402	10/16	0.59	0.27	60,61,62,63	0
5	1PE	A	403	16/16	0.67	0.26	71,72,74,74	0
5	1PE	C	503	5/16	0.68	0.35	67,67,69,70	0
5	1PE	C	504	11/16	0.68	0.20	63,65,65,66	0
5	1PE	B	402	11/16	0.70	0.25	70,71,72,74	0
5	1PE	B	404	16/16	0.74	0.26	75,77,80,80	0
5	1PE	A	404	10/16	0.76	0.17	58,59,59,61	0
7	PEG	C	501	7/7	0.77	0.16	63,63,63,64	0
3	GOL	A	401	6/6	0.85	0.26	67,67,67,67	6
6	CL	D	403	1/1	0.98	0.14	38,38,38,38	0
4	K	A	402	1/1	0.98	0.23	24,24,24,24	0
6	CL	A	405	1/1	0.98	0.16	30,30,30,30	0
6	CL	B	405	1/1	0.99	0.15	36,36,36,36	0
4	K	C	502	1/1	0.99	0.20	28,28,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.