



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 24, 2021 – 01:10 pm GMT

PDB ID : 7AZL
Title : DNA polymerase sliding clamp from Escherichia coli with peptide 38 bound
Authors : Monsarrat, C.; Compain, G.; Andre, C.; Martiel, I.; Engilberge, S.; Olieric, V.;
Wolff, P.; Brillet, K.; Landolfo, M.; Silva da Veiga, C.; Wagner, J.; Guichard,
G.; Burnouf, D.Y.
Deposited on : 2020-11-16
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4 (270009), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	<div> <div>2%</div> <div>90%</div> <div>6%</div> </div>
1	B	386	<div> <div>4%</div> <div>91%</div> <div>7%</div> </div>
1	C	386	<div> <div>3%</div> <div>91%</div> <div>6%</div> </div>
1	D	386	<div> <div>3%</div> <div>90%</div> <div>7%</div> </div>
2	E	6	<div> <div>83%</div> <div>17%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	6	 100%
2	G	6	 100%
2	H	6	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	1	0
			2808	1762	487	540	19			
1	B	358	Total	C	N	O	S	0	0	0
			2778	1747	484	528	19			
1	C	361	Total	C	N	O	S	0	2	0
			2798	1762	487	530	19			
1	D	359	Total	C	N	O	S	0	0	0
			2773	1750	482	522	19			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A073FMV0
A	-18	GLY	-	expression tag	UNP A0A073FMV0
A	-17	SER	-	expression tag	UNP A0A073FMV0
A	-16	SER	-	expression tag	UNP A0A073FMV0
A	-15	HIS	-	expression tag	UNP A0A073FMV0
A	-14	HIS	-	expression tag	UNP A0A073FMV0
A	-13	HIS	-	expression tag	UNP A0A073FMV0
A	-12	HIS	-	expression tag	UNP A0A073FMV0
A	-11	HIS	-	expression tag	UNP A0A073FMV0
A	-10	HIS	-	expression tag	UNP A0A073FMV0
A	-9	SER	-	expression tag	UNP A0A073FMV0
A	-8	SER	-	expression tag	UNP A0A073FMV0
A	-7	GLY	-	expression tag	UNP A0A073FMV0
A	-6	LEU	-	expression tag	UNP A0A073FMV0
A	-5	VAL	-	expression tag	UNP A0A073FMV0
A	-4	PRO	-	expression tag	UNP A0A073FMV0
A	-3	ARG	-	expression tag	UNP A0A073FMV0
A	-2	GLY	-	expression tag	UNP A0A073FMV0
A	-1	SER	-	expression tag	UNP A0A073FMV0
A	0	HIS	-	expression tag	UNP A0A073FMV0
B	-19	MET	-	initiating methionine	UNP A0A073FMV0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP A0A073FMV0
B	-17	SER	-	expression tag	UNP A0A073FMV0
B	-16	SER	-	expression tag	UNP A0A073FMV0
B	-15	HIS	-	expression tag	UNP A0A073FMV0
B	-14	HIS	-	expression tag	UNP A0A073FMV0
B	-13	HIS	-	expression tag	UNP A0A073FMV0
B	-12	HIS	-	expression tag	UNP A0A073FMV0
B	-11	HIS	-	expression tag	UNP A0A073FMV0
B	-10	HIS	-	expression tag	UNP A0A073FMV0
B	-9	SER	-	expression tag	UNP A0A073FMV0
B	-8	SER	-	expression tag	UNP A0A073FMV0
B	-7	GLY	-	expression tag	UNP A0A073FMV0
B	-6	LEU	-	expression tag	UNP A0A073FMV0
B	-5	VAL	-	expression tag	UNP A0A073FMV0
B	-4	PRO	-	expression tag	UNP A0A073FMV0
B	-3	ARG	-	expression tag	UNP A0A073FMV0
B	-2	GLY	-	expression tag	UNP A0A073FMV0
B	-1	SER	-	expression tag	UNP A0A073FMV0
B	0	HIS	-	expression tag	UNP A0A073FMV0
C	-19	MET	-	initiating methionine	UNP A0A073FMV0
C	-18	GLY	-	expression tag	UNP A0A073FMV0
C	-17	SER	-	expression tag	UNP A0A073FMV0
C	-16	SER	-	expression tag	UNP A0A073FMV0
C	-15	HIS	-	expression tag	UNP A0A073FMV0
C	-14	HIS	-	expression tag	UNP A0A073FMV0
C	-13	HIS	-	expression tag	UNP A0A073FMV0
C	-12	HIS	-	expression tag	UNP A0A073FMV0
C	-11	HIS	-	expression tag	UNP A0A073FMV0
C	-10	HIS	-	expression tag	UNP A0A073FMV0
C	-9	SER	-	expression tag	UNP A0A073FMV0
C	-8	SER	-	expression tag	UNP A0A073FMV0
C	-7	GLY	-	expression tag	UNP A0A073FMV0
C	-6	LEU	-	expression tag	UNP A0A073FMV0
C	-5	VAL	-	expression tag	UNP A0A073FMV0
C	-4	PRO	-	expression tag	UNP A0A073FMV0
C	-3	ARG	-	expression tag	UNP A0A073FMV0
C	-2	GLY	-	expression tag	UNP A0A073FMV0
C	-1	SER	-	expression tag	UNP A0A073FMV0
C	0	HIS	-	expression tag	UNP A0A073FMV0
D	-19	MET	-	initiating methionine	UNP A0A073FMV0
D	-18	GLY	-	expression tag	UNP A0A073FMV0
D	-17	SER	-	expression tag	UNP A0A073FMV0

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP A0A073FMV0
D	-15	HIS	-	expression tag	UNP A0A073FMV0
D	-14	HIS	-	expression tag	UNP A0A073FMV0
D	-13	HIS	-	expression tag	UNP A0A073FMV0
D	-12	HIS	-	expression tag	UNP A0A073FMV0
D	-11	HIS	-	expression tag	UNP A0A073FMV0
D	-10	HIS	-	expression tag	UNP A0A073FMV0
D	-9	SER	-	expression tag	UNP A0A073FMV0
D	-8	SER	-	expression tag	UNP A0A073FMV0
D	-7	GLY	-	expression tag	UNP A0A073FMV0
D	-6	LEU	-	expression tag	UNP A0A073FMV0
D	-5	VAL	-	expression tag	UNP A0A073FMV0
D	-4	PRO	-	expression tag	UNP A0A073FMV0
D	-3	ARG	-	expression tag	UNP A0A073FMV0
D	-2	GLY	-	expression tag	UNP A0A073FMV0
D	-1	SER	-	expression tag	UNP A0A073FMV0
D	0	HIS	-	expression tag	UNP A0A073FMV0

- Molecule 2 is a protein called Peptide 38.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	6	Total	C	Cl	N	O	0	0	0
			61	42	2	7	10			
2	G	6	Total	C	Cl	N	O	0	1	0
			70	47	2	8	13			
2	F	6	Total	C	Cl	N	O	0	0	0
			61	42	2	7	10			
2	H	6	Total	C	Cl	N	O	0	0	0
			61	42	2	7	10			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	4	3		
4	D	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	128	Total 129	O 129	0	1
5	B	98	Total 98	O 98	0	0
5	C	96	Total 96	O 96	0	0
5	D	95	Total 95	O 95	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp





- Molecule 2: Peptide 38

Chain E: 83% 17%



- Molecule 2: Peptide 38

Chain G: 100%

There are no outlier residues recorded for this chain.

- Molecule 2: Peptide 38

Chain F: 100%

There are no outlier residues recorded for this chain.

- Molecule 2: Peptide 38

Chain H: 100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.61Å 86.64Å 153.90Å 90.00° 92.56° 90.00°	Depositor
Resolution (Å)	70.69 – 2.42 153.74 – 2.42	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.69-2.42) 87.6 (153.74-2.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.43Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.208 , 0.233 0.212 , 0.241	Depositor DCC
R_{free} test set	3558 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.647	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.19$	Xtriage
Estimated twinning fraction	0.105 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11848	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALC, ZCL, SOQ, SJW, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2858	0.59	0/3871
1	B	0.37	0/2825	0.59	0/3823
1	C	0.36	0/2852	0.59	1/3862 (0.0%)
1	D	0.37	0/2821	0.59	0/3819
2	E	0.19	0/15	0.46	0/17
2	F	0.42	0/15	0.63	0/17
2	G	0.53	0/15	0.52	0/17
2	H	0.40	0/15	0.51	0/17
All	All	0.37	0/11416	0.59	1/15443 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	264	LYS	CG-CD-CE	6.28	130.75	111.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2808	0	2793	4	0
1	B	2778	0	2771	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2798	0	2792	2	0
1	D	2773	0	2781	5	0
2	E	61	0	39	1	0
2	F	61	0	39	0	0
2	G	70	0	38	0	0
2	H	61	0	39	0	0
3	A	6	0	8	0	0
4	B	7	0	10	0	0
4	D	7	0	10	1	0
5	A	129	0	0	0	0
5	B	98	0	0	0	0
5	C	96	0	0	0	0
5	D	95	0	0	1	0
All	All	11848	0	11320	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:184:ILE:HD11	1:D:188:LEU:HD21	1.93	0.51
1:A:184:ILE:HD11	1:A:188:LEU:HD21	1.93	0.50
1:C:184:ILE:HD11	1:C:188:LEU:HD21	1.93	0.48
1:D:136:LYS:HA	5:D:563:HOH:O	2.14	0.47
1:A:340:LEU:HD23	1:A:347:VAL:CG1	2.47	0.45
1:C:132:GLN:NE2	1:C:227:VAL:HG13	2.31	0.44
1:D:17:VAL:HG12	1:D:33:LEU:HD22	1.99	0.44
1:D:45:THR:HG22	1:D:54:VAL:HG22	2.00	0.44
1:D:58:ALA:HB2	4:D:401:PEG:H11	2.01	0.42
1:A:17:VAL:CG1	1:A:33:LEU:HD22	2.48	0.42
1:B:132:GLN:NE2	1:B:227:VAL:HG13	2.34	0.42
2:E:405:SOQ:O	2:E:405:SOQ:C1	2.64	0.42
1:A:245:ARG:HA	1:A:248:LEU:HD12	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/386 (93%)	352 (98%)	7 (2%)	1 (0%)	41	54
1	B	350/386 (91%)	341 (97%)	9 (3%)	0	100	100
1	C	357/386 (92%)	348 (98%)	9 (2%)	0	100	100
1	D	353/386 (92%)	345 (98%)	8 (2%)	0	100	100
2	E	2/6 (33%)	2 (100%)	0	0	100	100
2	F	2/6 (33%)	2 (100%)	0	0	100	100
2	G	2/6 (33%)	2 (100%)	0	0	100	100
2	H	2/6 (33%)	2 (100%)	0	0	100	100
All	All	1428/1568 (91%)	1394 (98%)	33 (2%)	1 (0%)	51	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/330 (93%)	303 (98%)	5 (2%)	62	78
1	B	305/330 (92%)	302 (99%)	3 (1%)	76	87
1	C	306/330 (93%)	301 (98%)	5 (2%)	62	78
1	D	303/330 (92%)	299 (99%)	4 (1%)	69	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	2/2 (100%)	2 (100%)	0	100	100
2	F	2/2 (100%)	2 (100%)	0	100	100
2	G	2/2 (100%)	2 (100%)	0	100	100
2	H	2/2 (100%)	2 (100%)	0	100	100
All	All	1230/1328 (93%)	1213 (99%)	17 (1%)	67	81

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	176	ARG
1	A	181	SER
1	A	214	LEU
1	A	362	MET
1	B	176	ARG
1	B	244	TYR
1	B	362	MET
1	C	16	GLN
1	C	176	ARG
1	C	181	SER
1	C	311	SER
1	C	362	MET
1	D	176	ARG
1	D	201	ILE
1	D	246	ARG
1	D	362	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	61	GLN
1	B	217	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/386 (94%)	0.01	6 (1%) 70 67	18, 37, 66, 92	0
1	B	358/386 (92%)	0.28	16 (4%) 33 31	29, 49, 83, 106	0
1	C	361/386 (93%)	0.26	11 (3%) 50 48	28, 50, 78, 103	0
1	D	359/386 (93%)	0.23	12 (3%) 46 44	26, 49, 80, 105	0
2	E	2/6 (33%)	0.28	0 100 100	38, 38, 38, 39	0
2	F	2/6 (33%)	0.18	0 100 100	36, 36, 36, 49	0
2	G	2/6 (33%)	-0.18	0 100 100	33, 33, 33, 46	0
2	H	2/6 (33%)	0.09	0 100 100	36, 36, 36, 40	0
All	All	1449/1568 (92%)	0.20	45 (3%) 49 46	18, 46, 78, 106	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	20	PRO	5.6
1	D	366	LEU	5.4
1	B	119	LEU	4.4
1	D	26	THR	4.3
1	C	185	GLY	4.2
1	B	187	SER	3.5
1	D	190	SER	3.4
1	B	120	ASP	3.2
1	B	26	THR	3.2
1	A	-1	SER	3.1
1	A	119	LEU	3.0
1	C	26	THR	3.0
1	D	227	VAL	2.9
1	B	213	PRO	2.9
1	B	183	PRO	2.9
1	A	208	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	27	LEU	2.7
1	D	20	PRO	2.7
1	B	214	LEU	2.7
1	C	312	GLY	2.6
1	C	248	LEU	2.6
1	C	132	GLN	2.6
1	D	118	ASN	2.6
1	B	353	ALA	2.5
1	A	251	ASN	2.5
1	B	149	GLN	2.4
1	B	153	TYR	2.4
1	C	65	PRO	2.4
1	B	117	PRO	2.4
1	B	126	VAL	2.4
1	C	352	ALA	2.4
1	B	247	VAL	2.3
1	B	27	LEU	2.3
1	C	247	VAL	2.3
1	D	119	LEU	2.3
1	D	114	ALA	2.3
1	D	37	VAL	2.2
1	B	238	ASP	2.2
1	D	238	ASP	2.1
1	D	38	ALA	2.1
1	C	148	HIS	2.1
1	A	19	GLY	2.1
1	C	311	SER	2.0
1	C	250	LYS	2.0
1	A	117	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZCL	F	406	14/14	0.74	0.24	56,73,85,86	0
2	ZCL	E	407	14/14	0.81	0.30	70,87,100,101	0
2	ZCL	G	407	14/14	0.83	0.20	47,74,82,98	0
2	ZCL	H	407	14/14	0.85	0.19	33,67,75,82	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ALC	F	403	11/12	0.88	0.20	38,42,54,55	0
2	ALC	H	404	11/12	0.91	0.15	36,42,49,62	0
2	ALC	E	404	11/12	0.93	0.17	32,37,44,50	0
2	ALC	G	404	11/12	0.96	0.15	36,49,52,66	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PEG	D	401	7/7	0.87	0.18	46,51,54,59	0
3	GOL	A	401	6/6	0.91	0.21	35,40,41,46	0
4	PEG	B	401	7/7	0.93	0.12	32,37,46,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.