



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 05:38 AM BST

PDB ID : 3B08
Title : Crystal structure of the mouse HOIL1-L-NZF in complex with linear di-ubiquitin
Authors : Sato, Y.; Fujita, H.; Yoshikawa, A.; Yamashita, M.; Yamagata, A.; Kaiser, S.E.; Iwai, K.; Fukai, S.
Deposited on : 2011-06-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	<div> <div>88%</div> <div>11% ..</div> </div>
1	D	152	<div> <div>80%</div> <div>17% ..</div> </div>
1	G	152	<div> <div>87%</div> <div>11% ..</div> </div>
1	J	152	<div> <div>%</div> <div>86%</div> <div>11% ..</div> </div>
2	B	64	<div> <div>5%</div> <div>78%</div> <div>13% 9%</div> </div>
2	E	64	<div> <div>2%</div> <div>83%</div> <div>9% 8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	64	<div><div></div><div>5%</div><div>78%</div><div>11%</div><div>•</div><div>8%</div></div>
2	K	64	<div><div></div><div>5%</div><div>83%</div><div>6%</div><div>•</div><div>8%</div></div>
3	C	2	<div><div></div><div>50%</div><div>50%</div></div>
3	F	2	<div><div></div><div>100%</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyubiquitin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1198	754	209	233	2			
1	D	151	Total	C	N	O	S	0	0	0
			1198	754	209	233	2			
1	G	150	Total	C	N	O	S	0	1	0
			1202	757	211	232	2			
1	J	149	Total	C	N	O	S	0	1	0
			1191	751	207	231	2			

- Molecule 2 is a protein called RanBP-type and C3HC4-type zinc finger-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	58	Total	C	N	O	S	0	0	0
			462	284	83	88	7			
2	E	59	Total	C	N	O	S	0	0	0
			471	289	85	90	7			
2	H	59	Total	C	N	O	S	0	0	0
			471	289	85	90	7			
2	K	59	Total	C	N	O	S	0	0	0
			472	290	86	89	7			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	187	GLY	-	expression tag	UNP Q9WUB0
B	188	PRO	-	expression tag	UNP Q9WUB0
B	189	GLY	-	expression tag	UNP Q9WUB0
B	190	HIS	-	expression tag	UNP Q9WUB0
B	191	MET	-	expression tag	UNP Q9WUB0
E	187	GLY	-	expression tag	UNP Q9WUB0
E	188	PRO	-	expression tag	UNP Q9WUB0
E	189	GLY	-	expression tag	UNP Q9WUB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	190	HIS	-	expression tag	UNP Q9WUB0
E	191	MET	-	expression tag	UNP Q9WUB0
H	187	GLY	-	expression tag	UNP Q9WUB0
H	188	PRO	-	expression tag	UNP Q9WUB0
H	189	GLY	-	expression tag	UNP Q9WUB0
H	190	HIS	-	expression tag	UNP Q9WUB0
H	191	MET	-	expression tag	UNP Q9WUB0
K	187	GLY	-	expression tag	UNP Q9WUB0
K	188	PRO	-	expression tag	UNP Q9WUB0
K	189	GLY	-	expression tag	UNP Q9WUB0
K	190	HIS	-	expression tag	UNP Q9WUB0
K	191	MET	-	expression tag	UNP Q9WUB0

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			23	12	11			
3	F	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		
4	K	1	Total	Zn	0	0
			1	1		
4	E	1	Total	Zn	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	155	Total	O	0	0
			155	155		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	59	Total 59	O 59	0	0
5	D	142	Total 142	O 142	0	0
5	E	66	Total 66	O 66	0	0
5	G	113	Total 113	O 113	0	0
5	H	60	Total 60	O 60	0	0
5	J	150	Total 150	O 150	0	0
5	K	85	Total 85	O 85	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyubiquitin-C

Chain A: 




- Molecule 1: Polyubiquitin-C

Chain D: 




- Molecule 1: Polyubiquitin-C

Chain G: 




- Molecule 1: Polyubiquitin-C

Chain J: 




- Molecule 2: RanBP-type and C3HC4-type zinc finger-containing protein 1


Chain B: 




- Molecule 2: RanBP-type and C3HC4-type zinc finger-containing protein 1



- Molecule 2: RanBP-type and C3HC4-type zinc finger-containing protein 1



- Molecule 2: RanBP-type and C3HC4-type zinc finger-containing protein 1



- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	104.94Å 104.94Å 170.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.93 – 1.70 46.93 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.5 (46.93-1.70) 97.5 (46.93-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.74 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0019, PHENIX 1.7.1_743	Depositor
R, R_{free}	0.173 , 0.211 0.173 , 0.211	Depositor DCC
R_{free} test set	5103 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7545	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 68.40 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.3943e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1211	0.54	0/1630
1	D	0.33	0/1211	0.53	0/1630
1	G	0.35	0/1218	0.53	0/1639
1	J	0.35	0/1207	0.52	0/1625
2	B	0.36	0/474	0.58	0/642
2	E	0.36	0/483	0.53	0/654
2	H	0.37	0/483	0.58	0/654
2	K	0.37	0/485	0.58	0/657
All	All	0.34	0/6772	0.54	0/9131

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1198	0	1253	12	0
1	D	1198	0	1253	19	0
1	G	1202	0	1263	9	0
1	J	1191	0	1250	14	0
2	B	462	0	430	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	471	0	438	7	0
2	H	471	0	438	10	0
2	K	472	0	437	7	0
3	C	23	0	21	0	0
3	F	23	0	21	0	0
4	B	1	0	0	0	0
4	E	1	0	0	0	0
4	H	1	0	0	0	0
4	K	1	0	0	0	0
5	A	155	0	0	1	0
5	B	59	0	0	1	0
5	D	142	0	0	2	0
5	E	66	0	0	3	0
5	G	113	0	0	0	0
5	H	60	0	0	2	0
5	J	150	0	0	0	0
5	K	85	0	0	0	0
All	All	7545	0	6804	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:206:PRO:O	2:E:218:ARG:NH2	1.93	1.00
1:D:118:ARG:HB2	1:D:146:VAL:HG22	1.46	0.95
2:H:208:ARG:NH1	5:H:610:HOH:O	1.99	0.94
1:A:118:ARG:NH1	2:H:242:GLU:OE2	2.06	0.88
1:D:63:LYS:HG2	1:D:64:GLU:HG2	1.54	0.87
2:K:191:MET:HG3	2:K:192:PRO:HD3	1.60	0.84
1:J:100:GLU:HG3	1:J:128:ASP:O	1.88	0.74
2:K:190:HIS:N	2:K:190:HIS:HD1	1.88	0.72
1:G:22:THR:OG1	1:G:24:GLU:HG3	1.90	0.72
1:D:118:ARG:HB2	1:D:146:VAL:CG2	2.19	0.71
2:K:190:HIS:ND1	2:K:190:HIS:N	2.38	0.71
1:J:83:THR:OG1	1:J:85:THR:HG22	1.92	0.70
1:A:115:ASP:OD2	5:A:184:HOH:O	2.09	0.69
1:A:82:LYS:HD3	1:A:144:HIS:CE1	2.27	0.69
2:E:247:GLN:HB2	2:K:248:TYR:HD2	1.57	0.68
1:A:100:GLU:HG2	1:A:128:ASP:HB3	1.75	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:112:ILE:HD13	1:D:147:LEU:HD11	1.79	0.65
2:B:230:PRO:O	5:B:739:HOH:O	2.14	0.65
2:E:241:GLU:OE1	5:E:388:HOH:O	2.14	0.64
2:B:245:LEU:HD23	2:H:249:GLN:HG2	1.81	0.62
1:J:118:ARG:CZ	1:J:148:ARG:HG3	2.31	0.60
2:E:226:ALA:HB1	1:J:116:GLN:NE2	2.18	0.59
1:D:82:LYS:HE2	5:E:305:HOH:O	2.04	0.58
1:D:120:ILE:HD13	1:D:125:GLN:HA	1.86	0.57
1:D:82:LYS:HD2	1:D:142:THR:HG21	1.85	0.57
2:K:191:MET:HG3	2:K:192:PRO:CD	2.34	0.56
1:A:45:PHE:HB2	1:A:67:LEU:HD22	1.87	0.56
1:G:112:ILE:HD13	1:G:147:LEU:HD11	1.88	0.56
1:G:23:ILE:HB	1:G:52:ASP:HA	1.88	0.55
2:E:247:GLN:HB2	2:K:248:TYR:CD2	2.41	0.55
1:A:121:PHE:HB2	1:A:143:LEU:HD22	1.87	0.55
1:J:100:GLU:HG2	1:J:128:ASP:HB3	1.88	0.55
1:A:82:LYS:HD3	1:A:144:HIS:HE1	1.71	0.54
1:D:7:THR:OG1	1:D:11:LYS:HG2	2.07	0.54
1:A:120:ILE:HD13	1:A:125:GLN:HA	1.89	0.53
1:G:45:PHE:HB2	1:G:67:LEU:HD22	1.89	0.53
2:H:249:GLN:HB3	5:H:512:HOH:O	2.09	0.53
1:J:118:ARG:HB2	1:J:146:VAL:CG2	2.38	0.53
1:D:6:LYS:HG3	1:D:66:THR:CG2	2.40	0.52
1:A:100:GLU:CG	1:A:128:ASP:HB3	2.39	0.52
2:E:242:GLU:OE2	1:J:148:ARG:NH1	2.43	0.51
1:J:45:PHE:HB2	1:J:67:LEU:HD22	1.93	0.51
1:J:39:ASP:OD2	1:J:74:ARG:HD3	2.11	0.50
2:H:239:ALA:O	2:H:243:GLU:HG3	2.12	0.50
1:J:36:ILE:HD13	1:J:71:LEU:HD21	1.95	0.49
1:A:118:ARG:NE	1:A:125:GLN:OE1	2.43	0.49
2:H:245:LEU:HD23	2:H:249:GLN:HB2	1.93	0.49
1:J:100:GLU:CG	1:J:128:ASP:HB3	2.43	0.48
2:H:191:MET:O	2:H:191:MET:HG3	2.13	0.48
1:J:115:ASP:N	1:J:115:ASP:OD1	2.34	0.48
1:J:118:ARG:HB2	1:J:146:VAL:HG23	1.95	0.47
2:B:195:TRP:CE3	2:B:219:PRO:HD3	2.49	0.47
1:G:124:LYS:HD2	1:G:124:LYS:HA	1.76	0.47
1:G:124:LYS:HD2	1:G:125:GLN:H	1.79	0.47
2:K:245:LEU:HD23	2:K:245:LEU:C	2.35	0.46
1:D:94:GLU:HG2	5:D:189:HOH:O	2.16	0.45
1:D:100:GLU:CG	1:D:128:ASP:HB3	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:244:ALA:O	2:H:249:GLN:HA	2.17	0.45
1:A:23:ILE:HB	1:A:52:ASP:HA	1.98	0.44
2:H:194:GLY:HA3	2:H:204:ASN:O	2.18	0.44
2:H:191:MET:O	2:H:191:MET:CG	2.66	0.43
2:E:218:ARG:HD3	5:E:124:HOH:O	2.18	0.43
1:G:87:LYS:HE2	1:G:89:ILE:HG22	2.01	0.42
1:G:121:PHE:HB2	1:G:143:LEU:HD22	2.01	0.42
1:D:45:PHE:HB2	1:D:67:LEU:HD22	2.02	0.42
1:D:148:ARG:HD3	1:D:148:ARG:HA	1.91	0.42
1:J:112:ILE:HG21	1:J:147:LEU:HD12	2.02	0.42
1:G:118:ARG:HB2	1:G:146:VAL:HG23	2.01	0.42
1:D:115:ASP:N	1:D:115:ASP:OD1	2.38	0.41
1:D:121:PHE:HB3	1:D:126:LEU:HD21	2.02	0.41
1:A:99:ILE:HB	1:A:128:ASP:HA	2.02	0.41
2:B:214:CYS:O	2:B:215:CYS:HB2	2.21	0.41
1:D:112:ILE:HD11	5:D:257:HOH:O	2.21	0.41
1:D:23:ILE:HB	1:D:52:ASP:HA	2.03	0.41
1:D:11:LYS:HE3	1:D:11:LYS:HB2	1.87	0.40
1:D:44:ILE:HD13	1:D:49:GLN:HA	2.03	0.40
2:B:218:ARG:HA	2:B:219:PRO:HD3	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/152 (98%)	148 (99%)	1 (1%)	0	100	100
1	D	149/152 (98%)	148 (99%)	1 (1%)	0	100	100
1	G	149/152 (98%)	149 (100%)	0	0	100	100
1	J	148/152 (97%)	148 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	56/64 (88%)	55 (98%)	1 (2%)	0	100	100
2	E	57/64 (89%)	57 (100%)	0	0	100	100
2	H	57/64 (89%)	57 (100%)	0	0	100	100
2	K	57/64 (89%)	57 (100%)	0	0	100	100
All	All	822/864 (95%)	819 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/136 (100%)	132 (97%)	4 (3%)	42	23
1	D	136/136 (100%)	131 (96%)	5 (4%)	34	15
1	G	137/136 (101%)	134 (98%)	3 (2%)	52	34
1	J	136/136 (100%)	134 (98%)	2 (2%)	65	51
2	B	49/53 (92%)	49 (100%)	0	100	100
2	E	50/53 (94%)	50 (100%)	0	100	100
2	H	50/53 (94%)	48 (96%)	2 (4%)	31	13
2	K	50/53 (94%)	47 (94%)	3 (6%)	19	6
All	All	744/756 (98%)	725 (97%)	19 (3%)	46	28

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LYS
1	A	74	ARG
1	A	115	ASP
1	A	127	GLU
1	D	6	LYS
1	D	87	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	94	GLU
1	D	115	ASP
1	D	127	GLU
1	G	24	GLU
1	G	115	ASP
1	G	134	ASP
2	H	191	MET
2	H	245	LEU
1	J	62	GLN
1	J	115	ASP
2	K	190	HIS
2	K	191	MET
2	K	247	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	144	HIS
1	G	25	ASN
1	J	116	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GLC	C	1	3	11,11,12	0.75	0	15,15,17	1.21	2 (13%)
3	GLC	C	2	3	12,12,12	0.81	0	17,17,17	0.81	0
3	GLC	F	1	3	11,11,12	0.71	0	15,15,17	1.10	1 (6%)
3	GLC	F	2	3	12,12,12	0.93	1 (8%)	17,17,17	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	C	1	3	-	0/2/19/22	0/1/1/1
3	GLC	C	2	3	-	0/2/22/22	0/1/1/1
3	GLC	F	1	3	-	0/2/19/22	0/1/1/1
3	GLC	F	2	3	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2	GLC	O5-C1	2.01	1.47	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	GLC	O5-C5-C6	2.77	111.54	107.20
3	F	1	GLC	O5-C5-C6	2.48	111.09	107.20
3	C	1	GLC	C1-O5-C5	2.03	114.94	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/152 (99%)	-0.18	0 100 100	13, 21, 38, 43	0
1	D	151/152 (99%)	-0.12	0 100 100	12, 21, 38, 51	0
1	G	150/152 (98%)	-0.11	0 100 100	12, 21, 37, 67	0
1	J	149/152 (98%)	-0.16	2 (1%) 77 81	12, 21, 38, 51	0
2	B	58/64 (90%)	0.41	3 (5%) 27 30	12, 21, 44, 54	0
2	E	59/64 (92%)	0.31	1 (1%) 70 74	11, 19, 45, 66	0
2	H	59/64 (92%)	0.29	3 (5%) 28 31	10, 19, 58, 62	0
2	K	59/64 (92%)	0.17	3 (5%) 28 31	9, 17, 58, 67	0
All	All	836/864 (96%)	-0.02	12 (1%) 75 79	9, 21, 39, 67	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	191	MET	4.4
2	K	190	HIS	4.4
2	K	248	TYR	3.7
2	H	248	TYR	3.3
2	K	191	MET	2.8
2	B	226	ALA	2.8
2	E	249	GLN	2.7
2	H	249	GLN	2.6
2	B	227	SER	2.5
1	J	149	LEU	2.3
2	B	248	TYR	2.0
1	J	115	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GLC	C	1	11/12	0.92	0.10	22,29,34,35	0
3	GLC	F	1	11/12	0.93	0.10	17,23,27,34	0
3	GLC	C	2	12/12	0.94	0.13	22,25,30,31	0
3	GLC	F	2	12/12	0.95	0.07	15,18,22,22	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ZN	K	1	1/1	1.00	0.03	21,21,21,21	0
4	ZN	E	1	1/1	1.00	0.03	23,23,23,23	0
4	ZN	H	1	1/1	1.00	0.03	21,21,21,21	0
4	ZN	B	1	1/1	1.00	0.03	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.