



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2021 – 04:53 PM EDT

PDB ID : 2B2D
Title : RNA stemloop operator from bacteriophage QBETA complexed with an N87S,E89K mutant MS2 capsid
Authors : Horn, W.T.; Tars, K.; Grahn, E.; Helgstrand, C.; Baron, A.J.; Lago, H.; Adams, C.J.; Peabody, D.S.; Phillips, S.E.V.; Stonehouse, N.J.; Liljas, L.; Stockley, P.G.
Deposited on : 2005-09-19
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RNA backbone	3102	1007 (3.16-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	R	20	
1	S	20	
2	A	129	
2	B	129	
2	C	129	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*AP*UP*GP*CP*AP*UP*GP*UP*CP*UP*AP*AP*GP*AP*CP*AP*GP*CP*AP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	9	Total	C	N	O	P	0	0	0
			191	86	35	61	9			
1	S	3	Total	C	N	O	P	0	0	0
			67	30	15	19	3			

- Molecule 2 is a protein called Coat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	129	Total	C	N	O	S	0	0	0
			962	606	165	187	4			
2	B	129	Total	C	N	O	S	0	0	0
			962	606	165	187	4			
2	C	129	Total	C	N	O	S	0	0	0
			962	606	165	187	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	SER	ASN	engineered mutation	UNP P03612
A	89	LYS	GLU	engineered mutation	UNP P03612
B	87	SER	ASN	engineered mutation	UNP P03612
B	89	LYS	GLU	engineered mutation	UNP P03612
C	87	SER	ASN	engineered mutation	UNP P03612
C	89	LYS	GLU	engineered mutation	UNP P03612

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	S	1	Total	O	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total 62	O 62	0	0
3	B	12	Total 12	O 12	0	0
3	C	44	Total 44	O 44	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-R(*AP*UP*GP*CP*AP*UP*GP*UP*CP*UP*AP*AP*GP*AP*CP*AP*GP*CP*AP*U)-3'

Chain R: 



- Molecule 1: 5'-R(*AP*UP*GP*CP*AP*UP*GP*UP*CP*UP*AP*AP*GP*AP*CP*AP*GP*CP*AP*U)-3'

Chain S: 




- Molecule 2: Coat protein

Chain A: 





- Molecule 2: Coat protein

Chain B: 



- Molecule 2: Coat protein

Chain C: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	287.80Å 287.80Å 652.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.21	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.197 , 0.203	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3263	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	R	0.71	1/212 (0.5%)	1.24	3/327 (0.9%)
1	S	0.44	0/75	1.86	5/115 (4.3%)
2	A	0.60	0/979	0.81	0/1333
2	B	0.39	0/979	0.67	0/1333
2	C	0.59	0/979	0.79	0/1333
All	All	0.55	1/3224 (0.0%)	0.85	8/4441 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	2	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	R	307	G	P-OP2	6.68	1.60	1.49

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	411	A	C2'-C3'-O3'	10.06	131.64	109.50
1	S	412	A	C2'-C3'-O3'	8.06	127.22	109.50
1	R	313	G	O4'-C1'-N9	7.26	114.01	108.20
1	R	307	G	OP1-P-OP2	-6.80	109.40	119.60
1	R	311	A	N9-C1'-C2'	6.15	121.99	114.00
1	S	412	A	C4'-C3'-O3'	5.65	124.31	113.00
1	S	412	A	C4'-C3'-C2'	5.60	108.20	102.60
1	S	413	G	C2'-C3'-O3'	5.46	122.44	113.70

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	S	411	A	C3'
1	S	412	A	C3'

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	191	0	96	8	0
1	S	67	0	34	1	0
2	A	962	0	970	26	0
2	B	962	0	970	29	0
2	C	962	0	970	12	0
3	A	62	0	0	1	0
3	B	12	0	0	0	0
3	C	44	0	0	1	0
3	S	1	0	0	0	0
All	All	3263	0	3040	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:20:VAL:HG13	2:C:32:TRP:HB3	1.35	1.07
1:R:310:U:H4'	1:R:311:A:OP2	1.58	0.99
2:A:20:VAL:HG13	2:A:32:TRP:HB3	1.49	0.93
2:A:129:TYR:H	2:B:3:ASN:HD21	1.17	0.90
2:B:20:VAL:HG22	2:B:32:TRP:HB3	1.55	0.89
2:A:97:THR:HG22	2:A:99:SER:H	1.41	0.85
2:A:129:TYR:H	2:B:3:ASN:ND2	1.83	0.77
2:B:3:ASN:ND2	2:B:3:ASN:H	1.88	0.71
2:A:117:PRO:HB3	2:B:3:ASN:O	1.91	0.69
2:A:3:ASN:O	2:B:117:PRO:HB3	1.93	0.69
2:A:112:LEU:HD12	2:B:112:LEU:HD12	1.75	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:313:G:O2'	1:R:314:A:O4'	2.13	0.67
2:A:79:VAL:HG22	2:A:80:ALA:H	1.60	0.66
2:C:56:ARG:HD2	3:C:137:HOH:O	1.95	0.65
1:R:310:U:C4'	1:R:311:A:OP2	2.40	0.62
2:C:20:VAL:CG1	2:C:32:TRP:HB3	2.22	0.62
1:R:308:U:OP1	2:B:57:LYS:HD2	1.99	0.61
2:A:97:THR:HG22	2:A:98:ASN:N	2.16	0.60
1:R:309:C:O2'	1:R:310:U:H2'	2.04	0.58
2:A:126:SER:O	2:B:58:TYR:OH	2.19	0.58
2:A:52:SER:OG	2:A:53:ALA:N	2.37	0.57
2:B:20:VAL:CG2	2:B:32:TRP:HB3	2.32	0.56
2:A:86:LEU:HD22	2:B:108:MET:HG2	1.88	0.56
2:A:79:VAL:HG22	2:A:80:ALA:N	2.21	0.55
2:B:23:SER:HB3	2:B:33:ILE:HG22	1.90	0.52
2:A:88:MET:HE1	2:B:88:MET:HG2	1.91	0.51
2:B:3:ASN:ND2	2:B:3:ASN:N	2.53	0.51
1:R:308:U:P	2:B:49:ARG:HH22	2.34	0.50
2:A:20:VAL:CG1	2:A:32:TRP:HB3	2.33	0.50
2:A:40:GLN:HB2	2:A:67:VAL:HG12	1.92	0.50
2:C:49:ARG:NH1	2:C:57:LYS:HE2	2.27	0.49
2:A:97:THR:CG2	2:A:98:ASN:N	2.75	0.48
2:A:97:THR:HG22	2:A:99:SER:N	2.20	0.48
2:B:105:VAL:O	2:B:109:GLN:HG3	2.13	0.48
2:C:15:THR:O	2:C:17:ASP:N	2.47	0.48
2:A:129:TYR:N	2:B:3:ASN:HD21	1.99	0.47
1:S:411:A:OP1	1:S:411:A:H8	1.97	0.47
1:R:313:G:O2'	1:R:314:A:O5'	2.33	0.46
2:B:62:VAL:HB	2:B:86:LEU:HB3	1.97	0.46
2:B:3:ASN:N	2:B:3:ASN:HD22	2.13	0.46
2:C:15:THR:C	2:C:17:ASP:H	2.19	0.45
2:C:49:ARG:HH12	2:C:57:LYS:HE2	1.81	0.45
3:A:135:HOH:O	2:B:125:ASN:HA	2.16	0.45
2:C:20:VAL:HG13	2:C:32:TRP:CB	2.26	0.45
2:B:10:VAL:HB	2:B:18:VAL:HB	1.99	0.45
1:R:310:U:H6	1:R:310:U:O5'	2.01	0.43
2:B:67:VAL:O	2:B:70:GLN:HB3	2.18	0.43
2:B:36:ASN:HB2	2:B:41:ALA:HB2	2.01	0.43
2:B:26:ALA:O	2:B:27:ASN:HB2	2.19	0.42
2:A:60:ILE:HB	2:A:88:MET:HG3	2.00	0.42
2:A:86:LEU:HD21	2:A:88:MET:HE3	2.00	0.42
2:A:103:LEU:HA	2:A:103:LEU:HD12	1.76	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:108:MET:HG2	2:B:86:LEU:HD22	2.01	0.42
2:C:36:ASN:O	2:C:37:SER:C	2.58	0.42
2:C:49:ARG:HH22	2:C:57:LYS:NZ	2.17	0.42
2:A:88:MET:HE1	2:B:88:MET:CG	2.50	0.41
2:A:8:VAL:HA	2:A:19:THR:HA	2.03	0.41
2:C:66:LYS:HG2	2:C:82:TRP:CE2	2.56	0.41
2:A:21:ALA:O	2:A:32:TRP:HA	2.20	0.41
2:B:32:TRP:HB2	2:B:44:VAL:HB	2.02	0.41
2:C:92:ILE:HA	2:C:93:PRO:HD3	1.93	0.41
2:B:70:GLN:HG2	2:B:71:THR:N	2.36	0.40
2:B:75:VAL:HG12	2:B:77:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	127/129 (98%)	118 (93%)	6 (5%)	3 (2%)	6	22
2	B	127/129 (98%)	122 (96%)	4 (3%)	1 (1%)	19	51
2	C	127/129 (98%)	117 (92%)	9 (7%)	1 (1%)	19	51
All	All	381/387 (98%)	357 (94%)	19 (5%)	5 (1%)	12	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	2	SER
2	A	2	SER
2	A	14	GLY
2	C	16	GLY
2	A	13	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	106/106 (100%)	96 (91%)	10 (9%)	8	26
2	B	106/106 (100%)	100 (94%)	6 (6%)	20	51
2	C	106/106 (100%)	98 (92%)	8 (8%)	13	37
All	All	318/318 (100%)	294 (92%)	24 (8%)	13	37

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	4	PHE
2	A	8	VAL
2	A	19	THR
2	A	20	VAL
2	A	25	PHE
2	A	38	ARG
2	A	48	VAL
2	A	52	SER
2	A	65	PRO
2	A	114	ASP
2	B	3	ASN
2	B	15	THR
2	B	38	ARG
2	B	50	GLN
2	B	54	GLN
2	B	114	ASP
2	C	20	VAL
2	C	23	SER
2	C	37	SER
2	C	38	ARG
2	C	48	VAL
2	C	54	GLN
2	C	65	PRO
2	C	99	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
2	A	40	GLN
2	B	3	ASN
2	B	6	GLN
2	B	36	ASN
2	B	54	GLN
2	C	6	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	8/20 (40%)	2 (25%)	2 (25%)
1	S	3/20 (15%)	2 (66%)	2 (66%)
All	All	11/40 (27%)	4 (36%)	4 (36%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	R	311	A
1	R	314	A
1	S	412	A
1	S	413	G

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	R	310	U
1	R	313	G
1	S	411	A
1	S	412	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.