



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 09:47 am BST

PDB ID : 6B2N
Title : Crystal structure of TEM-1 beta-lactamase mutant M182N
Authors : Jimah, J.R.; Tolia, N.H.
Deposited on : 2017-09-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>
1	B	263	<div> <div>7%</div> <div> <div></div> <div>88%</div> <div>9%</div> </div> </div>
1	C	263	<div> <div>9%</div> <div> <div></div> <div>91%</div> <div>6%</div> </div> </div>
1	D	263	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16199 atoms, of which 7992 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase TEM.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	260	Total	C	H	N	O	S	0	0	0
			4017	1255	2008	357	387	10			
1	B	255	Total	C	H	N	O	S	0	0	0
			3959	1237	1981	351	380	10			
1	C	257	Total	C	H	N	O	S	0	0	0
			3981	1243	1988	355	385	10			
1	D	261	Total	C	H	N	O	S	0	0	0
			4033	1259	2015	359	390	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	ASN	MET	engineered mutation	UNP P62593
B	182	ASN	MET	engineered mutation	UNP P62593
C	182	ASN	MET	engineered mutation	UNP P62593
D	182	ASN	MET	engineered mutation	UNP P62593

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		
3	B	43	Total	O	0	0
			43	43		
3	C	54	Total	O	0	0
			54	54		
3	D	48	Total	O	0	0
			48	48		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-lactamase TEM




• Molecule 1: Beta-lactamase TEM



• Molecule 1: Beta-lactamase TEM



• Molecule 1: Beta-lactamase TEM



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.91Å 49.66Å 122.16Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	19.77 – 2.00 46.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.7 (19.77-2.00) 97.8 (46.00-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.231 , 0.283 0.239 , 0.292	Depositor DCC
R_{free} test set	2003 reflections (2.82%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.601	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.408 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	16199	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.9409e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/2040	0.46	0/2758
1	B	0.25	0/2005	0.45	0/2704
1	C	0.25	0/2023	0.45	0/2734
1	D	0.26	0/2050	0.46	0/2773
All	All	0.25	0/8118	0.46	0/10969

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2009	2008	2014	11	1
1	B	1978	1981	1987	13	0
1	C	1993	1988	1994	10	0
1	D	2018	2015	2021	15	1
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	1	0
3	A	44	0	0	0	0
3	B	43	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	54	0	0	1	0
3	D	48	0	0	1	0
All	All	8207	7992	8016	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:SER:OG	1:B:133:THR:OG1	1.93	0.84
1:D:219:PRO:O	1:D:223:SER:OG	2.02	0.73
1:D:35:ASP:OD1	1:D:39:GLN:NE2	2.22	0.73
1:C:239:GLU:O	1:C:273:ARG:NH2	2.22	0.72
1:D:31:VAL:O	1:D:35:ASP:N	2.27	0.67
1:C:63:GLU:O	3:C:401:HOH:O	2.14	0.65
1:D:276:GLN:O	1:D:279:GLU:N	2.32	0.62
1:C:157:ASP:OD2	1:C:189:THR:OG1	2.18	0.61
1:D:166:GLU:OE1	1:D:170:ASN:ND2	2.34	0.61
1:A:50:ASP:OD2	1:A:53:SER:OG	2.16	0.60
1:A:26:HIS:N	1:A:58:GLU:OE2	2.38	0.57
1:D:157:ASP:OD2	1:D:189:THR:OG1	2.21	0.56
1:B:242:SER:OG	1:B:273:ARG:NH1	2.37	0.54
1:B:182:ASN:OD1	1:B:185:ALA:N	2.34	0.53
1:D:240:ARG:NH2	2:D:301:SO4:O4	2.43	0.52
1:B:269:THR:O	1:B:273:ARG:NH2	2.44	0.51
1:C:39:GLN:OE1	1:C:276:GLN:NE2	2.44	0.50
1:C:257:ARG:NH2	1:C:288:TRP:O	2.42	0.49
1:B:164:ARG:NH2	1:B:179:ASP:OD2	2.41	0.48
1:B:192:LYS:O	1:B:196:GLY:N	2.38	0.48
1:D:39:GLN:OE1	1:D:276:GLN:NE2	2.46	0.48
1:B:234:LYS:HG3	1:B:234:LYS:O	2.13	0.48
1:A:69:MET:HE1	1:A:243:ARG:HB2	1.95	0.47
1:D:175:ASN:N	3:D:404:HOH:O	2.46	0.47
1:D:192:LYS:O	1:D:196:GLY:N	2.45	0.46
1:C:270:MET:O	1:C:274:ASN:N	2.37	0.46
1:A:32:LYS:HE2	1:A:279:GLU:HB3	1.98	0.46
1:A:177:GLU:O	1:A:180:THR:OG1	2.24	0.45
1:B:181:THR:HG21	1:B:186:MET:HE2	1.98	0.45
1:C:164:ARG:NE	1:C:179:ASP:OD2	2.43	0.45
1:B:205:GLN:NE2	1:B:209:ASP:OD1	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ASP:OD2	1:A:189:THR:OG1	2.34	0.44
1:B:77:CYS:O	1:B:81:LEU:HG	2.18	0.44
1:D:276:GLN:O	1:D:280:ILE:N	2.40	0.44
1:B:93:ARG:O	1:B:118:THR:HA	2.18	0.44
1:A:49:LEU:HD21	1:A:191:ARG:HD3	1.99	0.43
1:A:199:LEU:O	1:A:204:ARG:NE	2.39	0.43
1:B:234:LYS:O	1:B:234:LYS:CG	2.67	0.43
1:A:73:LYS:NZ	1:A:130:SER:OG	2.45	0.42
1:D:218:GLY:N	1:D:219:PRO:HD2	2.34	0.42
1:A:65:ARG:HA	1:A:181:THR:O	2.20	0.42
1:B:173:ILE:HB	1:B:176:ASP:HB2	2.01	0.41
1:A:106:SER:HB3	1:A:109:THR:OG1	2.20	0.41
1:D:30:LEU:O	1:D:34:LYS:N	2.41	0.41
1:C:216:VAL:CG2	1:C:237:ALA:HA	2.50	0.41
1:C:257:ARG:NH1	1:C:288:TRP:O	2.53	0.41
1:D:233:ASP:OD1	1:D:233:ASP:N	2.54	0.41
1:D:65:ARG:HA	1:D:181:THR:O	2.21	0.40
1:C:148:LEU:HD23	1:C:162:LEU:HD22	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:LYS:O	1:D:275:ARG:NH1[2_554]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/263 (96%)	242 (96%)	10 (4%)	0	100	100
1	B	241/263 (92%)	232 (96%)	9 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	249/263 (95%)	244 (98%)	5 (2%)	0	100	100
1	D	255/263 (97%)	244 (96%)	11 (4%)	0	100	100
All	All	997/1052 (95%)	962 (96%)	35 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/217 (99%)	215 (100%)	0	100	100
1	B	213/217 (98%)	212 (100%)	1 (0%)	88	92
1	C	214/217 (99%)	214 (100%)	0	100	100
1	D	217/217 (100%)	217 (100%)	0	100	100
All	All	859/868 (99%)	858 (100%)	1 (0%)	93	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	226	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	26	HIS
1	C	287	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	C	301	-	4,4,4	0.15	0	6,6,6	0.14	0
2	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.09	0
2	SO4	D	302	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	D	301	-	4,4,4	0.14	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/263 (98%)	0.28	11 (4%) 36 35	11, 30, 60, 116	0
1	B	255/263 (96%)	0.48	19 (7%) 14 13	16, 33, 77, 137	0
1	C	257/263 (97%)	0.57	24 (9%) 8 8	13, 35, 85, 161	0
1	D	261/263 (99%)	0.31	10 (3%) 40 39	10, 31, 66, 100	0
All	All	1033/1052 (98%)	0.41	64 (6%) 20 19	10, 33, 74, 161	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	221	LEU	5.1
1	C	70	SER	4.8
1	B	51	LEU	4.6
1	C	71	THR	4.6
1	B	57	LEU	4.4
1	C	224	ALA	4.2
1	D	176	ASP	4.1
1	C	33	VAL	4.0
1	C	42	ALA	3.8
1	C	227	ALA	3.8
1	B	224	ALA	3.7
1	C	264	THR	3.7
1	C	98	SER	3.6
1	C	232	ALA	3.6
1	C	128	THR	3.4
1	B	252	ASP	3.4
1	C	276	GLN	3.4
1	B	27	PRO	3.3
1	B	213	ALA	3.3
1	B	256	SER	3.2
1	C	72	PHE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	271	ASP	3.0
1	B	223	SER	2.9
1	B	40	LEU	2.9
1	C	175	ASN	2.9
1	D	175	ASN	2.9
1	C	251	PRO	2.9
1	A	30	LEU	2.9
1	D	202	ALA	2.8
1	D	190	LEU	2.8
1	A	29	THR	2.7
1	C	48	GLU	2.7
1	B	288	TRP	2.6
1	B	227	ALA	2.6
1	D	227	ALA	2.6
1	D	279	GLU	2.6
1	C	87	GLY	2.5
1	C	193	LEU	2.5
1	C	36	ALA	2.5
1	C	57	LEU	2.5
1	A	267	GLN	2.5
1	B	42	ALA	2.4
1	A	51	LEU	2.3
1	A	279	GLU	2.3
1	B	55	LYS	2.3
1	C	222	ARG	2.3
1	B	277	ILE	2.3
1	C	287	HIS	2.3
1	D	26	HIS	2.3
1	B	99	GLN	2.3
1	D	220	LEU	2.2
1	C	274	ASN	2.2
1	A	219	PRO	2.2
1	B	243	ARG	2.2
1	A	173	ILE	2.1
1	D	31	VAL	2.1
1	B	272	GLU	2.1
1	D	282	ALA	2.1
1	A	50	ASP	2.1
1	B	257	ARG	2.1
1	A	67	PRO	2.0
1	A	254	LYS	2.0
1	B	60	PHE	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	D	302	5/5	0.83	0.45	172,173,174,177	0
2	SO4	B	301	5/5	0.93	0.16	35,43,48,53	0
2	SO4	C	301	5/5	0.97	0.07	19,20,30,33	0
2	SO4	D	301	5/5	0.98	0.09	23,28,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.