



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 01:44 AM BST

PDB ID : 2B39
Title : Structure of mammalian C3 with an intact thioester at 3Å resolution
Authors : Fredslund, F.; Jenner, L.; Husted, L.B.; Nyborg, J.; Andersen, G.R.; Sottrup-Jensen, L.
Deposited on : 2005-09-20
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1661	
1	B	1661	
2	C	3	
2	D	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	1	X	-	-	-
2	BMA	C	3	-	-	-	X
2	NAG	D	1	X	-	-	-

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 25624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1610	Total	C	N	O	S	0	0	0
			12773	8093	2187	2439	54			
1	B	1610	Total	C	N	O	S	0	0	0
			12773	8093	2187	2439	54			

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	D	3	Total	C	N	O	0	0	0
			39	22	2	15			

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: C3





T927	K860	L794	Q725	Q659	R528	P459	Q396	A268	T188	S126	A64
L928	V861	K795	C726	C660	L529	R460	S397	F269	L190	G127	K65
K929	R862	L796	C726	C660	V530	V461	L398	V270	S191	L128	R66
V930	V863	S797	I729	Q662	A531	E462	T399	L271	V192	F130	V68
V931	E864	T799	Q730	ALA	Y533	K463	Q400	F272	L197	Q132	L69
P932	L865	T800	T731	THR	K605	K464	D401	G273	V198	Q132	
E933	L866	T801	L732	ARG	K606	P465	D402	V274	N199	D134	
	V867	W801	Q733	ARG	L536	G466	Q403	D275	K200	D134	
T940		E802	Q734	ARG	L537	T468	V404	D276	Q200	D134	
T945	A870	L804	Q735	ARG	A538	L469	A405	G277	Q203	D134	
L946	F871	L804	Q735	ARG	A538	L469	A405	G277	Q203	D134	
	C872	W806	H736	ALA	Q541	W471	L407	R280	K204	D134	
		S807	A744	S670	Q541	W471	L407	R280	K204	D134	
		L808	L742	Q672	R542	M410	I409	I281	K205	D134	
			GLU	Q672	R542	M410	I409	I281	K205	D134	
			LEU	Q672	R542	M410	I409	I281	K205	D134	
			ALA	Q672	R542	M410	I409	I281	K205	D134	
			ARG	Q672	R542	M410	I409	I281	K205	D134	
			ASP	Q672	R542	M410	I409	I281	K205	D134	
			LEU	Q672	R542	M410	I409	I281	K205	D134	
			D750	Q683	V553	K483	V422	D294	K224	D134	
			D751	Q683	V553	K483	V422	D294	K224	D134	
			D752	Q683	V553	K483	V422	D294	K224	D134	
			D753	Q683	V553	K483	V422	D294	K224	D134	
			D754	Q683	V553	K483	V422	D294	K224	D134	
			D755	Q683	V553	K483	V422	D294	K224	D134	
			D756	Q683	V553	K483	V422	D294	K224	D134	
			D757	Q683	V553	K483	V422	D294	K224	D134	
			D758	Q683	V553	K483	V422	D294	K224	D134	
			D759	Q683	V553	K483	V422	D294	K224	D134	
			D760	Q683	V553	K483	V422	D294	K224	D134	
			D761	Q683	V553	K483	V422	D294	K224	D134	
			D762	Q683	V553	K483	V422	D294	K224	D134	
			D763	Q683	V553	K483	V422	D294	K224	D134	
			D764	Q683	V553	K483	V422	D294	K224	D134	
			D765	Q683	V553	K483	V422	D294	K224	D134	
			D766	Q683	V553	K483	V422	D294	K224	D134	
			D767	Q683	V553	K483	V422	D294	K224	D134	
			D768	Q683	V553	K483	V422	D294	K224	D134	
			D769	Q683	V553	K483	V422	D294	K224	D134	
			D770	Q683	V553	K483	V422	D294	K224	D134	
			D771	Q683	V553	K483	V422	D294	K224	D134	
			D772	Q683	V553	K483	V422	D294	K224	D134	
			D773	Q683	V553	K483	V422	D294	K224	D134	
			D774	Q683	V553	K483	V422	D294	K224	D134	
			D775	Q683	V553	K483	V422	D294	K224	D134	
			D776	Q683	V553	K483	V422	D294	K224	D134	
			D777	Q683	V553	K483	V422	D294	K224	D134	
			D778	Q683	V553	K483	V422	D294	K224	D134	
			D779	Q683	V553	K483	V422	D294	K224	D134	
			D780	Q683	V553	K483	V422	D294	K224	D134	
			D781	Q683	V553	K483	V422	D294	K224	D134	
			D782	Q683	V553	K483	V422	D294	K224	D134	
			D783	Q683	V553	K483	V422	D294	K224	D134	
			D784	Q683	V553	K483	V422	D294	K224	D134	
			D785	Q683	V553	K483	V422	D294	K224	D134	
			D786	Q683	V553	K483	V422	D294	K224	D134	
			D787	Q683	V553	K483	V422	D294	K224	D134	
			D788	Q683	V553	K483	V422	D294	K224	D134	
			D789	Q683	V553	K483	V422	D294	K224	D134	
			D790	Q683	V553	K483	V422	D294	K224	D134	
			D791	Q683	V553	K483	V422	D294	K224	D134	
			D792	Q683	V553	K483	V422	D294	K224	D134	
			D793	Q683	V553	K483	V422	D294	K224	D134	
			D794	Q683	V553	K483	V422	D294	K224	D134	
			D795	Q683	V553	K483	V422	D294	K224	D134	
			D796	Q683	V553	K483	V422	D294	K224	D134	
			D797	Q683	V553	K483	V422	D294	K224	D134	
			D798	Q683	V553	K483	V422	D294	K224	D134	
			D799	Q683	V553	K483	V422	D294	K224	D134	
			D800	Q683	V553	K483	V422	D294	K224	D134	
			D801	Q683	V553	K483	V422	D294	K224	D134	
			D802	Q683	V553	K483	V422	D294	K224	D134	
			D803	Q683	V553	K483	V422	D294	K224	D134	
			D804	Q683	V553	K483	V422	D294	K224	D134	
			D805	Q683	V553	K483	V422	D294	K224	D134	
			D806	Q683	V553	K483	V422	D294	K224	D134	
			D807	Q683	V553	K483	V422	D294	K224	D134	
			D808	Q683	V553	K483	V422	D294	K224	D134	
			D809	Q683	V553	K483	V422	D294	K224	D134	
			D810	Q683	V553	K483	V422	D294	K224	D134	
			D811	Q683	V553	K483	V422	D294	K224	D134	
			D812	Q683	V553	K483	V422	D294	K224	D134	
			D813	Q683	V553	K483	V422	D294	K224	D134	
			D814	Q683	V553	K483	V422	D294	K224	D134	
			D815	Q683	V553	K483	V422	D294	K224	D134	
			D816	Q683	V553	K483	V422	D294	K224	D134	
			D817	Q683	V553	K483	V422	D294	K224	D134	
			D818	Q683	V553	K483	V422	D294	K224	D134	
			D819	Q683	V553	K483	V422	D294	K224	D134	
			D820	Q683	V553	K483	V422	D294	K224	D134	
			D821	Q683	V553	K483	V422	D294	K224	D134	
			D822	Q683	V553	K483	V422	D294	K224	D134	
			D823	Q683	V553	K483	V422	D294	K224	D134	
			D824	Q683	V553	K483	V422	D294	K224	D134	
			D825	Q683	V553	K483	V422	D294	K224	D134	
			D826	Q683	V553	K483	V422	D294	K224	D134	
			D827	Q683	V553	K483	V422	D294	K224	D134	
			D828	Q683	V553	K483	V422	D294	K224	D134	
			D829	Q683	V553	K483	V422	D294	K224	D134	
			D830	Q683	V553	K483	V422	D294	K224	D134	
			D831	Q683	V553	K483	V422	D294	K224	D134	
			D832	Q683	V553	K483	V422	D294	K224	D134	
			D833	Q683	V553	K483	V422	D294	K224	D134	
			D834	Q683	V553	K483	V422	D294	K224	D134	
			D835	Q683	V553	K483	V422	D294	K224	D134	
			D836	Q683	V553	K483	V422	D294	K224	D134	
			D837	Q683	V553	K483	V422	D294	K224	D134	
			D838	Q683	V553	K483	V422	D294	K224	D134	
			D839	Q683	V553	K483	V422	D294	K224	D134	
			D840	Q683	V553	K483	V422	D294	K224	D134	
			D841	Q683	V553	K483	V422	D294	K224	D134	
			D842	Q683	V553	K483	V422	D294	K224	D134	
			D843	Q683	V553	K483	V422	D294	K224	D134	
			D844	Q683	V553	K483	V422	D294	K224	D134	
			D845	Q683	V553	K483	V422	D294	K224	D134	
			D846	Q683	V553	K483	V422	D294	K224	D134	
			D847	Q683	V553	K483	V422	D294	K224	D134	
			D848	Q683	V553	K483	V422	D294	K224	D134	
			D849	Q683	V553	K483	V422	D294	K224	D134	
			D850	Q683	V553	K483	V422	D294	K224	D134	
			D851	Q683	V553	K483	V422	D294	K224	D134	
			D852	Q683	V553	K483	V422	D294	K224	D134	
			D853	Q683	V553	K483	V422	D294	K224	D134	
			D854	Q683	V553	K483	V422	D294	K224	D134	
			D855	Q683	V553	K483	V422	D294	K224	D134	
			D856	Q683	V553	K483	V422	D294	K224	D134	
			D857	Q683	V553	K483	V422	D294	K224	D134	
			D858	Q683	V553	K483	V422	D294	K224	D134	
			D859	Q683	V553	K483	V422	D294	K224	D134	
			D860	Q683	V553	K483	V422	D294	K224	D134	
			D861	Q683	V553	K483	V422	D294	K224	D134	
			D862	Q683	V553	K483	V422	D294	K224	D134	
			D863	Q683	V553	K483	V422	D294	K224	D134	
			D864	Q683	V553	K483	V422	D294	K224	D134	
			D865	Q683	V553	K483	V422	D294	K224	D134	
			D866	Q683	V553	K483	V422	D294	K224	D134	
			D867	Q683	V553	K483	V422	D294	K224	D134	
			D868	Q683	V553	K483	V422	D294	K224	D134	
			D869	Q683	V553	K483	V422	D294	K224	D134	
			D870	Q683	V553	K483	V422	D294	K224	D134	
			D871	Q683	V553	K483	V422	D294	K224	D134	
			D872	Q683	V553	K483					



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 33% 67%

MAG1
MAG2
EWA3

• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 67% 33%

MAG1
MAG2
EWA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	254.25Å 246.86Å 113.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 37.85 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.00) 97.2 (37.85-3.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 3.01Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.278 , 0.286 0.273 , 0.285	Depositor DCC
R_{free} test set	1392 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	69.3	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 75.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	25624	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	1/13020 (0.0%)	0.82	15/17632 (0.1%)
1	B	0.57	0/13020	0.81	11/17632 (0.1%)
All	All	0.57	1/26040 (0.0%)	0.82	26/35264 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	102	HIS	CB-CG	6.64	1.61	1.50

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	LEU	CA-CB-CG	-9.75	92.87	115.30
1	A	91	PRO	N-CA-C	7.91	132.66	112.10
1	B	1362	LEU	CA-CB-CG	6.57	130.40	115.30
1	A	1362	LEU	CA-CB-CG	6.54	130.34	115.30
1	A	1130	GLY	N-CA-C	-6.49	96.87	113.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	138	TYR	Sidechain
1	B	820	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	12773	0	12782	1027	0
1	B	12773	0	12782	1002	0
2	C	39	0	34	3	0
2	D	39	0	34	3	0
All	All	25624	0	25632	2029	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 40.

The worst 5 of 2029 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:GLU:O	1:B:118:LYS:HG2	1.47	1.14
1:B:369:PHE:HB3	1:B:409:ILE:HD12	1.32	1.10
1:A:44:LEU:HD13	1:A:55:VAL:HG11	1.34	1.09
1:B:244:ILE:HD11	1:B:319:VAL:CG2	1.84	1.08
1:B:116:VAL:HG13	1:B:645:LYS:HG2	1.28	1.06

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1602/1661 (96%)	1285 (80%)	213 (13%)	104 (6%)	1	7
1	B	1602/1661 (96%)	1268 (79%)	239 (15%)	95 (6%)	1	9
All	All	3204/3322 (96%)	2553 (80%)	452 (14%)	199 (6%)	1	8

5 of 199 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	PRO
1	A	91	PRO
1	A	187	ILE
1	A	212	SER
1	A	244	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1420/1465 (97%)	1238 (87%)	182 (13%)	4	19
1	B	1420/1465 (97%)	1234 (87%)	186 (13%)	4	18
All	All	2840/2930 (97%)	2472 (87%)	368 (13%)	4	19

5 of 368 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1447	ASP
1	B	151	VAL
1	B	1351	LEU
1	A	1492	HIS
1	B	54	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 85 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	71	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	296	ASN
1	B	1298	GLN
1	B	79	ASN
1	B	184	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.57	0	17,19,21	1.00	1 (5%)
2	NAG	C	2	2	14,14,15	0.62	0	17,19,21	0.95	1 (5%)
2	BMA	C	3	2	11,11,12	0.47	0	15,15,17	0.55	0
2	NAG	D	1	1,2	14,14,15	0.57	0	17,19,21	1.00	0
2	NAG	D	2	2	14,14,15	0.61	0	17,19,21	1.03	2 (11%)
2	BMA	D	3	2	11,11,12	0.47	0	15,15,17	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	1/1/5/7	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	1/1/5/7	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	O5-C1-C2	-2.37	107.54	111.29
2	C	2	NAG	C3-C4-C5	-2.34	106.06	110.24
2	D	2	NAG	O5-C1-C2	-2.06	108.04	111.29
2	D	2	NAG	O5-C5-C6	2.05	110.42	107.20

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	1	NAG	C1
2	D	1	NAG	C1



There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3	BMA	1	0
2	D	2	NAG	2	0
2	C	2	NAG	2	0
2	C	1	NAG	3	0
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	1610/1661 (96%)	-0.07	55 (3%)	45	19	15, 101, 169, 200	0
1	B	1610/1661 (96%)	-0.12	47 (2%)	51	23	17, 96, 169, 200	0
All	All	3220/3322 (96%)	-0.09	102 (3%)	47	20	15, 99, 169, 200	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	671	VAL	6.0
1	A	742	LEU	5.6
1	A	741	ALA	5.2
1	A	276	ASP	5.0
1	A	348	VAL	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	C	3	11/12	0.72	0.41	81,81,82,82	0
2	BMA	D	3	11/12	0.79	0.23	80,81,81,82	0
2	NAG	C	2	14/15	0.80	0.32	80,81,82,82	0
2	NAG	C	1	14/15	0.83	0.24	74,80,81,82	0
2	NAG	D	2	14/15	0.88	0.26	80,80,81,82	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	D	1	14/15	0.93	0.15	73,80,81,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.