



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 04:35 PM BST

PDB ID : 2B3B
Title : Thermus thermophilus Glucose/Galactose Binding Protein With Bound Glucose
Authors : Cuneo, M.J.; Changela, A.; Warren, J.J.; Beese, L.S.; Hellinga, H.W.
Deposited on : 2005-09-20
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	<div> <div>2%</div> <div>94%</div> <div>5%</div> <div>• •</div> </div>
1	B	400	<div> <div>2%</div> <div>93%</div> <div>5%</div> <div>•</div> </div>
1	C	400	<div> <div>3%</div> <div>94%</div> <div>5%</div> <div>• •</div> </div>
1	D	400	<div> <div>0%</div> <div>95%</div> <div>5%</div> <div>• •</div> </div>
1	E	400	<div> <div>0%</div> <div>95%</div> <div>5%</div> <div>• •</div> </div>
1	F	400	<div> <div>2%</div> <div>93%</div> <div>5%</div> <div>•</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glucose-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	B	391	Total	C	N	O	S	0	0	0
			3028	1936	534	544	14			
1	C	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	D	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	E	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	F	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q72KX2
A	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
A	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
A	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
A	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
A	395	HIS	-	expression tag	UNP Q72KX2
A	396	HIS	-	expression tag	UNP Q72KX2
A	397	HIS	-	expression tag	UNP Q72KX2
A	398	HIS	-	expression tag	UNP Q72KX2
A	399	HIS	-	expression tag	UNP Q72KX2
A	400	HIS	-	expression tag	UNP Q72KX2
B	1	MET	-	initiating methionine	UNP Q72KX2
B	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
B	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
B	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
B	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
B	395	HIS	-	expression tag	UNP Q72KX2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	396	HIS	-	expression tag	UNP Q72KX2
B	397	HIS	-	expression tag	UNP Q72KX2
B	398	HIS	-	expression tag	UNP Q72KX2
B	399	HIS	-	expression tag	UNP Q72KX2
B	400	HIS	-	expression tag	UNP Q72KX2
C	1	MET	-	initiating methionine	UNP Q72KX2
C	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
C	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
C	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
C	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
C	395	HIS	-	expression tag	UNP Q72KX2
C	396	HIS	-	expression tag	UNP Q72KX2
C	397	HIS	-	expression tag	UNP Q72KX2
C	398	HIS	-	expression tag	UNP Q72KX2
C	399	HIS	-	expression tag	UNP Q72KX2
C	400	HIS	-	expression tag	UNP Q72KX2
D	1	MET	-	initiating methionine	UNP Q72KX2
D	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
D	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
D	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
D	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
D	395	HIS	-	expression tag	UNP Q72KX2
D	396	HIS	-	expression tag	UNP Q72KX2
D	397	HIS	-	expression tag	UNP Q72KX2
D	398	HIS	-	expression tag	UNP Q72KX2
D	399	HIS	-	expression tag	UNP Q72KX2
D	400	HIS	-	expression tag	UNP Q72KX2
E	1	MET	-	initiating methionine	UNP Q72KX2
E	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
E	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
E	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
E	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
E	395	HIS	-	expression tag	UNP Q72KX2
E	396	HIS	-	expression tag	UNP Q72KX2
E	397	HIS	-	expression tag	UNP Q72KX2
E	398	HIS	-	expression tag	UNP Q72KX2
E	399	HIS	-	expression tag	UNP Q72KX2
E	400	HIS	-	expression tag	UNP Q72KX2
F	1	MET	-	initiating methionine	UNP Q72KX2
F	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
F	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
F	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
F	395	HIS	-	expression tag	UNP Q72KX2
F	396	HIS	-	expression tag	UNP Q72KX2
F	397	HIS	-	expression tag	UNP Q72KX2
F	398	HIS	-	expression tag	UNP Q72KX2
F	399	HIS	-	expression tag	UNP Q72KX2
F	400	HIS	-	expression tag	UNP Q72KX2

- Molecule 2 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	E	1	Total	C	O	0	0
			12	6	6		
2	F	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			12	6	6		
3	D	1	Total	C	O	0	0
			12	6	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	302	Total	O	0	0
			302	302		
4	B	341	Total	O	0	0
			341	341		
4	C	315	Total	O	0	0
			315	315		
4	D	326	Total	O	0	0
			326	326		
4	E	325	Total	O	0	0
			325	325		
4	F	328	Total	O	0	0
			328	328		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: glucose-binding protein




- Molecule 1: glucose-binding protein



- Molecule 1: glucose-binding protein



- Molecule 1: glucose-binding protein




- Molecule 1: glucose-binding protein





● Molecule 1: glucose-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	121.64Å 134.87Å 159.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 49.48 – 1.95	Depositor EDS
% Data completeness (in resolution range)	95.6 (50.00-1.95) 95.6 (49.48-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.95Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.185 , 0.225 0.194 , 0.234	Depositor DCC
R_{free} test set	9120 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtrriage
Anisotropy	0.225	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20217	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 75.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3501e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.61	0/3116	0.68	1/4235 (0.0%)
1	B	0.59	0/3108	0.69	4/4224 (0.1%)
1	C	0.60	0/3116	0.67	2/4235 (0.0%)
1	D	0.61	0/3116	0.68	0/4235
1	E	0.59	0/3116	0.68	1/4235 (0.0%)
1	F	0.59	0/3116	0.69	1/4235 (0.0%)
All	All	0.60	0/18688	0.68	9/25399 (0.0%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	80	ARG	NE-CZ-NH1	6.69	123.65	120.30
1	B	297	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	B	297	ARG	NE-CZ-NH2	-5.86	117.37	120.30
1	C	47	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	F	210	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	218	ASP	CB-CG-OD1	5.61	123.35	118.30
1	C	47	ARG	NE-CZ-NH2	-5.57	117.52	120.30
1	B	47	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	B	47	ARG	NE-CZ-NH1	5.32	122.96	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3036	0	3012	8	0
1	B	3028	0	3001	11	0
1	C	3036	0	3012	10	0
1	D	3036	0	3012	6	0
1	E	3036	0	3012	7	0
1	F	3036	0	3012	12	0
2	A	12	0	12	2	0
2	B	12	0	12	2	0
2	E	12	0	12	1	0
2	F	12	0	12	5	0
3	C	12	0	12	0	0
3	D	12	0	12	0	0
4	A	302	0	0	0	0
4	B	341	0	0	2	0
4	C	315	0	0	1	0
4	D	326	0	0	0	0
4	E	325	0	0	2	0
4	F	328	0	0	1	0
All	All	20217	0	18133	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:169:GLN:OE1	1:F:368:LEU:HD11	1.62	0.98
1:C:169:GLN:OE1	1:C:368:LEU:HD11	1.91	0.71
1:D:47:ARG:HD3	1:D:354:GLU:OE2	1.95	0.67
1:C:364:MET:O	1:C:368:LEU:HG	2.04	0.57
1:A:348:HIS:HD1	2:A:2605:GLC:HO1	1.50	0.57
1:E:348:HIS:ND1	2:E:2609:GLC:O1	2.40	0.55
1:F:20:LEU:HG	1:F:308:SER:HB3	1.89	0.54
1:A:20:LEU:HG	1:A:308:SER:HB3	1.90	0.54
1:A:34:ILE:HG12	1:C:154:GLN:HA	1.91	0.51
1:C:43:GLY:O	1:C:47:ARG:HG3	2.11	0.51
1:F:270:GLN:H	1:F:270:GLN:CD	2.14	0.50
1:D:20:LEU:HG	1:D:308:SER:HB3	1.93	0.50
1:F:8:TRP:CZ2	2:F:2610:GLC:H1	2.46	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:HIS:CE1	1:A:314:SER:HB3	2.48	0.49
1:F:119:HIS:CE1	1:F:314:SER:HB3	2.48	0.49
1:F:8:TRP:HE1	2:F:2610:GLC:H1	1.77	0.48
1:A:363:VAL:O	1:A:366:ILE:HG13	2.13	0.48
1:A:8:TRP:CZ2	2:A:2605:GLC:H1	2.49	0.47
1:B:169:GLN:OE1	1:B:368:LEU:HD11	2.13	0.47
1:E:54:MET:O	1:E:80:ARG:NH2	2.47	0.47
1:D:14:GLY:N	1:D:15:PRO:CD	2.78	0.47
1:B:386:GLN:NE2	4:B:2899:HOH:O	2.48	0.47
1:E:94:LEU:CD1	1:E:95:GLN:HG3	2.45	0.46
1:D:126:TYR:CE1	1:D:262:ALA:HB3	2.50	0.46
1:B:126:TYR:CD1	1:B:131:LEU:HD22	2.51	0.46
1:A:127:LEU:HD13	1:A:231:VAL:HG12	1.98	0.46
1:B:300:GLY:O	1:B:318:ARG:HD2	2.16	0.46
1:C:143:ASP:OD2	1:D:369:GLN:NE2	2.50	0.44
1:E:155:LYS:NZ	4:E:2923:HOH:O	2.21	0.44
1:F:348:HIS:ND1	2:F:2610:GLC:O1	2.46	0.44
1:B:34:ILE:N	1:B:34:ILE:HD12	2.32	0.44
1:F:99:LYS:HG3	4:F:2854:HOH:O	2.18	0.44
1:C:265:PRO:HG2	1:C:270:GLN:HG2	2.00	0.43
1:B:348:HIS:ND1	2:B:2606:GLC:O1	2.50	0.43
1:A:14:GLY:N	1:A:15:PRO:CD	2.81	0.43
1:E:257:PRO:HB2	1:E:328:ALA:HB3	2.00	0.43
1:F:14:GLY:N	1:F:15:PRO:CD	2.82	0.43
1:B:335:ARG:HG2	4:B:2795:HOH:O	2.18	0.42
1:F:323:PRO:O	1:F:331:GLN:NE2	2.47	0.42
1:F:8:TRP:NE1	2:F:2610:GLC:H1	2.35	0.42
1:B:119:HIS:CE1	1:B:314:SER:HB3	2.55	0.41
1:C:138:PRO:HA	1:C:139:PRO:HD2	1.97	0.41
1:E:25:LYS:HE2	4:E:2673:HOH:O	2.20	0.41
1:C:338:ARG:NE	4:C:2751:HOH:O	2.53	0.41
1:B:364:MET:O	1:B:368:LEU:HG	2.21	0.41
1:F:8:TRP:HZ2	2:F:2610:GLC:H1	1.86	0.41
1:B:8:TRP:HE1	2:B:2606:GLC:H1	1.85	0.41
1:C:7:SER:HB2	1:C:64:GLN:OE1	2.21	0.41
1:C:126:TYR:CE1	1:C:262:ALA:HB3	2.56	0.40
1:D:72:ILE:HA	1:D:76:VAL:HB	2.04	0.40
1:B:149:CYS:O	1:B:153:LYS:HG3	2.21	0.40
1:E:94:LEU:HD12	1:E:95:GLN:CG	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/400 (98%)	386 (99%)	4 (1%)	0	100	100
1	B	389/400 (97%)	383 (98%)	6 (2%)	0	100	100
1	C	390/400 (98%)	383 (98%)	7 (2%)	0	100	100
1	D	390/400 (98%)	384 (98%)	6 (2%)	0	100	100
1	E	390/400 (98%)	385 (99%)	5 (1%)	0	100	100
1	F	390/400 (98%)	386 (99%)	4 (1%)	0	100	100
All	All	2339/2400 (98%)	2307 (99%)	32 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/316 (98%)	307 (99%)	2 (1%)	86	85
1	B	308/316 (98%)	307 (100%)	1 (0%)	92	92
1	C	309/316 (98%)	307 (99%)	2 (1%)	86	85
1	D	309/316 (98%)	306 (99%)	3 (1%)	76	74
1	E	309/316 (98%)	305 (99%)	4 (1%)	69	65
1	F	309/316 (98%)	305 (99%)	4 (1%)	69	65
All	All	1853/1896 (98%)	1837 (99%)	16 (1%)	78	77

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	273	PHE
1	A	276	LEU
1	B	276	LEU
1	C	276	LEU
1	C	341	ARG
1	D	1	MET
1	D	276	LEU
1	D	368	LEU
1	E	52	THR
1	E	213	ASP
1	E	276	LEU
1	E	341	ARG
1	F	212	LEU
1	F	213	ASP
1	F	273	PHE
1	F	276	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	369	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	E	2609	-	12,12,12	0.73	0	17,17,17	2.25	6 (35%)
2	GLC	F	2610	-	12,12,12	0.71	0	17,17,17	2.10	5 (29%)
3	BGC	C	2607	-	12,12,12	0.65	0	17,17,17	0.69	0
2	GLC	A	2605	-	12,12,12	0.80	1 (8%)	17,17,17	2.14	6 (35%)
3	BGC	D	2608	-	12,12,12	0.76	0	17,17,17	0.79	0
2	GLC	B	2606	-	12,12,12	0.70	0	17,17,17	2.41	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	2609	-	-	0/2/22/22	0/1/1/1
2	GLC	F	2610	-	-	0/2/22/22	0/1/1/1
3	BGC	C	2607	-	-	0/2/22/22	0/1/1/1
2	GLC	A	2605	-	-	0/2/22/22	0/1/1/1
3	BGC	D	2608	-	-	0/2/22/22	0/1/1/1
2	GLC	B	2606	-	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2605	GLC	C1-C2	2.11	1.57	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2609	GLC	O5-C1-C2	5.75	120.54	110.28
2	F	2610	GLC	C1-O5-C5	5.36	123.77	113.66
2	A	2605	GLC	O5-C1-C2	5.24	119.64	110.28
2	B	2606	GLC	O5-C1-C2	5.10	119.39	110.28
2	B	2606	GLC	C1-O5-C5	4.92	122.95	113.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2609	GLC	C1-O5-C5	4.74	122.60	113.66
2	F	2610	GLC	O5-C1-C2	4.41	118.15	110.28
2	A	2605	GLC	C1-O5-C5	4.34	121.84	113.66
2	B	2606	GLC	O2-C2-C1	3.64	117.59	109.16
2	B	2606	GLC	O5-C5-C4	3.06	115.26	109.69
2	F	2610	GLC	C1-C2-C3	2.93	116.39	110.31
2	A	2605	GLC	C1-C2-C3	2.83	116.19	110.31
2	B	2606	GLC	C1-C2-C3	2.81	116.14	110.31
2	B	2606	GLC	O1-C1-C2	2.74	116.73	109.03
2	E	2609	GLC	O2-C2-C1	2.58	115.15	109.16
2	A	2605	GLC	O1-C1-C2	2.58	116.29	109.03
2	E	2609	GLC	O5-C5-C4	2.54	114.31	109.69
2	A	2605	GLC	O2-C2-C1	2.48	114.92	109.16
2	E	2609	GLC	O3-C3-C2	-2.24	105.17	110.35
2	B	2606	GLC	O3-C3-C2	-2.21	105.25	110.35
2	A	2605	GLC	O5-C5-C4	2.20	113.69	109.69
2	E	2609	GLC	C1-C2-C3	2.15	114.78	110.31
2	F	2610	GLC	O1-C1-C2	2.12	115.01	109.03
2	F	2610	GLC	O2-C2-C1	2.07	113.96	109.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2609	GLC	1	0
2	F	2610	GLC	5	0
2	A	2605	GLC	2	0
2	B	2606	GLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/400 (98%)	0.10	8 (2%) 65 73	15, 25, 37, 44	0
1	B	391/400 (97%)	0.11	7 (1%) 68 76	14, 23, 35, 52	0
1	C	392/400 (98%)	0.19	13 (3%) 46 56	16, 25, 38, 61	0
1	D	392/400 (98%)	-0.07	3 (0%) 86 90	16, 25, 33, 43	0
1	E	392/400 (98%)	0.03	4 (1%) 82 87	16, 26, 35, 41	0
1	F	392/400 (98%)	0.11	9 (2%) 60 69	15, 25, 38, 44	0
All	All	2351/2400 (97%)	0.08	44 (1%) 66 74	14, 25, 36, 61	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	392	LEU	8.3
1	E	94	LEU	6.1
1	C	391	ARG	4.8
1	B	391	ARG	4.4
1	B	132	LYS	3.7
1	F	132	LYS	3.6
1	C	258	GLY	3.4
1	C	259	THR	3.0
1	A	133	GLY	2.9
1	A	136	VAL	2.9
1	C	256	LYS	2.9
1	A	129	ALA	2.8
1	C	390	GLY	2.8
1	C	79	ASN	2.8
1	C	129	ALA	2.7
1	C	132	LYS	2.7
1	A	50	LEU	2.6
1	F	1	MET	2.6
1	E	132	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	131	LEU	2.5
1	D	203	ARG	2.5
1	F	134	TRP	2.5
1	F	129	ALA	2.5
1	B	256	LYS	2.5
1	B	99	LYS	2.4
1	D	30	GLY	2.4
1	B	259	THR	2.4
1	A	32	GLU	2.3
1	E	140	ARG	2.3
1	F	32	GLU	2.3
1	A	134	TRP	2.2
1	F	136	VAL	2.2
1	E	134	TRP	2.2
1	F	131	LEU	2.2
1	F	133	GLY	2.2
1	C	108	LYS	2.2
1	B	129	ALA	2.1
1	C	270	GLN	2.1
1	F	140	ARG	2.1
1	C	77	VAL	2.1
1	A	140	ARG	2.0
1	A	154	GLN	2.0
1	B	79	ASN	2.0
1	D	338	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	A	2605	12/12	0.95	0.12	14,15,20,21	0
2	GLC	F	2610	12/12	0.96	0.10	14,17,18,21	0
2	GLC	E	2609	12/12	0.96	0.10	13,17,19,21	0
3	BGC	D	2608	12/12	0.97	0.07	11,15,17,19	0
2	GLC	B	2606	12/12	0.97	0.12	15,17,18,19	0
3	BGC	C	2607	12/12	0.98	0.10	12,14,15,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.