



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 09:38 AM BST

PDB ID : 6B4K
Title : Crystal structure of human DDX19B(AMPPNP)
Authors : Lin, D.H.; Correia, A.R.; Cai, S.W.; Huber, F.M.; Jette, C.A.; Hoelz, A.
Deposited on : 2017-09-26
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	430	<div> <div>6%</div> <div>92%</div> <div>5%</div> </div>
1	B	430	<div> <div>19%</div> <div>92%</div> <div>6%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13924 atoms, of which 6898 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase DDX19B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	419	Total	C	H	N	O	S	0	11	0
			6844	2141	3457	591	635	20			
1	B	420	Total	C	H	N	O	S	0	1	0
			6737	2106	3402	581	628	20			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	GLY	-	expression tag	UNP Q9UMR2
A	51	PRO	-	expression tag	UNP Q9UMR2
A	52	HIS	-	expression tag	UNP Q9UMR2
A	53	MET	-	expression tag	UNP Q9UMR2
B	50	GLY	-	expression tag	UNP Q9UMR2
B	51	PRO	-	expression tag	UNP Q9UMR2
B	52	HIS	-	expression tag	UNP Q9UMR2
B	53	MET	-	expression tag	UNP Q9UMR2

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	H	N	O	P	0	0
			48	10	17	6	12	3		
3	B	1	Total	C	H	N	O	P	0	0
			47	10	16	6	12	3		

- Molecule 4 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			9	3	2	4		
4	A	1	Total	C	H	O	0	0
			9	3	2	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			9	3	2	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total	O	0	2
			139	139		
5	B	78	Total	O	0	2
			80	80		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent RNA helicase DDX19B



- Molecule 1: ATP-dependent RNA helicase DDX19B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.23 Å 45.36 Å 127.39 Å 90.00° 97.03° 90.00°	Depositor
Resolution (Å)	42.14 – 2.20 126.43 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (42.14-2.20) 99.3 (126.43-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.20 Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.207 , 0.248 0.209 , 0.250	Depositor DCC
R_{free} test set	1998 reflections (4.14%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13924	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3168e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MLI, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/3475	0.46	0/4688
1	B	0.27	0/3387	0.46	0/4571
All	All	0.27	0/6862	0.46	0/9259

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3387	3457	3404	15	0
1	B	3335	3402	3400	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	17	13	1	0
3	B	31	16	13	0	0
4	A	14	4	4	1	0
4	B	7	2	2	0	0
5	A	139	0	0	1	0
5	B	80	0	0	0	0
All	All	7026	6898	6836	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:LEU:HD11	1:A:215:ILE:HG12	1.79	0.64
1:A:251:GLN:NE2	5:A:602:HOH:O	2.32	0.61
1:B:443:VAL:HG11	1:B:449:MET:HA	1.89	0.55
1:B:316:SER:N	1:B:444:ASP:OD2	2.42	0.53
1:B:443:VAL:HG11	1:B:449:MET:CA	2.41	0.50
1:A:366:GLU:CB	1:B:203:LEU:HD11	2.42	0.49
1:B:316:SER:CA	1:B:444:ASP:OD2	2.61	0.49
1:B:342:THR:HG22	1:B:344:LYS:H	1.77	0.48
1:B:466:LEU:HD13	1:B:477:ILE:HD11	1.96	0.48
1:A:251:GLN:NE2	4:A:504:MLI:O8	2.47	0.47
1:B:467:ASP:OD2	1:B:469:ASP:OD2	2.31	0.47
1:A:64:LYS:NZ	3:A:502:ANP:O3G	2.45	0.46
1:B:323:ALA:HA	1:B:468:THR:HG21	1.98	0.46
1:B:203:LEU:HD23	1:B:230:PHE:CD2	2.51	0.45
1:B:277:PHE:CZ	1:B:296:LYS:HE2	2.52	0.44
1:A:67:ARG:NH2	1:A:140:SER:OG	2.50	0.44
1:B:66:ILE:O	1:B:380:ARG:NH1	2.50	0.43
1:A:308:LYS:HE2	1:A:310:TYR:OH	2.18	0.43
1:A:310:TYR:CZ	1:A:477:ILE:HG22	2.53	0.42
1:A:82:ARG:NE	1:A:291:ASP:HB3	2.34	0.42
1:B:316:SER:C	1:B:444:ASP:OD2	2.58	0.42
1:B:304:LEU:HD22	1:B:424:LEU:CD1	2.50	0.42
1:B:203:LEU:HD23	1:B:230:PHE:CE2	2.55	0.42
1:A:368:MET:HE1	1:B:201:ASN:CG	2.41	0.41
1:A:474:ILE:O	1:A:477:ILE:HG13	2.21	0.41
1:B:380:ARG:NE	1:B:400:GLU:HG2	2.36	0.41
1:A:72:ASP:N	1:A:72:ASP:OD1	2.54	0.41
1:A:145:THR:HA	1:A:148:PHE:CE2	2.56	0.40
1:A:361:ALA:HB1	1:A:375:VAL:HG13	2.03	0.40
1:A:331:ILE:HD13	1:A:404:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/430 (99%)	421 (99%)	3 (1%)	0	100	100
1	B	417/430 (97%)	411 (99%)	6 (1%)	0	100	100
All	All	841/860 (98%)	832 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/378 (100%)	379 (100%)	1 (0%)	92	97
1	B	371/378 (98%)	369 (100%)	2 (0%)	88	94
All	All	751/756 (99%)	748 (100%)	3 (0%)	91	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	277	PHE
1	B	64	LYS
1	B	343	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MLI	B	503	-	0,6,6	0.00	-	0,7,7	0.00	-
3	ANP	B	502	2	29,33,33	0.65	0	31,52,52	0.64	1 (3%)
4	MLI	A	504	-	0,6,6	0.00	-	0,7,7	0.00	-
3	ANP	A	502	2	29,33,33	0.72	0	31,52,52	0.69	1 (3%)
4	MLI	A	503	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	B	503	-	-	0/0/4/4	-
3	ANP	B	502	2	-	5/14/38/38	0/3/3/3
4	MLI	A	504	-	-	0/0/4/4	-
3	ANP	A	502	2	-	4/14/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	A	503	-	-	0/0/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	ANP	C5-C6-N6	2.41	124.01	120.35
3	A	502	ANP	C5-C6-N6	2.36	123.94	120.35

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	ANP	PB-N3B-PG-O1G
3	B	502	ANP	PG-N3B-PB-O1B
3	B	502	ANP	PA-O3A-PB-O1B
3	A	502	ANP	PB-N3B-PG-O1G
3	A	502	ANP	PG-N3B-PB-O1B
3	A	502	ANP	PA-O3A-PB-O1B
3	A	502	ANP	PA-O3A-PB-O2B
3	B	502	ANP	PA-O3A-PB-O2B
3	B	502	ANP	PG-N3B-PB-O3A


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	MLI	1	0
3	A	502	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/430 (97%)	0.50	25 (5%) 21 20	24, 61, 124, 154	0
1	B	420/430 (97%)	1.05	81 (19%) 1 1	30, 76, 164, 249	0
All	All	839/860 (97%)	0.77	106 (12%) 3 3	24, 67, 151, 249	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	329	GLY	11.1
1	B	478	ALA	10.3
1	A	334	ALA	8.0
1	B	316	SER	7.4
1	B	313	LEU	7.2
1	B	330	ALA	7.1
1	B	324	LEU	7.1
1	B	331	ILE	6.8
1	B	311	TYR	6.5
1	A	328	TYR	6.4
1	B	477	ILE	6.0
1	B	315	SER	5.6
1	B	451	ILE	5.5
1	B	357	GLY	5.4
1	B	314	CYS	5.3
1	B	465	ARG	5.2
1	B	332	THR	4.9
1	B	474	ILE	4.7
1	B	408	PHE	4.5
1	B	360	VAL	4.3
1	B	453	ASN	4.2
1	B	385	LYS	4.2
1	A	479	ASN	4.2
1	B	446	LYS	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	307	ILE	3.7
1	B	343	ARG	3.6
1	B	470	ASP	3.6
1	A	331	ILE	3.6
1	B	333	ILE	3.5
1	B	412	VAL	3.4
1	B	328	TYR	3.4
1	B	277	PHE	3.3
1	B	321	PHE	3.3
1	B	459	PHE	3.2
1	B	410	LEU	3.2
1	B	469	ASP	3.2
1	B	472	ASP	3.2
1	B	448	SER	3.2
1	B	386	VAL	3.1
1	B	351	ALA	3.1
1	B	83	ASP	3.1
1	B	398	ASP	3.1
1	A	311	TYR	3.1
1	B	303	THR	3.0
1	B	413	ASP	3.0
1	B	302	GLU	2.9
1	B	475	GLU	2.9
1	B	82	ARG	2.9
1	A	89[A]	TYR	2.9
1	B	319	GLU	2.9
1	B	348	TRP	2.9
1	B	402	VAL	2.8
1	A	462	LYS	2.8
1	B	473	GLU	2.8
1	A	460	ASN	2.8
1	B	476	LYS	2.8
1	A	446	LYS	2.7
1	B	306	THR	2.7
1	A	478	ALA	2.7
1	B	452	LEU	2.7
1	B	359	GLN	2.7
1	B	468	THR	2.7
1	B	471	LEU	2.7
1	A	443	VAL	2.6
1	A	205	ARG	2.6
1	B	327	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	455	ILE	2.6
1	A	444	ASP	2.6
1	B	460	ASN	2.6
1	A	465	ARG	2.6
1	B	353	LEU	2.5
1	B	349	LEU	2.4
1	B	85	ASN	2.4
1	B	87	PRO	2.4
1	A	301	GLU	2.4
1	B	443	VAL	2.4
1	B	282	TRP	2.3
1	B	318	ASP	2.3
1	B	425	HIS	2.3
1	A	445	SER	2.3
1	A	459	PHE	2.3
1	A	327	LEU	2.2
1	A	335	GLN	2.2
1	B	355	LYS	2.2
1	B	428	GLY	2.2
1	B	322	GLN	2.2
1	B	84	PRO	2.2
1	B	401	GLN	2.2
1	A	476	LYS	2.1
1	B	88	LEU	2.1
1	B	335	GLN	2.1
1	B	416	GLY	2.1
1	B	463	ILE	2.1
1	A	313	LEU	2.1
1	A	464	GLU	2.1
1	B	399	VAL	2.1
1	A	358	HIS	2.1
1	B	346	ALA	2.0
1	B	350	ALA	2.0
1	A	317	ARG	2.0
1	B	89	TYR	2.0
1	B	86	SER	2.0
1	B	424	LEU	2.0
1	B	337	MET	2.0
1	A	343	ARG	2.0
1	B	262	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MLI	A	504	7/7	0.66	0.35	57,87,110,113	9
2	MG	A	501	1/1	0.85	0.23	54,54,54,54	0
2	MG	B	501	1/1	0.91	0.30	69,69,69,69	0
3	ANP	B	502	31/31	0.96	0.15	40,45,88,121	0
4	MLI	A	503	7/7	0.96	0.12	43,50,60,64	0
4	MLI	B	503	7/7	0.97	0.14	43,50,54,57	0
3	ANP	A	502	31/31	0.97	0.18	37,46,73,111	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.