



wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 09:26 pm BST

PDB ID : 2B5L
Title : Crystal Structure of DDB1 In Complex with Simian Virus 5 V Protein
Authors : Li, T.; Chen, X.; Garbutt, K.C.; Zhou, P.; Zheng, N.
Deposited on : 2005-09-28
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 54% 37% 7% .. </div> </div>
1	B	1140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 41% 48% 10% .. </div> </div>
2	C	222	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 30% 37% 9% 22% </div> </div>
2	D	222	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 25% 37% 13% 21% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called damage-specific DNA binding protein 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1132	Total	C	N	O	S	0	0	0
			8860	5610	1493	1708	49			
1	B	1134	Total	C	N	O	S	0	0	0
			8876	5619	1495	1713	49			

- Molecule 2 is a protein called Nonstructural protein V.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	174	Total	C	N	O	S	0	0	0
			1324	826	229	261	8			
2	D	175	Total	C	N	O	S	0	0	0
			1335	834	233	260	8			


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		




• Molecule 1: damage-specific DNA binding protein 1







• Molecule 2: Nonstructural protein V



• Molecule 2: Nonstructural protein V





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.76Å 240.79Å 117.18Å 90.00° 101.79° 90.00°	Depositor
Resolution (Å)	47.80 – 2.85 47.77 – 2.63	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.80-2.85) 86.6 (47.77-2.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.65Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.229 , 0.299 0.226 , 0.293	Depositor DCC
R_{free} test set	3805 reflections (4.13%)	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.604	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 62.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	20399	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	1/9022 (0.0%)	0.77	7/12219 (0.1%)
1	B	0.45	3/9039 (0.0%)	0.91	41/12244 (0.3%)
2	C	0.56	0/1351	0.95	9/1832 (0.5%)
2	D	0.71	2/1361 (0.1%)	1.24	19/1843 (1.0%)
All	All	0.48	6/20773 (0.0%)	0.88	76/28138 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	1
2	D	0	1
All	All	1	2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	189	TRP	C-N	-15.06	0.99	1.34
2	D	190	CYS	C-N	11.09	1.59	1.34
1	B	748	GLY	N-CA	-7.48	1.34	1.46
1	B	431	GLY	N-CA	-6.93	1.35	1.46
1	A	725	CYS	CB-SG	-5.43	1.73	1.81

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	749	THR	N-CA-C	-19.59	58.10	111.00
1	B	35	LYS	CB-CA-C	18.15	146.70	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	15	LYS	N-CA-C	-16.98	65.16	111.00
1	B	431	GLY	N-CA-C	-16.54	71.75	113.10
1	B	749	THR	N-CA-CB	14.29	137.44	110.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	1	MET	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	430	VAL	Peptide
2	D	190	CYS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8860	0	8828	466	0
1	B	8876	0	8842	793	0
2	C	1324	0	1306	128	0
2	D	1335	0	1323	177	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
All	All	20399	0	20299	1549	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1549 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:LEU:HD21	1:B:374:GLN:CG	1.51	1.38
1:B:367:LEU:CD2	1:B:374:GLN:HB2	1.53	1.37
1:B:367:LEU:HD21	1:B:374:GLN:CB	1.50	1.37

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:742:VAL:CG2	1:B:750:THR:HG21	1.57	1.35
1:B:367:LEU:CD2	1:B:374:GLN:CG	2.06	1.34

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1126/1140 (99%)	960 (85%)	128 (11%)	38 (3%)	3	12
1	B	1130/1140 (99%)	892 (79%)	145 (13%)	93 (8%)	1	1
2	C	168/222 (76%)	123 (73%)	33 (20%)	12 (7%)	1	2
2	D	169/222 (76%)	115 (68%)	32 (19%)	22 (13%)	0	0
All	All	2593/2724 (95%)	2090 (81%)	338 (13%)	165 (6%)	1	3

5 of 165 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	LYS
1	A	269	SER
1	A	290	GLN
1	A	341	ASN
1	A	370	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	992/999 (99%)	900 (91%)	92 (9%)	9	24
1	B	994/999 (100%)	925 (93%)	69 (7%)	15	38
2	C	151/191 (79%)	134 (89%)	17 (11%)	6	16
2	D	151/191 (79%)	133 (88%)	18 (12%)	5	13
All	All	2288/2380 (96%)	2092 (91%)	196 (9%)	10	27

5 of 196 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1069	GLU
1	B	402	ILE
2	D	51	LEU
1	A	1135	GLU
1	B	147	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 82 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1056	ASN
1	B	255	GLN
2	C	210	GLN
1	A	1070	HIS
1	B	22	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	189:TRP	C	190:CYS	N	0.99

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1132/1140 (99%)	-0.29	10 (0%) 84 84	18, 47, 99, 160	0
1	B	1134/1140 (99%)	0.01	35 (3%) 49 44	29, 68, 122, 173	0
2	C	174/222 (78%)	0.22	12 (6%) 16 12	32, 62, 118, 183	0
2	D	175/222 (78%)	0.56	19 (10%) 5 4	39, 81, 138, 168	0
All	All	2615/2724 (95%)	-0.07	76 (2%) 51 47	18, 59, 117, 183	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1018	GLY	5.9
1	B	750	THR	5.2
1	B	746	SER	5.0
1	B	780	THR	4.9
2	D	16	LEU	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	C	3001	1/1	0.83	0.09	35,35,35,35	0
3	ZN	C	3002	1/1	0.89	0.07	35,35,35,35	0
3	ZN	D	3003	1/1	0.96	0.05	35,35,35,35	0
3	ZN	D	3004	1/1	0.96	0.08	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.