



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 07:09 pm BST

PDB ID : 3B77  
Title : Crystal structure of a ph domain containing bacterial protein (exig\_2160) from  
exiguobacterium sibiricum 255-15 at 2.42 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-10-30  
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>•</div> </div> </div>
1	B	193	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>• 5%</div> </div> </div>
1	C	193	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>• 5%</div> </div> </div>
1	D	193	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>6%</div> <div>• 5%</div> </div> </div>
1	E	193	<div> <div>10%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
1	F	193	<div> <div>10%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>5%</div> </div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9166 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	188	Total	C	N	O	S	Se	0	0	0
			1525	983	247	291	1	3			
1	B	184	Total	C	N	O	S	Se	0	1	0
			1507	975	244	284	1	3			
1	C	184	Total	C	N	O	S	Se	0	0	0
			1490	965	239	282	1	3			
1	D	184	Total	C	N	O	S	Se	0	0	0
			1494	963	242	285	1	3			
1	E	183	Total	C	N	O	S	Se	0	0	1
			1468	953	234	277	1	3			
1	F	183	Total	C	N	O	S	Se	0	2	0
			1509	976	245	284	1	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	-	LEADER SEQUENCE	UNP Q41E03
B	12	GLY	-	LEADER SEQUENCE	UNP Q41E03
C	12	GLY	-	LEADER SEQUENCE	UNP Q41E03
D	12	GLY	-	LEADER SEQUENCE	UNP Q41E03
E	12	GLY	-	LEADER SEQUENCE	UNP Q41E03
F	12	GLY	-	LEADER SEQUENCE	UNP Q41E03

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	34	Total	O	0	0
			34	34		
2	B	43	Total	O	0	0
			43	43		
2	C	37	Total	O	0	0
			37	37		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	20	Total	O	0	0
			20	20		
2	E	17	Total	O	0	0
			17	17		
2	F	22	Total	O	0	0
			22	22		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein





● Molecule 1: Uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.99Å 150.99Å 76.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.73 – 2.42 47.75 – 2.42	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.73-2.42) 99.7 (47.75-2.42)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, $R_{free}$	0.214 , 0.254 0.219 , 0.257	Depositor DCC
$R_{free}$ test set	3326 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.5	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 51.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.045 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9166	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/1561	0.75	3/2104 (0.1%)
1	B	0.85	0/1546	0.85	4/2081 (0.2%)
1	C	0.70	0/1526	0.76	0/2058
1	D	0.61	0/1529	0.69	1/2059 (0.0%)
1	E	0.64	0/1504	0.72	2/2027 (0.1%)
1	F	0.67	0/1551	0.74	2/2088 (0.1%)
All	All	0.70	0/9217	0.75	12/12417 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	157	LEU	CB-CG-CD1	-6.37	100.18	111.00
1	E	157	LEU	CB-CG-CD1	-6.21	100.44	111.00
1	B	106	GLY	N-CA-C	-5.98	98.16	113.10
1	F	106	GLY	N-CA-C	-5.84	98.51	113.10
1	B	160	LEU	CA-CB-CG	5.80	128.64	115.30
1	E	106	GLY	N-CA-C	-5.74	98.76	113.10
1	B	180	ASP	CB-CG-OD1	-5.73	113.14	118.30
1	A	173	ASP	CB-CG-OD1	5.65	123.39	118.30
1	A	106	GLY	N-CA-C	-5.51	99.31	113.10
1	A	130	LEU	CA-CB-CG	5.42	127.77	115.30
1	D	106	GLY	N-CA-C	-5.07	100.42	113.10
1	B	63	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1525	0	1425	14	0
1	B	1507	0	1419	10	0
1	C	1490	0	1391	11	0
1	D	1494	0	1395	9	0
1	E	1468	0	1369	11	0
1	F	1509	0	1433	10	0
2	A	34	0	0	0	0
2	B	43	0	0	1	0
2	C	37	0	0	0	0
2	D	20	0	0	0	0
2	E	17	0	0	0	0
2	F	22	0	0	0	0
All	All	9166	0	8432	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:MSE:CE	1:B:88:MSE:SE	2.20	1.40
1:A:88:MSE:SE	1:A:88:MSE:CE	2.17	1.39
1:D:159:GLY:O	1:D:160:LEU:CB	2.43	0.66
1:B:100:GLU:OE1	1:B:102:LYS:HD2	1.95	0.66
1:C:15:ILE:N	1:C:15:ILE:HD12	2.15	0.61
1:D:130:LEU:HD21	1:E:76:ARG:HD2	1.85	0.59
1:C:188:LYS:HD3	1:D:95:VAL:HG12	1.85	0.57
1:C:15:ILE:N	1:C:15:ILE:CD1	2.69	0.56
1:A:166:ASN:OD1	1:A:168:PRO:HD2	2.06	0.55
1:C:188:LYS:CD	1:D:95:VAL:HG12	2.37	0.54
1:A:156:ILE:HD12	1:B:174:LEU:HD11	1.89	0.54
1:E:152:HIS:CD2	1:F:150:LEU:HD21	2.45	0.52
1:A:150:LEU:O	1:A:154:VAL:HG23	2.10	0.52
1:A:150:LEU:HD23	1:A:150:LEU:C	2.31	0.51
1:A:84:ILE:HD13	1:A:129:LEU:HD22	1.93	0.51
1:E:88:MSE:HB2	1:E:102:LYS:HB2	1.92	0.51
1:A:39:GLU:OE2	1:A:76:ARG:NH2	2.47	0.48
1:F:177:GLU:HA	1:F:177:GLU:OE1	2.14	0.47
1:B:199:GLU:O	1:B:203:ASN:HB2	2.15	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:88:MSE:HB2	1:F:102:LYS:HB2	1.97	0.46
1:C:95:VAL:HG12	1:C:95:VAL:O	2.15	0.46
1:D:100:GLU:OE1	1:D:102:LYS:HD2	2.16	0.46
1:E:156:ILE:HD12	1:F:174:LEU:HD11	1.98	0.46
1:F:32:TYR:HB3	1:F:74:LEU:HD21	1.97	0.46
1:D:119:LEU:C	1:D:119:LEU:HD23	2.37	0.45
1:E:202:ILE:O	1:E:203:ASN:C	2.56	0.45
1:B:165:MSE:HE3	1:C:162:GLN:HE22	1.81	0.44
1:A:202:ILE:O	1:A:203:ASN:C	2.56	0.44
1:D:156:ILE:HG21	1:E:174:LEU:HD21	2.00	0.44
1:B:203:ASN:C	2:B:244:HOH:O	2.55	0.44
1:E:166:ASN:OD1	1:E:168:PRO:HD2	2.17	0.44
1:F:62:LEU:HG	1:F:74:LEU:HD12	2.00	0.43
1:B:166:ASN:OD1	1:B:168:PRO:HD2	2.19	0.43
1:F:166:ASN:OD1	1:F:168:PRO:HD2	2.18	0.43
1:B:15:ILE:N	1:B:15:ILE:HD12	2.33	0.43
1:A:162:GLN:O	1:A:163:GLY:C	2.58	0.42
1:E:165:MSE:CE	1:F:162:GLN:HG3	2.49	0.42
1:E:119:LEU:HD23	1:E:119:LEU:C	2.39	0.42
1:B:143:LEU:HG	1:B:189:TRP:HB2	2.01	0.42
1:C:56:ASN:OD1	1:C:56:ASN:N	2.53	0.42
1:A:44:LEU:HD11	1:A:51:GLU:HG2	2.02	0.42
1:E:100:GLU:OE1	1:E:102:LYS:HD2	2.20	0.42
1:C:184:GLY:C	1:D:95:VAL:HG21	2.40	0.41
1:A:202:ILE:CG2	1:A:202:ILE:O	2.66	0.41
1:C:105:ILE:HG23	1:C:105:ILE:O	2.20	0.41
1:D:63:ASP:OD1	1:D:64:GLY:N	2.53	0.41
1:A:88:MSE:HB2	1:A:102:LYS:HB2	2.02	0.41
1:C:88:MSE:HB2	1:C:102:LYS:HB2	2.03	0.41
1:F:58:ALA:HB2	1:F:78:PRO:HA	2.02	0.41
1:C:170:THR:O	1:C:174:LEU:HB2	2.21	0.40
1:A:61:HIS:CE1	1:A:63:ASP:HB2	2.56	0.40
1:B:39:GLU:OE2	1:B:76:ARG:NH2	2.54	0.40
1:E:104:GLU:OE2	1:E:109:HIS:NE2	2.54	0.40
1:A:15:ILE:CD1	1:A:15:ILE:N	2.85	0.40
1:F:202:ILE:O	1:F:202:ILE:CG2	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/193 (95%)	181 (98%)	3 (2%)	0	100	100
1	B	179/193 (93%)	174 (97%)	3 (2%)	2 (1%)	14	19
1	C	178/193 (92%)	173 (97%)	3 (2%)	2 (1%)	14	19
1	D	178/193 (92%)	173 (97%)	4 (2%)	1 (1%)	25	35
1	E	177/193 (92%)	172 (97%)	4 (2%)	1 (1%)	25	35
1	F	181/193 (94%)	177 (98%)	4 (2%)	0	100	100
All	All	1077/1158 (93%)	1050 (98%)	21 (2%)	6 (1%)	25	35

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	159	GLY
1	D	160	LEU
1	C	159	GLY
1	E	159	GLY
1	C	161	ARG
1	B	158	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/166 (95%)	154 (98%)	3 (2%)	57	74
1	B	157/166 (95%)	154 (98%)	3 (2%)	57	74

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	153/166 (92%)	149 (97%)	4 (3%)	46	64
1	D	155/166 (93%)	153 (99%)	2 (1%)	69	83
1	E	149/166 (90%)	145 (97%)	4 (3%)	44	63
1	F	158/166 (95%)	156 (99%)	2 (1%)	69	83
All	All	929/996 (93%)	911 (98%)	18 (2%)	57	74

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	SER
1	A	138	GLU
1	A	199	GLU
1	B	47	SER
1	B	70	SER
1	B	174	LEU
1	C	15	ILE
1	C	47	SER
1	C	138	GLU
1	C	199	GLU
1	D	47	SER
1	D	138	GLU
1	E	47	SER
1	E	50	ASP
1	E	70	SER
1	E	138	GLU
1	F	47	SER
1	F	138	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	HIS
1	B	190	ASN
1	D	109	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/193 (95%)	0.24	4 (2%) 62 59	42, 65, 90, 103	0
1	B	181/193 (93%)	0.38	7 (3%) 39 37	42, 65, 88, 95	0
1	C	181/193 (93%)	0.31	6 (3%) 46 44	42, 65, 88, 94	0
1	D	181/193 (93%)	0.32	7 (3%) 39 37	43, 65, 87, 95	0
1	E	180/193 (93%)	0.59	19 (10%) 6 5	43, 65, 87, 95	0
1	F	180/193 (93%)	0.68	20 (11%) 5 4	43, 65, 87, 94	0
All	All	1088/1158 (93%)	0.42	63 (5%) 23 21	42, 65, 88, 103	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	202	ILE	6.3
1	E	203	ASN	5.2
1	F	22	ASP	5.0
1	F	97	LEU	4.8
1	E	121	HIS	4.4
1	E	89	PHE	4.2
1	D	203	ASN	4.1
1	B	203	ASN	4.0
1	F	25	ASP	4.0
1	E	43	PHE	3.9
1	E	122	VAL	3.9
1	E	97	LEU	3.7
1	C	20	HIS	3.7
1	F	95	VAL	3.6
1	E	19	ILE	3.6
1	D	97	LEU	3.5
1	E	44	LEU	3.5
1	A	67	ALA	3.4
1	D	29	ALA	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	23	ASP	3.3
1	F	96	ASP	3.2
1	D	62	LEU	3.2
1	E	119	LEU	3.1
1	B	22	ASP	3.0
1	B	21	PRO	3.0
1	E	95	VAL	2.9
1	F	89	PHE	2.9
1	C	44	LEU	2.9
1	F	108	LYS	2.8
1	C	22	ASP	2.8
1	B	20	HIS	2.8
1	E	22	ASP	2.8
1	F	62	LEU	2.8
1	E	33	VAL	2.8
1	F	28	ALA	2.7
1	B	160	LEU	2.7
1	F	75	TYR	2.7
1	B	70	SER	2.6
1	C	24	PHE	2.6
1	A	65	GLU	2.6
1	D	60	VAL	2.6
1	F	46	LYS	2.6
1	E	96	ASP	2.5
1	F	21	PRO	2.4
1	A	66	SER	2.4
1	E	70	SER	2.4
1	F	92	ALA	2.3
1	F	74	LEU	2.3
1	E	25	ASP	2.3
1	F	32	TYR	2.2
1	F	31	ASP	2.2
1	D	75	TYR	2.2
1	E	94	THR	2.2
1	C	14	ASP	2.1
1	C	15	ILE	2.1
1	E	24	PHE	2.1
1	D	160	LEU	2.1
1	F	93	GLY	2.1
1	E	124	ASP	2.0
1	A	29	ALA	2.0
1	F	27	ALA	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	52	TYR	2.0
1	B	24	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.