



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 10:53 am BST

PDB ID : 6B8V  
Title : Crystal structure of adenylyl-sulfate kinase from *Cryptococcus neoformans*  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2017-10-09  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>•</div> <div>22%</div> </div> </div>
1	B	211	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>•</div> <div>22%</div> </div> </div>
1	C	211	<div> <div>3%</div> <div> <div></div> <div>75%</div> <div>•</div> <div>22%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylylsulfate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	0	0
			1253	794	217	241	1			
1	B	164	Total	C	N	O	S	0	0	0
			1264	800	219	244	1			
1	C	164	Total	C	N	O	S	0	0	0
			1254	795	217	241	1			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A0A212GZX6
A	-6	ALA	-	expression tag	UNP A0A212GZX6
A	-5	HIS	-	expression tag	UNP A0A212GZX6
A	-4	HIS	-	expression tag	UNP A0A212GZX6
A	-3	HIS	-	expression tag	UNP A0A212GZX6
A	-2	HIS	-	expression tag	UNP A0A212GZX6
A	-1	HIS	-	expression tag	UNP A0A212GZX6
A	0	HIS	-	expression tag	UNP A0A212GZX6
B	-7	MET	-	expression tag	UNP A0A212GZX6
B	-6	ALA	-	expression tag	UNP A0A212GZX6
B	-5	HIS	-	expression tag	UNP A0A212GZX6
B	-4	HIS	-	expression tag	UNP A0A212GZX6
B	-3	HIS	-	expression tag	UNP A0A212GZX6
B	-2	HIS	-	expression tag	UNP A0A212GZX6
B	-1	HIS	-	expression tag	UNP A0A212GZX6
B	0	HIS	-	expression tag	UNP A0A212GZX6
C	-7	MET	-	expression tag	UNP A0A212GZX6
C	-6	ALA	-	expression tag	UNP A0A212GZX6
C	-5	HIS	-	expression tag	UNP A0A212GZX6
C	-4	HIS	-	expression tag	UNP A0A212GZX6
C	-3	HIS	-	expression tag	UNP A0A212GZX6
C	-2	HIS	-	expression tag	UNP A0A212GZX6
C	-1	HIS	-	expression tag	UNP A0A212GZX6

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	HIS	-	expression tag	UNP A0A212GZX6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	1
			12	6	6		
2	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total	O	0	0
			46	46		
3	B	50	Total	O	0	1
			51	51		
3	C	43	Total	O	0	0
			43	43		



- Molecule 1: Adenylylsulfate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.68Å 87.68Å 173.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.84 – 2.55 46.03 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.84-2.55) 99.9 (46.03-2.55)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.01 (at 2.54Å)	Xtriage
Refinement program	PHENIX (dev_2499: ???)	Depositor
R, $R_{free}$	0.184 , 0.240 0.184 , 0.240	Depositor DCC
$R_{free}$ test set	1854 reflections (7.34%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.5	Xtriage
Anisotropy	0.674	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.55$ , $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3929	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.8261e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.39	0/1276	0.53	0/1733
1	B	0.40	0/1287	0.54	0/1746
1	C	0.40	0/1277	0.53	0/1734
All	All	0.40	0/3840	0.54	0/5213

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1253	0	1236	6	0
1	B	1264	0	1255	3	0
1	C	1254	0	1238	3	0
2	A	12	0	16	1	0
2	C	6	0	8	0	0
3	A	46	0	0	3	0
3	B	51	0	0	2	0
3	C	43	0	0	1	0
All	All	3929	0	3753	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:17:ARG:NH2	3:C:401:HOH:O	2.21	0.65
1:B:17:ARG:NH1	3:B:301:HOH:O	2.34	0.60
1:A:51:HIS:ND1	3:A:401:HOH:O	2.36	0.45
1:A:17:ARG:NH1	3:A:402:HOH:O	2.42	0.44
1:B:37:GLY:HA2	3:B:338:HOH:O	2.18	0.42
1:B:83:ILE:HD13	1:B:111:ASP:HB3	2.01	0.42
1:A:55:HIS:CE1	2:A:300[A]:GOL:H2	2.55	0.41
1:A:37:GLY:HA2	3:A:437:HOH:O	2.19	0.41
1:A:25:GLY:HA3	1:A:94:ALA:O	2.21	0.41
1:A:83:ILE:HD13	1:A:111:ASP:HB3	2.03	0.40
1:C:75:ASP:OD1	1:C:75:ASP:N	2.51	0.40
1:C:175:ILE:HD11	1:C:194:TYR:CG	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/211 (76%)	157 (98%)	3 (2%)	0	100	100
1	B	160/211 (76%)	155 (97%)	5 (3%)	0	100	100
1	C	160/211 (76%)	157 (98%)	3 (2%)	0	100	100
All	All	480/633 (76%)	469 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/175 (75%)	131 (99%)	1 (1%)	81	88
1	B	135/175 (77%)	131 (97%)	4 (3%)	41	55
1	C	132/175 (75%)	131 (99%)	1 (1%)	81	88
All	All	399/525 (76%)	393 (98%)	6 (2%)	65	77

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ARG
1	B	112	ARG
1	B	172	ASN
1	B	179	THR
1	B	180	ASP
1	C	112	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	C	300	-	5,5,5	0.43	0	5,5,5	0.26	0
2	GOL	A	300[A]	-	5,5,5	0.37	0	5,5,5	0.24	0
2	GOL	A	300[B]	-	5,5,5	0.37	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	300	-	-	2/4/4/4	-
2	GOL	A	300[A]	-	-	0/4/4/4	-
2	GOL	A	300[B]	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	300[B]	GOL	O1-C1-C2-O2
2	A	300[B]	GOL	O1-C1-C2-C3
2	C	300	GOL	O1-C1-C2-C3
2	C	300	GOL	O1-C1-C2-O2
2	A	300[B]	GOL	O2-C2-C3-O3
2	A	300[B]	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300[A]	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	164/211 (77%)	0.03	5 (3%)	50 57	32, 45, 75, 98	0
1	B	164/211 (77%)	0.08	8 (4%)	29 35	31, 45, 70, 98	0
1	C	164/211 (77%)	0.08	7 (4%)	35 42	30, 47, 75, 104	0
All	All	492/633 (77%)	0.06	20 (4%)	37 44	30, 46, 75, 104	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	180	ASP	4.1
1	C	135	ALA	3.1
1	B	135	ALA	2.7
1	C	167	TYR	2.7
1	A	36	SER	2.5
1	B	136	PRO	2.5
1	A	135	ALA	2.4
1	A	180	ASP	2.3
1	C	34	SER	2.3
1	C	35	ALA	2.3
1	C	136	PRO	2.3
1	C	179	THR	2.2
1	B	36	SER	2.2
1	C	169	ALA	2.1
1	B	182	VAL	2.1
1	B	179	THR	2.1
1	A	35	ALA	2.1
1	B	172	ASN	2.1
1	A	37	GLY	2.0
1	B	33	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	300	6/6	0.93	0.18	50,61,65,65	6
2	GOL	A	300[A]	6/6	0.93	0.39	62,65,65,66	6
2	GOL	A	300[B]	6/6	0.93	0.39	64,64,65,66	6

## 6.5 Other polymers [i](#)

There are no such residues in this entry.