



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 03:53 pm BST

PDB ID : 4B9B
Title : The structure of the omega aminotransferase from Pseudomonas aeruginosa
Authors : Sayer, C.; Isupov, M.N.; Westlake, A.; Littlechild, J.A.
Deposited on : 2012-09-03
Resolution : 1.64 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>85%</div> <div>11% ..</div> </div>
1	B	448	<div> <div>88%</div> <div>9% .</div> </div>
1	C	448	<div> <div>87%</div> <div>10% ..</div> </div>
1	D	448	<div> <div>84%</div> <div>12% ..</div> </div>
1	E	448	<div> <div>88%</div> <div>9% .</div> </div>
1	F	448	<div> <div>87%</div> <div>9% ..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	448	 87% 9% . .
1	H	448	 86% 11% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLP	A	500	-	-	X	-
2	PLP	D	500	-	X	-	-
3	GOL	C	504	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 33210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-ALANINE-PYRUVATE TRANSAMINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	38	0
			3537	2261	615	644	17			
1	B	435	Total	C	N	O	S	0	41	0
			3549	2274	614	642	19			
1	C	436	Total	C	N	O	S	0	42	0
			3563	2276	620	647	20			
1	D	436	Total	C	N	O	S	0	47	0
			3607	2305	632	651	19			
1	E	435	Total	C	N	O	S	0	43	0
			3569	2280	623	647	19			
1	F	436	Total	C	N	O	S	0	37	0
			3536	2255	622	642	17			
1	G	436	Total	C	N	O	S	0	34	0
			3528	2249	619	641	19			
1	H	436	Total	C	N	O	S	0	39	0
			3547	2264	622	643	18			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	C	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	D	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	E	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	F	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	G	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	H	1	Total	C	N	O	P	0	0
			16	8	1	6	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		
3	G	1	Total	C	O	0	0
			6	3	3		
3	H	1	Total	C	O	0	0
			6	3	3		
3	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total 1	Cl 1	0	0
4	H	1	Total 1	Cl 1	0	0
4	B	1	Total 1	Cl 1	0	0
4	C	1	Total 1	Cl 1	0	0
4	A	1	Total 1	Cl 1	0	0
4	F	1	Total 1	Cl 1	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total 1	Ca 1	0	0
5	G	1	Total 1	Ca 1	0	0
5	F	1	Total 1	Ca 1	0	0
5	E	1	Total 1	Ca 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	654	Total 654	O 654	0	0
6	B	536	Total 536	O 536	0	0
6	C	718	Total 718	O 718	0	0
6	D	570	Total 570	O 570	0	0
6	E	608	Total 608	O 608	0	0
6	F	515	Total 515	O 515	0	0
6	G	532	Total 532	O 532	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	441	Total 441	O 441	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE




- Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE



- Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE



- Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE






• Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE

Chain E: 88% 9% .




• Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE

Chain F: 87% 9% . .



• Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE

Chain G: 87% 9% . .



• Molecule 1: BETA-ALANINE-PYRUVATE TRANSAMINASE

Chain H: 86% 11% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.38Å 133.16Å 161.96Å 90.00° 91.75° 90.00°	Depositor
Resolution (Å)	66.58 – 1.64 66.58 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.3 (66.58-1.64) 99.3 (66.58-1.64)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.174 , 0.219 0.173 , 0.217	Depositor DCC
R_{free} test set	20738 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	13.0	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	33210	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5891e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, CL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	4/3721 (0.1%)	0.85	8/5034 (0.2%)
1	B	0.85	3/3736 (0.1%)	0.85	2/5049 (0.0%)
1	C	0.89	2/3753 (0.1%)	0.85	4/5074 (0.1%)
1	D	0.86	2/3809 (0.1%)	0.84	3/5147 (0.1%)
1	E	0.88	3/3762 (0.1%)	0.87	4/5084 (0.1%)
1	F	0.91	3/3717 (0.1%)	0.90	6/5027 (0.1%)
1	G	0.84	1/3691 (0.0%)	0.83	3/4989 (0.1%)
1	H	0.82	2/3734 (0.1%)	0.83	1/5048 (0.0%)
All	All	0.87	20/29923 (0.1%)	0.85	31/40452 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	G	0	1
All	All	0	2

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	407	TRP	CD2-CE2	6.56	1.49	1.41
1	C	61	TRP	CD2-CE2	6.33	1.49	1.41
1	G	61	TRP	CD2-CE2	6.14	1.48	1.41
1	F	61	TRP	CD2-CE2	5.71	1.48	1.41
1	H	61	TRP	CD2-CE2	5.69	1.48	1.41

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	78[A]	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	F	78[B]	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	E	84	ASP	CB-CG-OD1	7.02	124.61	118.30
1	E	228	MET	CG-SD-CE	6.96	111.34	100.20
1	A	391	ARG	NE-CZ-NH2	-6.71	116.95	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	419	THR	Peptide
1	G	419	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3537	0	3600	55	0
1	B	3549	0	3632	55	0
1	C	3563	0	3627	59	0
1	D	3607	0	3691	56	0
1	E	3569	0	3642	41	0
1	F	3536	0	3591	45	0
1	G	3528	0	3573	44	0
1	H	3547	0	3612	51	0
2	A	16	0	7	6	0
2	B	16	0	7	1	0
2	C	16	0	7	1	0
2	D	16	0	7	3	0
2	E	16	0	7	1	0
2	F	16	0	7	1	0
2	G	16	0	7	4	0
2	H	16	0	7	1	0
3	A	12	0	16	1	0
3	B	6	0	8	3	0
3	C	12	0	16	7	0
3	E	6	0	8	2	0
3	F	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	6	0	8	1	0
3	H	12	0	16	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
6	A	654	0	0	27	0
6	B	536	0	0	37	0
6	C	718	0	0	38	0
6	D	570	0	0	29	0
6	E	608	0	0	27	0
6	F	515	0	0	20	0
6	G	532	0	0	29	0
6	H	441	0	0	26	0
All	All	33210	0	29104	411	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 411 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:70[A]:LYS:HE2	6:G:2092:HOH:O	1.23	1.35
1:H:52[A]:LYS:CE	6:H:2051:HOH:O	1.69	1.33
1:D:378[B]:ARG:NH2	1:D:387[B]:GLN:OE1	1.62	1.30
1:B:442[B]:GLU:OE2	6:B:2527:HOH:O	1.52	1.27
1:F:18[B]:ARG:NH2	6:F:2009:HOH:O	1.62	1.26

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/448 (105%)	453 (96%)	16 (3%)	3 (1%)	25	8
1	B	474/448 (106%)	458 (97%)	15 (3%)	1 (0%)	47	26
1	C	476/448 (106%)	459 (96%)	15 (3%)	2 (0%)	34	15
1	D	481/448 (107%)	466 (97%)	11 (2%)	4 (1%)	19	4
1	E	476/448 (106%)	462 (97%)	10 (2%)	4 (1%)	19	4
1	F	471/448 (105%)	454 (96%)	14 (3%)	3 (1%)	25	8
1	G	468/448 (104%)	448 (96%)	17 (4%)	3 (1%)	25	8
1	H	473/448 (106%)	456 (96%)	14 (3%)	3 (1%)	25	8
All	All	3791/3584 (106%)	3656 (96%)	112 (3%)	23 (1%)	22	8

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	LYS
1	C	288	LYS
1	D	288	LYS
1	G	288	LYS
1	H	288	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	376/350 (107%)	369 (98%)	7 (2%)	57 32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	379/350 (108%)	375 (99%)	4 (1%)	73	55
1	C	380/350 (109%)	373 (98%)	7 (2%)	59	34
1	D	386/350 (110%)	378 (98%)	8 (2%)	53	26
1	E	381/350 (109%)	377 (99%)	4 (1%)	76	59
1	F	375/350 (107%)	371 (99%)	4 (1%)	73	55
1	G	373/350 (107%)	365 (98%)	8 (2%)	53	26
1	H	377/350 (108%)	368 (98%)	9 (2%)	49	21
All	All	3027/2800 (108%)	2976 (98%)	51 (2%)	73	36

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	228[B]	MET
1	E	405[A]	LYS
1	H	408[A]	GLN
1	D	253[B]	ASN
1	E	405[B]	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	373	ASN
1	F	409	GLN
1	H	252	HIS
1	E	169	ASN
1	F	79	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 12 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GOL	C	504	-	5,5,5	0.31	0	5,5,5	1.07	0
2	PLP	F	500	-	16,16,16	2.74	5 (31%)	20,23,23	2.46	10 (50%)
3	GOL	H	501	-	5,5,5	0.52	0	5,5,5	0.60	0
2	PLP	D	500	-	16,16,16	2.70	6 (37%)	20,23,23	2.16	9 (45%)
3	GOL	G	504	-	5,5,5	0.43	0	5,5,5	1.29	0
3	GOL	F	501	-	5,5,5	0.65	0	5,5,5	0.68	0
3	GOL	H	504	-	5,5,5	0.28	0	5,5,5	0.45	0
3	GOL	A	501	-	5,5,5	0.58	0	5,5,5	0.39	0
2	PLP	C	500	-	16,16,16	3.06	6 (37%)	20,23,23	2.43	8 (40%)
3	GOL	C	501	-	5,5,5	0.38	0	5,5,5	1.06	1 (20%)
2	PLP	A	500	-	16,16,16	3.40	4 (25%)	20,23,23	2.26	9 (45%)
2	PLP	G	500	-	16,16,16	2.94	5 (31%)	20,23,23	2.24	7 (35%)
3	GOL	B	504	-	5,5,5	0.41	0	5,5,5	1.34	1 (20%)
2	PLP	E	500	-	16,16,16	2.94	5 (31%)	20,23,23	2.06	8 (40%)
2	PLP	B	500	-	16,16,16	2.85	7 (43%)	20,23,23	1.82	6 (30%)
2	PLP	H	500	-	16,16,16	2.95	4 (25%)	20,23,23	1.66	6 (30%)
3	GOL	E	504	-	5,5,5	0.48	0	5,5,5	0.62	0
3	GOL	A	504	-	5,5,5	0.30	0	5,5,5	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	504	-	-	0/4/4/4	-
2	PLP	F	500	-	-	1/8/8/8	0/1/1/1
3	GOL	H	501	-	-	0/4/4/4	-
2	PLP	D	500	-	-	5/8/8/8	0/1/1/1
3	GOL	G	504	-	-	1/4/4/4	-
3	GOL	F	501	-	-	2/4/4/4	-
3	GOL	H	504	-	-	2/4/4/4	-
3	GOL	A	501	-	-	0/4/4/4	-
2	PLP	C	500	-	-	5/8/8/8	0/1/1/1
3	GOL	C	501	-	-	2/4/4/4	-
2	PLP	A	500	-	-	2/8/8/8	0/1/1/1
2	PLP	G	500	-	-	2/8/8/8	0/1/1/1
3	GOL	B	504	-	-	2/4/4/4	-
2	PLP	E	500	-	-	3/8/8/8	0/1/1/1
2	PLP	B	500	-	-	3/8/8/8	0/1/1/1
2	PLP	H	500	-	-	4/8/8/8	0/1/1/1
3	GOL	E	504	-	-	2/4/4/4	-
3	GOL	A	504	-	-	4/4/4/4	-

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	PLP	C3-C2	10.35	1.51	1.40
2	G	500	PLP	C3-C2	8.03	1.48	1.40
2	E	500	PLP	C3-C2	7.71	1.48	1.40
2	H	500	PLP	C3-C2	6.86	1.47	1.40
2	H	500	PLP	C4-C5	6.73	1.50	1.42

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	500	PLP	O4P-P-O1P	5.96	123.20	106.47
2	C	500	PLP	C6-N1-C2	5.06	128.53	119.17
2	C	500	PLP	O3P-P-O4P	4.98	119.98	106.73
2	D	500	PLP	O3P-P-O4P	4.86	119.66	106.73
2	E	500	PLP	O2P-P-O4P	4.52	118.76	106.73

There are no chirality outliers.

5 of 40 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	501	GOL	O1-C1-C2-C3
2	C	500	PLP	C5A-O4P-P-O2P
2	C	500	PLP	C5A-O4P-P-O3P
3	H	504	GOL	O1-C1-C2-C3
3	C	501	GOL	C1-C2-C3-O3

There are no ring outliers.

14 monomers are involved in 33 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	504	GOL	7	0
2	F	500	PLP	1	0
2	D	500	PLP	3	0
3	G	504	GOL	1	0
3	H	504	GOL	1	0
2	C	500	PLP	1	0
2	A	500	PLP	6	0
2	G	500	PLP	4	0
3	B	504	GOL	3	0
2	E	500	PLP	1	0
2	B	500	PLP	1	0
2	H	500	PLP	1	0
3	E	504	GOL	2	0
3	A	504	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/448 (97%)	-0.23	0 100 100	7, 13, 24, 64	0
1	B	435/448 (97%)	-0.37	1 (0%) 95 94	7, 14, 22, 64	0
1	C	436/448 (97%)	-0.43	0 100 100	6, 11, 21, 63	0
1	D	436/448 (97%)	-0.37	1 (0%) 95 94	6, 12, 21, 76	0
1	E	435/448 (97%)	-0.46	1 (0%) 95 94	7, 11, 20, 75	0
1	F	436/448 (97%)	-0.24	2 (0%) 91 91	6, 12, 22, 76	0
1	G	436/448 (97%)	-0.26	1 (0%) 95 94	8, 15, 26, 83	0
1	H	436/448 (97%)	-0.10	4 (0%) 84 85	7, 16, 32, 91	0
All	All	3486/3584 (97%)	-0.31	10 (0%) 94 94	6, 13, 25, 91	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	13	SER	4.2
1	H	13	SER	4.0
1	D	13	SER	4.0
1	H	367	GLY	3.0
1	E	14	GLU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	G	504	6/6	0.91	0.15	18,27,38,46	0
2	PLP	B	500	16/16	0.91	0.16	8,17,21,37	16
2	PLP	A	500	16/16	0.91	0.18	9,16,23,28	16
3	GOL	H	501	6/6	0.92	0.12	18,20,25,29	0
3	GOL	C	504	6/6	0.92	0.15	13,19,31,32	0
2	PLP	D	500	16/16	0.92	0.17	6,15,20,30	16
2	PLP	G	500	16/16	0.93	0.17	9,17,24,46	16
2	PLP	F	500	16/16	0.93	0.15	7,14,18,23	16
3	GOL	H	504	6/6	0.93	0.17	14,21,29,30	0
2	PLP	C	500	16/16	0.93	0.16	5,13,18,28	16
3	GOL	F	501	6/6	0.93	0.15	15,22,30,34	0
3	GOL	C	501	6/6	0.94	0.15	17,25,32,52	0
2	PLP	E	500	16/16	0.94	0.18	6,14,18,29	16
2	PLP	H	500	16/16	0.94	0.15	11,19,25,33	16
3	GOL	A	501	6/6	0.95	0.11	13,14,20,25	0
3	GOL	E	504	6/6	0.95	0.12	12,18,28,35	0
3	GOL	A	504	6/6	0.95	0.16	14,27,38,43	0
3	GOL	B	504	6/6	0.96	0.15	15,32,38,39	0
4	CL	G	502	1/1	0.96	0.05	22,22,22,22	0
4	CL	C	502	1/1	0.98	0.06	13,13,13,13	0
5	CA	H	503	1/1	0.99	0.05	19,19,19,19	0
4	CL	E	502	1/1	0.99	0.04	18,18,18,18	0
4	CL	F	502	1/1	0.99	0.04	14,14,14,14	0
4	CL	B	502	1/1	0.99	0.04	18,18,18,18	0
4	CL	D	502	1/1	0.99	0.05	18,18,18,18	0
5	CA	E	503	1/1	0.99	0.04	17,17,17,17	0
4	CL	H	502	1/1	0.99	0.05	16,16,16,16	0
4	CL	A	502	1/1	0.99	0.05	14,14,14,14	0
5	CA	F	503	1/1	1.00	0.09	18,18,18,18	0
5	CA	G	503	1/1	1.00	0.06	20,20,20,20	0

6.5 Other polymers ⓘ

There are no such residues in this entry.