



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:24 pm BST

PDB ID : 1BD8
Title : STRUCTURE OF CDK INHIBITOR P19INK4D
Authors : Baumgartner, R.; Fernandez-Catalan, C.; Winoto, A.; Huber, R.; Engh, R.;
Holak, T.A.
Deposited on : 1998-05-12
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	156	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called P19INK4D CDK4/6 INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	0	0
			1168	721	222	222	3			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	195	Total	O	14	0
			195	195		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: P19INK4D CDK4/6 INHIBITOR

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	28.47 Å 54.56 Å 45.59 Å 90.00° 90.16° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.80)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.190 , 0.260	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1363	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	3/1185 (0.3%)	1.85	21/1605 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	162	MET	C-OXT	-20.50	0.84	1.23
1	A	23	GLU	CD-OE2	5.97	1.32	1.25
1	A	11	ARG	CD-NE	-5.23	1.37	1.46

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	ARG	NE-CZ-NH2	-36.24	102.18	120.30
1	A	149	ARG	NE-CZ-NH1	20.35	130.47	120.30
1	A	11	ARG	CD-NE-CZ	19.21	150.49	123.60
1	A	149	ARG	CD-NE-CZ	14.42	143.79	123.60
1	A	145	LEU	CA-CB-CG	13.86	147.18	115.30
1	A	11	ARG	CG-CD-NE	11.54	136.04	111.80
1	A	146	ALA	CA-C-N	9.66	138.46	117.20
1	A	36	ASP	CB-CG-OD1	9.47	126.83	118.30
1	A	146	ALA	O-C-N	-8.90	108.45	122.70
1	A	80	ASP	CB-CG-OD1	8.00	125.50	118.30
1	A	11	ARG	NE-CZ-NH2	-7.87	116.36	120.30
1	A	147	LEU	N-CA-CB	7.72	125.85	110.40
1	A	36	ASP	CA-CB-CG	7.50	129.90	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	ARG	NH1-CZ-NH2	7.21	127.33	119.40
1	A	58	LEU	CB-CG-CD2	-7.01	99.08	111.00
1	A	10	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	149	ARG	CG-CD-NE	-5.95	99.31	111.80
1	A	11	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	A	71	ASP	CB-CG-OD2	-5.72	113.15	118.30
1	A	26	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	36	ASP	OD1-CG-OD2	-5.02	113.75	123.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	128	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1168	0	1171	37	5
2	A	195	0	0	18	11
All	All	1363	0	1171	37	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ALA:O	1:A:129:GLU:HG3	1.75	0.86
1:A:43:LYS:HE3	2:A:323:HOH:O	1.81	0.79
1:A:84:THR:O	2:A:243:HOH:O	2.06	0.73
1:A:78:VAL:HG21	1:A:100:VAL:HG12	1.75	0.69
1:A:146:ALA:O	1:A:147:LEU:HB2	1.94	0.67
1:A:161:HIS:O	1:A:162:MET:HB2	1.96	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:HIS:HB3	2:A:252:HOH:O	2.00	0.61
1:A:7:ARG:HG3	1:A:11:ARG:HD3	1.83	0.61
1:A:129:GLU:OE1	2:A:331:HOH:O	2.16	0.60
1:A:146:ALA:O	1:A:147:LEU:CB	2.47	0.59
1:A:145:LEU:O	1:A:149:ARG:HG3	2.03	0.59
1:A:11:ARG:NH1	2:A:171:HOH:O	2.36	0.58
1:A:104:ASP:OD2	2:A:178:HOH:O	2.18	0.56
1:A:109:LEU:HB3	2:A:163:HOH:O	2.06	0.56
1:A:140:LEU:HB2	1:A:145:LEU:HD23	1.88	0.55
1:A:145:LEU:HB3	2:A:349:HOH:O	2.06	0.54
1:A:78:VAL:HG23	1:A:90:LEU:CD1	2.38	0.54
1:A:87:LEU:HB2	2:A:332:HOH:O	2.07	0.53
1:A:128:ALA:HB2	1:A:161:HIS:CE1	2.43	0.53
1:A:77:PRO:HD3	2:A:172:HOH:O	2.09	0.52
1:A:112:HIS:HB3	2:A:234:HOH:O	2.10	0.50
1:A:138:ARG:HG2	1:A:138:ARG:HH11	1.76	0.50
1:A:143:LEU:O	1:A:146:ALA:O	2.31	0.49
1:A:149:ARG:CZ	1:A:149:ARG:HB3	2.41	0.47
1:A:60:LEU:HD13	2:A:166:HOH:O	2.13	0.47
1:A:71:ASP:OD2	2:A:324:HOH:O	2.20	0.47
1:A:21:VAL:HG23	2:A:306:HOH:O	2.16	0.44
1:A:76:SER:HA	2:A:172:HOH:O	2.18	0.44
1:A:78:VAL:HG23	1:A:90:LEU:HD11	2.00	0.44
1:A:26:ARG:HD3	2:A:253:HOH:O	2.18	0.43
1:A:7:ARG:HH11	1:A:7:ARG:HD3	1.72	0.43
1:A:58:LEU:HD23	1:A:59:GLU:N	2.34	0.42
1:A:46:LEU:HB3	2:A:169:HOH:O	2.19	0.42
1:A:34:HIS:CG	1:A:35:PRO:HD2	2.55	0.42
1:A:161:HIS:O	1:A:162:MET:CB	2.65	0.41
1:A:12:LEU:HA	1:A:27:LEU:HD23	2.02	0.41
1:A:79:HIS:CE1	2:A:197:HOH:O	2.73	0.40

All (11) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:187:HOH:O	2:A:238:HOH:O[1_455]	1.39	0.81
1:A:149:ARG:NH2	2:A:170:HOH:O[1_455]	1.49	0.71
2:A:188:HOH:O	2:A:274:HOH:O[1_655]	1.56	0.64
1:A:117:GLU:O	2:A:235:HOH:O[1_455]	1.87	0.33

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:304:HOH:O	2:A:356:HOH:O[1_656]	1.95	0.25
1:A:91:LYS:NZ	2:A:248:HOH:O[2_656]	1.96	0.24
2:A:322:HOH:O	2:A:346:HOH:O[1_455]	1.99	0.21
1:A:149:ARG:CZ	2:A:170:HOH:O[1_455]	2.00	0.20
2:A:272:HOH:O	2:A:302:HOH:O[1_554]	2.06	0.14
2:A:207:HOH:O	2:A:250:HOH:O[1_655]	2.08	0.12
1:A:156:ASP:OD1	2:A:222:HOH:O[1_454]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	154/156 (99%)	149 (97%)	3 (2%)	2 (1%)	12 3

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	GLU
1	A	147	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	121/121 (100%)	112 (93%)	9 (7%)	13 4

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	11	ARG
1	A	27	LEU
1	A	50	MET
1	A	58	LEU
1	A	135	ARG
1	A	148	GLN
1	A	149	ARG
1	A	162	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	119	HIS
1	A	148	GLN
1	A	159	GLN
1	A	161	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.