



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:43 am BST

PDB ID : 2BFM  
Title : Leishmania major pteridine reductase 1 in complex with NADP and trimethoprim  
Authors : Schuettelkopf, A.W.; Hunter, W.N.  
Deposited on : 2004-12-09  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>10%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	288	<div> <div>5%</div> <div> <div></div> <div>79%</div> <div>12%</div> <div>•</div> <div>8%</div> </div> </div>
1	C	288	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>•</div> <div>12%</div> </div> </div>
1	D	288	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>•</div> <div>11%</div> </div> </div>

2 Entry composition ⓘ


There are 4 unique types of molecules in this entry. The entry contains 8188 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PTERIDINE REDUCTASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	0
			1984	1250	353	370	11			
1	B	266	Total	C	N	O	S	0	0	0
			1987	1253	354	369	11			
1	C	254	Total	C	N	O	S	0	0	0
			1892	1192	340	350	10			
1	D	256	Total	C	N	O	S	0	0	0
			1906	1201	342	353	10			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	
			48	21	7	17	3	
2	D	1	Total	C	N	O	P	
			48	21	7	17	3	

- Molecule 3 is TRIMETHOPRIM (three-letter code: TOP) (formula: C<sub>14</sub>H<sub>18</sub>N<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O		
			21	14	4	3		
3	B	1	Total	C	N	O		
			21	14	4	3		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	60	Total	O		
			60	60	0	0
4	B	45	Total	O		
			45	45	0	0
4	C	44	Total	O		
			44	44	0	0
4	D	36	Total	O		
			36	36	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: PTERIDINE REDUCTASE 1

Chain A: 





#### • Molecule 1: PTERIDINE REDUCTASE 1

Chain B: 




#### • Molecule 1: PTERIDINE REDUCTASE 1


Chain C: 



#### • Molecule 1: PTERIDINE REDUCTASE 1

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.70Å 104.44Å 138.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.50 – 2.60 24.76 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.50-2.60) 97.7 (24.76-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.23 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.253 , 0.288 0.253 , 0.287	Depositor DCC
$R_{free}$ test set	2085 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	8188	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0083e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, TOP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/2023	0.85	6/2755 (0.2%)
1	B	0.51	0/2026	0.77	4/2760 (0.1%)
1	C	0.47	0/1925	0.76	5/2616 (0.2%)
1	D	0.49	0/1940	0.74	3/2638 (0.1%)
All	All	0.50	0/7914	0.78	18/10769 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	ASP	CB-CG-OD2	14.62	131.45	118.30
1	A	104	ASP	OD1-CG-OD2	-11.04	102.33	123.30
1	A	104	ASP	CB-CG-OD1	8.80	126.22	118.30
1	B	104	ASP	CB-CG-OD2	8.48	125.94	118.30
1	C	104	ASP	CB-CG-OD2	8.34	125.81	118.30
1	D	104	ASP	CB-CG-OD2	7.41	124.97	118.30
1	C	259	ASP	CB-CG-OD2	7.16	124.75	118.30
1	B	142	ASP	CB-CG-OD2	6.73	124.36	118.30
1	A	259	ASP	CB-CG-OD2	6.58	124.22	118.30
1	D	259	ASP	CB-CG-OD2	6.03	123.72	118.30
1	D	65	ASP	CB-CG-OD2	5.63	123.37	118.30
1	B	65	ASP	CB-CG-OD2	5.50	123.25	118.30
1	B	104	ASP	OD1-CG-OD2	-5.48	112.89	123.30
1	C	251	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	231	ASP	CB-CG-OD2	5.17	122.95	118.30
1	C	104	ASP	OD1-CG-OD2	-5.16	113.50	123.30
1	C	142	ASP	CB-CG-OD2	5.03	122.83	118.30
1	A	65	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.



There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1984	0	1990	17	0
1	B	1987	0	1998	21	0
1	C	1892	0	1914	13	0
1	D	1906	0	1928	14	0
2	A	48	0	26	0	0
2	B	48	0	26	0	0
2	C	48	0	26	0	0
2	D	48	0	26	0	0
3	A	21	0	18	0	0
3	B	21	0	18	0	0
4	A	60	0	0	1	0
4	B	45	0	0	2	0
4	C	44	0	0	0	0
4	D	36	0	0	3	0
All	All	8188	0	7970	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:12:THR:HA	1:C:36:HIS:HB3	1.69	0.73
1:B:168:LYS:H	1:B:168:LYS:HD3	1.56	0.70
1:B:166:PRO:HB3	1:B:168:LYS:NZ	2.08	0.68
1:B:156:LYS:HE2	4:B:2017:HOH:O	1.95	0.67
1:B:166:PRO:HB3	1:B:168:LYS:HZ3	1.62	0.63
1:A:120:ARG:NH1	1:B:160:HIS:HB2	2.16	0.61
1:A:20:ARG:HE	1:A:47:LEU:HD13	1.65	0.60
1:B:229:LEU:HD23	1:B:238:TRP:CD1	2.36	0.60
1:A:120:ARG:CZ	1:B:160:HIS:HB2	2.32	0.58
1:B:12:THR:HA	1:B:36:HIS:HB3	1.85	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:ARG:HH11	1:C:39:ARG:HG2	1.69	0.57
1:C:133:ARG:O	1:C:136:MET:HG2	2.03	0.57
1:A:120:ARG:N	1:A:120:ARG:HD2	2.20	0.57
1:A:252:SER:HB3	1:C:271:TYR:CE2	2.40	0.57
1:A:242:ARG:HB2	1:A:250:ARG:HA	1.86	0.57
1:C:81:ALA:HB3	1:C:82:PRO:HD3	1.87	0.56
1:D:181:ASP:HB3	1:D:184:THR:HG23	1.87	0.55
1:B:73:PRO:HD3	1:B:82:PRO:HA	1.86	0.55
1:C:181:ASP:HB3	1:C:184:THR:HG23	1.87	0.55
1:A:134:GLU:C	1:A:136:MET:H	2.12	0.53
1:D:140:THR:O	1:D:144:PHE:HB2	2.09	0.53
1:A:271:TYR:CE2	1:C:252:SER:HB3	2.44	0.52
1:A:73:PRO:HA	1:A:81:ALA:O	2.09	0.52
4:B:2025:HOH:O	1:C:285:LEU:HD23	2.09	0.52
1:D:161:ARG:NH1	4:D:2013:HOH:O	2.42	0.52
1:D:156:LYS:HG2	1:D:160:HIS:HE1	1.75	0.52
1:C:150:ALA:HB3	1:C:151:PRO:HD3	1.91	0.52
1:A:181:ASP:HB3	1:A:184:THR:HG23	1.92	0.50
1:D:150:ALA:HB3	1:D:151:PRO:HD3	1.93	0.50
1:B:252:SER:HB3	1:D:271:TYR:CE2	2.45	0.50
1:A:20:ARG:NH2	4:A:2001:HOH:O	2.43	0.50
1:B:73:PRO:HG3	1:B:81:ALA:O	2.12	0.49
1:B:168:LYS:H	1:B:168:LYS:CD	2.19	0.48
1:B:271:TYR:CE2	1:D:252:SER:HB3	2.49	0.47
1:B:166:PRO:CB	1:B:168:LYS:HZ3	2.27	0.47
1:C:39:ARG:NH1	1:C:39:ARG:HG2	2.28	0.47
1:D:156:LYS:HG2	1:D:160:HIS:CE1	2.50	0.47
1:B:181:ASP:HB3	1:B:184:THR:HG23	1.98	0.46
1:C:140:THR:O	1:C:144:PHE:HB2	2.16	0.45
1:D:71:THR:O	1:D:83:VAL:N	2.47	0.45
1:C:242:ARG:H	1:C:242:ARG:HG3	1.59	0.45
1:C:160:HIS:HB2	4:D:2012:HOH:O	2.16	0.45
1:B:145:GLY:HA2	1:B:149:ILE:HB	1.97	0.45
1:D:12:THR:HA	1:D:36:HIS:HB3	1.99	0.45
1:B:119:LEU:O	1:B:121:ASN:N	2.49	0.45
1:A:120:ARG:CD	1:A:120:ARG:N	2.81	0.44
1:B:140:THR:O	1:B:144:PHE:HB2	2.17	0.44
1:B:151:PRO:O	1:B:155:ILE:HG13	2.18	0.44
1:D:70:ALA:HB1	1:D:82:PRO:HB2	1.99	0.43
1:B:238:TRP:CE2	1:B:242:ARG:HD2	2.54	0.43
1:A:133:ARG:HA	1:A:136:MET:HB3	1.99	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:LEU:HB3	1:B:120:ARG:H	1.52	0.42
1:D:244:LYS:HA	1:D:244:LYS:HD3	1.59	0.42
1:A:150:ALA:HB3	1:A:151:PRO:HD3	2.02	0.42
1:A:12:THR:HA	1:A:36:HIS:HB3	2.01	0.42
1:D:148:ALA:C	1:D:151:PRO:HD2	2.41	0.41
1:D:256:GLU:HA	4:D:2029:HOH:O	2.21	0.41
1:A:120:ARG:CD	1:A:120:ARG:H	2.33	0.41
1:A:140:THR:O	1:A:144:PHE:HB2	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/288 (90%)	241 (93%)	18 (7%)	1 (0%)	34	57
1	B	260/288 (90%)	242 (93%)	14 (5%)	4 (2%)	10	21
1	C	246/288 (85%)	234 (95%)	10 (4%)	2 (1%)	19	39
1	D	248/288 (86%)	235 (95%)	12 (5%)	1 (0%)	34	57
All	All	1014/1152 (88%)	952 (94%)	54 (5%)	8 (1%)	19	39

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	ARG
1	B	136	MET
1	A	180	VAL
1	B	180	VAL
1	C	228	VAL
1	D	180	VAL
1	C	180	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	73	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/223 (92%)	195 (95%)	11 (5%)	22	45
1	B	207/223 (93%)	196 (95%)	11 (5%)	22	45
1	C	196/223 (88%)	187 (95%)	9 (5%)	27	51
1	D	198/223 (89%)	184 (93%)	14 (7%)	14	29
All	All	807/892 (90%)	762 (94%)	45 (6%)	21	42

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	16	LYS
1	A	132	ASP
1	A	134	GLU
1	A	136	MET
1	A	144	PHE
1	A	168	LYS
1	A	242	ARG
1	A	262	ILE
1	A	270	LYS
1	A	275	THR
1	B	16	LYS
1	B	62	VAL
1	B	120	ARG
1	B	133	ARG
1	B	144	PHE
1	B	168	LYS
1	B	228	VAL
1	B	230	VAL
1	B	262	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	268	LYS
1	B	275	THR
1	C	112	SER
1	C	133	ARG
1	C	134	GLU
1	C	144	PHE
1	C	228	VAL
1	C	241	HIS
1	C	242	ARG
1	C	262	ILE
1	C	275	THR
1	D	5	THR
1	D	16	LYS
1	D	20	ARG
1	D	36	HIS
1	D	133	ARG
1	D	144	PHE
1	D	156	LYS
1	D	169	HIS
1	D	228	VAL
1	D	241	HIS
1	D	242	ARG
1	D	262	ILE
1	D	270	LYS
1	D	275	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	216	GLN
1	B	68	ASN
1	B	216	GLN
1	C	216	GLN
1	D	216	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDP	C	1289	-	45,52,52	1.71	5 (11%)	53,80,80	1.11	2 (3%)
2	NDP	D	1289	-	45,52,52	1.71	5 (11%)	53,80,80	1.12	1 (1%)
2	NDP	A	1289	-	45,52,52	1.74	5 (11%)	53,80,80	1.10	1 (1%)
2	NDP	B	1289	-	45,52,52	1.70	5 (11%)	53,80,80	1.07	1 (1%)
3	TOP	A	1290	-	22,22,22	2.42	3 (13%)	30,30,30	2.19	13 (43%)
3	TOP	B	1290	-	22,22,22	2.46	4 (18%)	30,30,30	2.32	13 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	C	1289	-	-	2/30/77/77	0/5/5/5
2	NDP	D	1289	-	-	2/30/77/77	0/5/5/5
2	NDP	A	1289	-	-	2/30/77/77	0/5/5/5
2	NDP	B	1289	-	-	4/30/77/77	0/5/5/5
3	TOP	A	1290	-	-	2/10/10/10	0/2/2/2
3	TOP	B	1290	-	-	2/10/10/10	0/2/2/2

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1289	NDP	O7N-C7N	7.03	1.41	1.24
2	A	1289	NDP	O7N-C7N	7.02	1.41	1.24
2	B	1289	NDP	O7N-C7N	6.98	1.41	1.24
2	C	1289	NDP	O7N-C7N	6.94	1.40	1.24
3	A	1290	TOP	O13-C12	-6.73	1.26	1.37
3	B	1290	TOP	O13-C12	-6.52	1.26	1.37
3	B	1290	TOP	O19-C18	-6.44	1.26	1.37
3	A	1290	TOP	O19-C18	-6.23	1.27	1.37
3	B	1290	TOP	O16-C15	-6.15	1.27	1.38
3	A	1290	TOP	O16-C15	-5.84	1.27	1.38
2	B	1289	NDP	C4N-C3N	-5.49	1.39	1.49
2	A	1289	NDP	C4N-C3N	-5.35	1.39	1.49
2	C	1289	NDP	C4N-C3N	-5.26	1.39	1.49
2	D	1289	NDP	C4N-C3N	-5.18	1.39	1.49
2	C	1289	NDP	C2A-N3A	4.08	1.38	1.32
2	D	1289	NDP	C2A-N3A	3.78	1.38	1.32
2	B	1289	NDP	C2A-N3A	3.77	1.38	1.32
2	A	1289	NDP	C2A-N3A	3.64	1.38	1.32
2	A	1289	NDP	C6N-C5N	3.45	1.39	1.33
2	D	1289	NDP	C6N-C5N	3.35	1.39	1.33
2	C	1289	NDP	C6N-C5N	3.17	1.39	1.33
2	B	1289	NDP	C6N-C5N	2.89	1.38	1.33
2	C	1289	NDP	C2A-N1A	2.89	1.39	1.33
2	A	1289	NDP	C2A-N1A	2.60	1.38	1.33
2	B	1289	NDP	C2A-N1A	2.40	1.38	1.33
2	D	1289	NDP	C2A-N1A	2.31	1.38	1.33
3	B	1290	TOP	C8-C6	-2.02	1.39	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1289	NDP	N3A-C2A-N1A	-5.74	119.70	128.68
2	A	1289	NDP	N3A-C2A-N1A	-5.64	119.86	128.68
2	B	1289	NDP	N3A-C2A-N1A	-5.63	119.89	128.68
2	C	1289	NDP	N3A-C2A-N1A	-5.56	119.98	128.68
3	B	1290	TOP	N2-C3-N5	-4.64	120.06	125.70
3	A	1290	TOP	N2-C3-N5	-4.27	120.50	125.70
3	B	1290	TOP	C14-O13-C12	4.15	123.80	117.53
3	B	1290	TOP	O13-C12-C15	3.89	122.00	115.16
3	A	1290	TOP	C20-O19-C18	3.55	122.88	117.53
3	A	1290	TOP	C17-O16-C15	3.55	124.50	114.78
3	A	1290	TOP	O19-C18-C15	3.45	121.22	115.16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1290	TOP	O13-C12-C11	-3.44	118.20	124.12
3	A	1290	TOP	O13-C12-C11	-3.43	118.22	124.12
3	A	1290	TOP	O13-C12-C15	3.36	121.07	115.16
3	B	1290	TOP	C20-O19-C18	3.28	122.48	117.53
3	B	1290	TOP	O19-C18-C15	3.25	120.87	115.16
3	A	1290	TOP	C8-C1-N2	-3.24	118.42	123.82
3	B	1290	TOP	C17-O16-C15	3.19	123.52	114.78
3	B	1290	TOP	N4-C3-N2	3.12	120.60	117.44
3	A	1290	TOP	N4-C3-N2	3.08	120.57	117.44
3	B	1290	TOP	C8-C1-N2	-3.01	118.81	123.82
3	B	1290	TOP	C3-N5-C6	2.94	120.23	116.99
3	A	1290	TOP	C14-O13-C12	2.89	121.89	117.53
3	A	1290	TOP	C1-N2-C3	2.86	121.74	116.44
3	B	1290	TOP	C1-N2-C3	2.77	121.56	116.44
3	B	1290	TOP	C1-C8-C6	2.75	119.46	115.72
3	A	1290	TOP	O19-C18-C21	-2.56	119.72	124.12
3	A	1290	TOP	C3-N5-C6	2.51	119.76	116.99
3	A	1290	TOP	C1-C8-C6	2.42	119.02	115.72
2	C	1289	NDP	C4A-C5A-N7A	-2.35	106.94	109.40
3	B	1290	TOP	O19-C18-C21	-2.33	120.10	124.12

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1290	TOP	C15-C12-O13-C14
2	C	1289	NDP	O4D-C1D-N1N-C6N
2	D	1289	NDP	O4D-C1D-N1N-C6N
2	A	1289	NDP	O4D-C1D-N1N-C6N
2	B	1289	NDP	O4D-C1D-N1N-C6N
3	B	1290	TOP	C11-C12-O13-C14
3	A	1290	TOP	C11-C12-O13-C14
3	B	1290	TOP	C15-C12-O13-C14
2	B	1289	NDP	C2B-O2B-P2B-O3X
2	C	1289	NDP	C2N-C3N-C7N-N7N
2	D	1289	NDP	C2N-C3N-C7N-N7N
2	A	1289	NDP	C2N-C3N-C7N-N7N
2	B	1289	NDP	C2N-C3N-C7N-N7N
2	B	1289	NDP	O4D-C4D-C5D-O5D


There are no ring outliers.


No monomer is involved in short contacts.




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## Ligand NDP B 1289



## Ligand TOP A 1290





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	266/288 (92%)	0.05	11 (4%)	37 30	17, 27, 42, 74	1 (0%)
1	B	266/288 (92%)	0.17	13 (4%)	29 23	17, 27, 43, 66	1 (0%)
1	C	254/288 (88%)	-0.03	13 (5%)	28 22	17, 27, 42, 71	3 (1%)
1	D	256/288 (88%)	0.11	17 (6%)	18 13	17, 28, 45, 71	3 (1%)
All	All	1042/1152 (90%)	0.08	54 (5%)	27 21	17, 27, 43, 74	8 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	81	ALA	5.8
1	C	134	GLU	5.7
1	C	133	ARG	4.5
1	A	132	ASP	4.5
1	D	72	ALA	4.3
1	B	74	VAL	4.0
1	B	164	GLY	4.0
1	B	73	PRO	3.9
1	A	133	ARG	3.9
1	C	120	ARG	3.8
1	C	81	ALA	3.7
1	A	137	GLU	3.7
1	C	136	MET	3.7
1	D	133	ARG	3.6
1	C	230	VAL	3.5
1	B	121	ASN	3.5
1	A	184	THR	3.3
1	A	134	GLU	3.3
1	C	135	ALA	3.3
1	D	5	THR	3.3
1	D	120	ARG	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	136	MET	3.2
1	B	120	ARG	3.2
1	A	73	PRO	3.1
1	B	165	THR	3.0
1	C	72	ALA	2.9
1	D	73	PRO	2.9
1	D	169	HIS	2.8
1	C	241	HIS	2.8
1	A	72	ALA	2.8
1	D	230	VAL	2.7
1	D	167	ALA	2.7
1	C	160	HIS	2.6
1	A	136	MET	2.6
1	B	184	THR	2.5
1	A	138	THR	2.5
1	D	240	GLY	2.4
1	B	136	MET	2.4
1	B	72	ALA	2.4
1	B	53	ALA	2.3
1	D	134	GLU	2.3
1	D	241	HIS	2.3
1	B	137	GLU	2.2
1	A	120	ARG	2.2
1	B	169	HIS	2.2
1	C	169	HIS	2.2
1	D	168	LYS	2.2
1	C	166	PRO	2.1
1	D	82	PRO	2.1
1	A	163	ALA	2.1
1	D	135	ALA	2.1
1	C	167	ALA	2.0
1	B	86	PHE	2.0
1	D	166	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TOP	B	1290	21/21	0.83	0.23	48,54,58,59	0
3	TOP	A	1290	21/21	0.84	0.22	42,50,54,57	0
2	NDP	C	1289	48/48	0.94	0.16	37,40,44,44	0
2	NDP	B	1289	48/48	0.95	0.18	37,40,44,44	0
2	NDP	D	1289	48/48	0.95	0.15	37,40,44,45	0
2	NDP	A	1289	48/48	0.95	0.18	37,40,44,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around TOP A 1290:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NDP C 1289:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NDP B 1289:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NDP D 1289:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around NDP A 1289:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.