



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 02:36 pm BST

PDB ID : 2BG8  
Title : Bacillus cereus metallo-beta-lactamase (BcII) Arg (121) Cys mutant. Solved at pH4.5 using 20 Micromolar ZnSO4 in the buffer. 1mM DTT and 1mM TCEP-HCl were used as reducing agents.  
Authors : Davies, A.M.; Rasia, R.M.; Vila, A.J.; Sutton, B.J.; Fabiane, S.M.  
Deposited on : 2004-12-17  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	
2	B	227	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1293	-	X	-	-
3	GOL	A	1294	-	X	-	-
3	GOL	A	1297	-	X	-	-
3	GOL	A	1303	-	X	-	-
3	GOL	A	1308	-	X	-	-
3	GOL	B	1292	-	X	-	-
3	GOL	B	1294	-	X	-	-
3	GOL	B	1296	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3776 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-LACTAMASE II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1680	1063	289	324	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	CYS	ARG	engineered mutation	UNP P04190

- Molecule 2 is a protein called BETA-LACTAMASE II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	219	Total	C	N	O	S	0	1	0
			1689	1068	290	327	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	121	CYS	ARG	engineered mutation	UNP P04190

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	143	Total 143	O 143	0	0
6	B	145	Total 145	O 145	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BETA-LACTAMASE II



#### • Molecule 2: BETA-LACTAMASE II





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.25Å 67.25Å 177.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.50 58.24 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.6 (6.00-2.50) 97.6 (58.24-2.50)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.66 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.185 , 0.237 0.176 , 0.177	Depositor DCC
$R_{free}$ test set	848 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 63.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3776	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, CSO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/1698	0.64	0/2296
2	B	0.35	0/1715	0.64	0/2321
All	All	0.36	0/3413	0.64	0/4617

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1680	0	1709	48	0
2	B	1689	0	1713	45	0
3	A	42	0	39	5	0
3	B	30	0	26	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	20	0	0	0	0
5	B	25	0	0	1	0
6	A	143	0	0	4	0
6	B	145	0	0	3	0
All	All	3776	0	3487	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:83:VAL:O	2:B:84[B]:ASP:HB3	1.85	0.75
1:A:38:GLU:CD	1:A:38:GLU:H	1.89	0.75
1:A:243:SER:O	1:A:247:GLU:HG2	1.87	0.75
2:B:78:LYS:HD3	2:B:182:PHE:HD1	1.56	0.70
1:A:34:VAL:HG22	1:A:45:SER:HB3	1.73	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/227 (94%)	203 (95%)	10 (5%)	1 (0%)	29	48
2	B	216/227 (95%)	205 (95%)	11 (5%)	0	100	100
All	All	430/454 (95%)	408 (95%)	21 (5%)	1 (0%)	47	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	50	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/195 (95%)	181 (98%)	4 (2%)	52	77
2	B	188/196 (96%)	186 (99%)	2 (1%)	73	89
All	All	373/391 (95%)	367 (98%)	6 (2%)	62	84

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	LYS
2	B	200	ASN
1	A	200	ASN
1	A	45	SER
2	B	147	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	215	ASN
2	B	233	ASN
1	A	255	ASN
1	A	116	HIS
2	B	215	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSO	A	221	1	3,6,7	0.88	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	221	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 2 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	1293	-	5,5,5	3.70	5 (100%)	5,5,5	0.44	0
5	SO4	B	1302	-	4,4,4	0.27	0	6,6,6	0.06	0
3	GOL	A	1297	-	5,5,5	3.87	5 (100%)	5,5,5	0.33	0
3	GOL	A	1293	-	5,5,5	3.75	5 (100%)	5,5,5	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	B	1301	-	4,4,4	0.28	0	6,6,6	0.12	0
5	SO4	B	1299	-	4,4,4	0.28	0	6,6,6	0.08	0
3	GOL	B	1292	-	5,5,5	3.77	5 (100%)	5,5,5	0.42	0
3	GOL	A	1294	-	5,5,5	3.89	5 (100%)	5,5,5	0.42	0
5	SO4	A	1300	-	4,4,4	0.23	0	6,6,6	0.14	0
3	GOL	B	1296	-	5,5,5	3.56	5 (100%)	5,5,5	0.41	0
5	SO4	A	1299	-	4,4,4	0.23	0	6,6,6	0.20	0
3	GOL	B	1295	-	5,5,5	3.75	4 (80%)	5,5,5	0.31	0
3	GOL	A	1308	-	5,5,5	3.71	5 (100%)	5,5,5	0.34	0
5	SO4	B	1298	-	4,4,4	0.23	0	6,6,6	0.11	0
3	GOL	A	1295	-	5,5,5	3.83	4 (80%)	5,5,5	0.29	0
5	SO4	A	1302	-	4,4,4	0.24	0	6,6,6	0.08	0
5	SO4	A	1301	-	4,4,4	0.28	0	6,6,6	0.08	0
3	GOL	A	1296	-	5,5,5	3.83	5 (100%)	5,5,5	0.56	0
5	SO4	B	1300	-	4,4,4	0.26	0	6,6,6	0.07	0
3	GOL	B	1294	-	5,5,5	3.60	5 (100%)	5,5,5	0.50	0
3	GOL	A	1303	-	5,5,5	3.67	5 (100%)	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1293	-	-	0/4/4/4	-
3	GOL	A	1293	-	-	4/4/4/4	-
3	GOL	A	1297	-	-	2/4/4/4	-
3	GOL	A	1308	-	-	1/4/4/4	-
3	GOL	B	1292	-	-	2/4/4/4	-
3	GOL	A	1294	-	-	2/4/4/4	-
3	GOL	B	1296	-	-	2/4/4/4	-
3	GOL	B	1295	-	-	0/4/4/4	-
3	GOL	A	1295	-	-	1/4/4/4	-
3	GOL	A	1296	-	-	0/4/4/4	-
3	GOL	B	1294	-	-	2/4/4/4	-
3	GOL	A	1303	-	-	3/4/4/4	-

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1294	GOL	C3-C2	-6.60	1.24	1.51
3	A	1296	GOL	C3-C2	-6.32	1.25	1.51
3	B	1295	GOL	C3-C2	-6.31	1.25	1.51
3	A	1297	GOL	C3-C2	-6.21	1.26	1.51
3	B	1292	GOL	C3-C2	-6.18	1.26	1.51

There are no bond angle outliers.

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1294	GOL	O1-C1-C2-O2
3	A	1294	GOL	O1-C1-C2-C3
3	B	1296	GOL	C1-C2-C3-O3
3	A	1293	GOL	C1-C2-C3-O3
3	A	1303	GOL	C1-C2-C3-O3

There are no ring outliers.

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1297	GOL	1	0
3	A	1293	GOL	1	0
3	B	1292	GOL	1	0
3	A	1294	GOL	1	0
3	B	1296	GOL	1	0
5	B	1298	SO4	1	0
3	A	1295	GOL	2	0
3	B	1294	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/227 (96%)	-0.60	0 100 100	11, 20, 45, 57	0
2	B	219/227 (96%)	-0.58	0 100 100	12, 21, 38, 52	0
All	All	437/454 (96%)	-0.59	0 100 100	11, 20, 43, 57	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	221	7/8	0.91	0.11	21,22,36,37	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	1294	6/6	0.78	0.24	47,50,52,52	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	1297	6/6	0.81	0.27	55,60,60,60	0
3	GOL	A	1294	6/6	0.82	0.20	49,53,55,56	0
5	SO4	B	1301	5/5	0.83	0.25	99,99,99,99	0
3	GOL	B	1292	6/6	0.87	0.16	40,41,42,42	0
5	SO4	A	1302	5/5	0.88	0.27	81,81,81,83	0
3	GOL	A	1296	6/6	0.88	0.18	61,62,62,62	0
3	GOL	A	1303	6/6	0.88	0.21	42,44,44,45	0
3	GOL	B	1293	6/6	0.89	0.14	49,51,51,51	0
3	GOL	B	1296	6/6	0.90	0.15	58,59,59,60	0
3	GOL	A	1295	6/6	0.90	0.15	51,53,54,54	0
5	SO4	B	1300	5/5	0.91	0.13	83,83,84,85	0
3	GOL	B	1295	6/6	0.91	0.15	57,58,59,60	0
3	GOL	A	1308	6/6	0.92	0.18	50,50,51,53	0
5	SO4	A	1301	5/5	0.93	0.19	84,85,85,85	0
5	SO4	B	1302	5/5	0.93	0.25	99,99,100,100	0
5	SO4	A	1300	5/5	0.94	0.17	73,73,73,74	0
3	GOL	A	1293	6/6	0.94	0.17	38,39,40,41	0
4	ZN	B	1297	1/1	0.96	0.05	37,37,37,37	0
5	SO4	B	1299	5/5	0.97	0.10	59,59,60,60	0
5	SO4	B	1298	5/5	0.97	0.19	47,47,49,49	0
5	SO4	A	1299	5/5	0.99	0.13	23,25,28,28	0
4	ZN	A	1298	1/1	0.99	0.04	28,28,28,28	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.