



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 06:06 PM BST

PDB ID : 4BH6  
Title : Insights into degron recognition by APC coactivators from the structure of an Acm1-Cdh1 complex  
Authors : He, J.; Chao, W.C.H.; Zhang, Z.; Yang, J.; Cronin, N.; Barford, D.  
Deposited on : 2013-03-29  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	<div> <div style="width: 73%;"></div> <div style="width: 23%;"></div> <div style="width: 4%;"></div> </div> <div> <div style="width: 73%;"></div> <div style="width: 23%;"></div> <div style="width: 4%;"></div> </div>
1	B	308	<div> <div style="width: 74%;"></div> <div style="width: 21%;"></div> <div style="width: 5%;"></div> </div> <div> <div style="width: 74%;"></div> <div style="width: 21%;"></div> <div style="width: 5%;"></div> </div>
1	C	308	<div> <div style="width: 74%;"></div> <div style="width: 21%;"></div> <div style="width: 5%;"></div> </div> <div> <div style="width: 74%;"></div> <div style="width: 21%;"></div> <div style="width: 5%;"></div> </div>
1	D	308	<div> <div style="width: 77%;"></div> <div style="width: 20%;"></div> <div style="width: 3%;"></div> </div> <div> <div style="width: 77%;"></div> <div style="width: 20%;"></div> <div style="width: 3%;"></div> </div>
1	E	308	<div> <div style="width: 70%;"></div> <div style="width: 26%;"></div> <div style="width: 4%;"></div> </div> <div> <div style="width: 70%;"></div> <div style="width: 26%;"></div> <div style="width: 4%;"></div> </div>
1	F	308	<div> <div style="width: 71%;"></div> <div style="width: 24%;"></div> <div style="width: 5%;"></div> </div> <div> <div style="width: 71%;"></div> <div style="width: 24%;"></div> <div style="width: 5%;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	308	 3% 73% 22% ...
1	H	308	 12% 70% 24% . .
2	I	70	 % 51% 33% 9% 7%
2	J	70	 % 56% 26% 9% 10%
2	K	70	 57% 24% 11% 7%
2	L	70	 51% 16% 6% . 26%
2	M	70	 7% 20% 6% . 70%
2	N	70	 21% 6% . 71%
2	O	70	 21% 7% 71%
2	P	70	 4% 24% . . 70%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 21491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APC/C ACTIVATOR PROTEIN CDH1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	0	0
			2371	1488	419	452	12			
1	B	304	Total	C	N	O	S	0	0	0
			2358	1480	415	452	11			
1	C	304	Total	C	N	O	S	0	0	0
			2359	1480	416	451	12			
1	D	305	Total	C	N	O	S	0	0	0
			2364	1483	417	452	12			
1	E	303	Total	C	N	O	S	0	0	0
			2344	1472	410	450	12			
1	F	304	Total	C	N	O	S	0	0	0
			2347	1475	411	449	12			
1	G	303	Total	C	N	O	S	0	0	0
			2350	1474	413	451	12			
1	H	299	Total	C	N	O	S	0	0	0
			2321	1457	406	446	12			

- Molecule 2 is a protein called APC/C-CDH1 MODULATOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	65	Total	C	N	O	P	S	0	0
			528	320	98	107	1	2		
2	J	63	Total	C	N	O	P	S	0	0
			508	309	93	103	1	2		
2	K	65	Total	C	N	O	P	S	0	0
			528	320	98	107	1	2		
2	L	52	Total	C	N	O	P	S	0	0
			430	259	78	90	1	2		
2	M	21	Total	C	N	O	S		0	0
			179	110	33	35	1			
2	N	20	Total	C	N	O	S		0	0
			157	97	29	30	1			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	20	Total	C	N	O	S	0	0	0
			170	105	32	32	1			
2	P	21	Total	C	N	O		0	0	0
			149	93	24	32				


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O	0	0
			7	7		
3	B	8	Total	O	0	0
			8	8		
3	C	5	Total	O	0	0
			5	5		
3	D	3	Total	O	0	0
			3	3		
3	E	4	Total	O	0	0
			4	4		
3	F	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: APC/C ACTIVATOR PROTEIN CDH1



#### • Molecule 1: APC/C ACTIVATOR PROTEIN CDH1




#### • Molecule 1: APC/C ACTIVATOR PROTEIN CDH1






• Molecule 1: APC/C ACTIVATOR PROTEIN CDH1

Chain D: 77% 20%




• Molecule 1: APC/C ACTIVATOR PROTEIN CDH1

Chain E: 70% 26%




• Molecule 1: APC/C ACTIVATOR PROTEIN CDH1

Chain F: 71% 24%



• Molecule 1: APC/C ACTIVATOR PROTEIN CDH1

Chain G: 73% 22%



• Molecule 1: APC/C ACTIVATOR PROTEIN CDH1



• Molecule 2: APC/C-CDH1 MODULATOR 1



• Molecule 2: APC/C-CDH1 MODULATOR 1



• Molecule 2: APC/C-CDH1 MODULATOR 1



• Molecule 2: APC/C-CDH1 MODULATOR 1




Chain L:  51% 16% 6% 26%




• Molecule 2: APC/C-CDH1 MODULATOR 1

Chain M:  7% 20% 6% 70%




• Molecule 2: APC/C-CDH1 MODULATOR 1

Chain N:  21% 6% 71%




• Molecule 2: APC/C-CDH1 MODULATOR 1

Chain O:  21% 7% 71%



• Molecule 2: APC/C-CDH1 MODULATOR 1

Chain P:  4% 24% 70%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	198.43Å 188.14Å 93.44Å 90.00° 92.13° 90.00°	Depositor
Resolution (Å)	69.27 – 2.90 69.27 – 2.90	Depositor EDS
% Data completeness (in resolution range)	96.1 (69.27-2.90) 96.1 (69.27-2.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.219 , 0.257 0.218 , 0.256	Depositor DCC
$R_{free}$ test set	3671 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.7	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 62.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	21491	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7776e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.63	0/2425	0.79	2/3297 (0.1%)
1	B	0.71	4/2412 (0.2%)	0.80	6/3283 (0.2%)
1	C	0.62	0/2413	0.80	2/3284 (0.1%)
1	D	0.56	0/2418	0.75	1/3291 (0.0%)
1	E	0.56	0/2397	0.75	1/3262 (0.0%)
1	F	0.51	0/2401	0.72	2/3269 (0.1%)
1	G	0.52	0/2403	0.72	1/3270 (0.0%)
1	H	0.46	0/2374	0.67	1/3230 (0.0%)
2	I	0.79	4/524 (0.8%)	0.87	0/703
2	J	0.58	0/504	0.80	0/677
2	K	0.66	0/524	0.88	1/703 (0.1%)
2	L	0.61	0/427	0.80	0/576
2	M	0.46	0/182	0.72	1/244 (0.4%)
2	N	0.53	0/160	0.76	0/216
2	O	0.44	0/173	0.71	0/232
2	P	0.52	0/151	0.64	0/206
All	All	0.58	8/21888 (0.0%)	0.76	18/29743 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	I	0	1
2	J	0	1
2	K	0	1
2	L	0	2
All	All	0	6

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	545	PHE	CG-CD2	-9.56	1.24	1.38
1	B	545	PHE	CG-CD1	-9.13	1.25	1.38
2	I	92	ASN	CG-ND2	-6.11	1.17	1.32
2	I	100	ASN	CG-ND2	-5.72	1.18	1.32
2	I	100	ASN	CG-OD1	-5.69	1.11	1.24

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	452	ARG	NE-CZ-NH2	-7.36	116.62	120.30
1	A	452	ARG	NE-CZ-NH2	-7.08	116.76	120.30
1	H	497	LEU	CA-CB-CG	7.00	131.40	115.30
1	C	497	LEU	CA-CB-CG	6.60	130.47	115.30
1	F	497	LEU	CA-CB-CG	6.53	130.32	115.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	362	ARG	Peptide
2	I	107	CYS	Peptide
2	J	107	CYS	Peptide
2	K	107	CYS	Peptide
2	L	107	CYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2309	45	0
1	B	2358	0	2280	54	0
1	C	2359	0	2281	42	0
1	D	2364	0	2283	38	0
1	E	2344	0	2262	59	0
1	F	2347	0	2266	55	0
1	G	2350	0	2269	52	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	2321	0	2246	57	0
2	I	528	0	498	24	0
2	J	508	0	476	23	0
2	K	528	0	495	24	0
2	L	430	0	390	24	0
2	M	179	0	158	7	0
2	N	157	0	128	6	0
2	O	170	0	152	2	0
2	P	149	0	107	4	0
3	A	7	0	0	3	0
3	B	8	0	0	2	0
3	C	5	0	0	1	0
3	D	3	0	0	0	0
3	E	4	0	0	1	0
3	F	1	0	0	0	0
All	All	21491	0	20600	433	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 433 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:264:PHE:H	2:L:92:ASN:HD21	1.08	0.99
1:G:264:PHE:H	2:J:92:ASN:HD21	1.14	0.92
1:B:544:LEU:HB3	1:B:545:PHE:CE1	2.04	0.91
1:E:494:LYS:HE3	3:E:2003:HOH:O	1.75	0.86
1:B:544:LEU:C	1:B:545:PHE:HD1	1.77	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/308 (98%)	282 (93%)	16 (5%)	4 (1%)	12	37
1	B	302/308 (98%)	278 (92%)	20 (7%)	4 (1%)	12	37
1	C	302/308 (98%)	279 (92%)	20 (7%)	3 (1%)	15	45
1	D	303/308 (98%)	278 (92%)	22 (7%)	3 (1%)	15	45
1	E	301/308 (98%)	274 (91%)	24 (8%)	3 (1%)	15	45
1	F	302/308 (98%)	277 (92%)	21 (7%)	4 (1%)	12	37
1	G	301/308 (98%)	278 (92%)	19 (6%)	4 (1%)	12	37
1	H	297/308 (96%)	274 (92%)	20 (7%)	3 (1%)	15	45
2	I	60/70 (86%)	51 (85%)	6 (10%)	3 (5%)	2	7
2	J	58/70 (83%)	50 (86%)	6 (10%)	2 (3%)	3	15
2	K	60/70 (86%)	51 (85%)	6 (10%)	3 (5%)	2	7
2	L	49/70 (70%)	41 (84%)	6 (12%)	2 (4%)	3	11
2	M	19/70 (27%)	18 (95%)	1 (5%)	0	100	100
2	N	18/70 (26%)	18 (100%)	0	0	100	100
2	O	18/70 (26%)	18 (100%)	0	0	100	100
2	P	19/70 (27%)	18 (95%)	0	1 (5%)	2	6
All	All	2711/3024 (90%)	2485 (92%)	187 (7%)	39 (1%)	11	36

5 of 39 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	CYS
1	A	460	ASN
1	B	299	CYS
1	C	299	CYS
1	D	299	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/266 (98%)	240 (92%)	22 (8%)	11	31
1	B	259/266 (97%)	239 (92%)	20 (8%)	13	35
1	C	259/266 (97%)	238 (92%)	21 (8%)	11	33
1	D	259/266 (97%)	240 (93%)	19 (7%)	14	38
1	E	257/266 (97%)	238 (93%)	19 (7%)	13	38
1	F	257/266 (97%)	236 (92%)	21 (8%)	11	32
1	G	258/266 (97%)	238 (92%)	20 (8%)	12	34
1	H	256/266 (96%)	237 (93%)	19 (7%)	13	38
2	I	56/61 (92%)	49 (88%)	7 (12%)	4	14
2	J	54/61 (88%)	47 (87%)	7 (13%)	4	12
2	K	56/61 (92%)	49 (88%)	7 (12%)	4	14
2	L	47/61 (77%)	42 (89%)	5 (11%)	6	20
2	M	18/61 (30%)	15 (83%)	3 (17%)	2	6
2	N	14/61 (23%)	12 (86%)	2 (14%)	3	10
2	O	17/61 (28%)	14 (82%)	3 (18%)	2	5
2	P	11/61 (18%)	9 (82%)	2 (18%)	1	5
All	All	2340/2616 (89%)	2143 (92%)	197 (8%)	11	31

5 of 197 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	373	MET
1	F	373	MET
2	L	88	VAL
1	E	386	GLN
1	E	544	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	316	HIS
1	H	514	HIS
2	M	79	ASN
1	G	520	HIS
1	H	520	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SEP	L	102	2	8,9,10	0.78	0	8,12,14	0.97	0
2	SEP	K	102	2	8,9,10	0.60	0	8,12,14	1.49	1 (12%)
2	SEP	J	102	2	8,9,10	0.76	0	8,12,14	1.47	2 (25%)
2	SEP	I	102	2	8,9,10	0.95	0	8,12,14	1.73	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	L	102	2	-	0/5/8/10	-
2	SEP	K	102	2	-	0/5/8/10	-
2	SEP	J	102	2	-	0/5/8/10	-
2	SEP	I	102	2	-	0/5/8/10	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	102	SEP	O3P-P-OG	-3.34	97.84	106.73
2	J	102	SEP	O3P-P-OG	-2.96	98.86	106.73
2	K	102	SEP	O3P-P-OG	-2.56	99.93	106.73
2	J	102	SEP	O3P-P-O2P	2.10	115.66	107.64



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	102	SEP	1	0
2	J	102	SEP	1	0
2	I	102	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	304/308 (98%)	-0.21	1 (0%) 94 94	23, 45, 70, 87	0
1	B	304/308 (98%)	0.00	2 (0%) 87 87	38, 59, 80, 95	0
1	C	304/308 (98%)	-0.29	0 100 100	26, 46, 72, 89	0
1	D	305/308 (99%)	-0.19	0 100 100	31, 55, 87, 101	0
1	E	303/308 (98%)	-0.01	2 (0%) 87 87	27, 59, 99, 121	0
1	F	304/308 (98%)	0.06	6 (1%) 65 63	33, 62, 96, 115	0
1	G	303/308 (98%)	0.02	8 (2%) 56 52	40, 62, 87, 104	0
1	H	299/308 (97%)	0.66	38 (12%) 3 2	36, 98, 183, 227	0
2	I	64/70 (91%)	-0.10	1 (1%) 72 71	29, 62, 95, 102	0
2	J	62/70 (88%)	0.08	1 (1%) 72 71	41, 77, 111, 115	0
2	K	64/70 (91%)	-0.15	0 100 100	29, 62, 96, 109	0
2	L	51/70 (72%)	-0.18	0 100 100	41, 71, 103, 106	0
2	M	21/70 (30%)	0.99	5 (23%) 0 0	127, 146, 159, 166	0
2	N	20/70 (28%)	0.31	0 100 100	98, 113, 129, 145	0
2	O	20/70 (28%)	0.45	0 100 100	100, 112, 142, 166	0
2	P	21/70 (30%)	1.00	3 (14%) 2 2	98, 130, 155, 179	0
All	All	2749/3024 (90%)	0.02	67 (2%) 59 56	23, 59, 121, 227	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	329	TYR	4.8
1	H	357	LEU	4.4
1	H	318	ALA	4.4
1	E	548	PRO	4.3
1	H	546	ASP	4.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SEP	L	102	10/11	0.93	0.17	52,54,59,60	0
2	SEP	K	102	10/11	0.96	0.15	39,42,44,45	0
2	SEP	J	102	10/11	0.96	0.14	47,48,49,51	0
2	SEP	I	102	10/11	0.98	0.15	29,31,32,32	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.