



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:07 am BST

PDB ID : 3BIT
Title : Crystal structure of yeast Spt16 N-terminal Domain
Authors : VanDemark, A.P.; Xin, H.; McCullough, L.; Rawlins, R.; Bentley, S.; Heroux, A.; David, S.J.; Hill, C.P.; Formosa, T.
Deposited on : 2007-11-30
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>..</div> </div> </div>
1	B	453	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>.</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FACT complex subunit SPT16.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	Se	4	17	0
			3738	2391	611	724	6	6			
1	B	438	Total	C	N	O	S	Se	0	14	0
			3651	2342	593	705	6	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP P32558
A	0	HIS	-	EXPRESSION TAG	UNP P32558
B	-1	GLY	-	EXPRESSION TAG	UNP P32558
B	0	HIS	-	EXPRESSION TAG	UNP P32558

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Cl	0	0
			2	2		
2	A	2	Total	Cl	0	0
			2	2		

- Molecule 3 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O		
			6	3	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	517	Total 517	O 517	0	0
5	B	517	Total 517	O 517	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FACT complex subunit SPT16



• Molecule 1: FACT complex subunit SPT16



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.68Å 144.63Å 88.51Å 90.00° 101.44° 90.00°	Depositor
Resolution (Å)	39.88 – 1.90 39.87 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.8 (39.88-1.90) 93.8 (39.87-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.89Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.167 , 0.224 0.178 , 0.225	Depositor DCC
R_{free} test set	3741 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.067 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8445	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	1/3810 (0.0%)	0.69	1/5146 (0.0%)
1	B	0.66	0/3721	0.67	3/5026 (0.1%)
All	All	0.66	1/7531 (0.0%)	0.68	4/10172 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	SER	CB-OG	-5.63	1.34	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	100	LEU	CA-CB-CG	5.71	128.42	115.30
1	B	253	LEU	CA-CB-CG	5.55	128.07	115.30
1	B	36	LEU	CA-CB-CG	5.41	127.73	115.30
1	A	274	ARG	NE-CZ-NH1	5.25	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3738	0	3715	48	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3651	0	3626	42	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	4	0	6	0	0
3	B	8	0	12	1	0
4	B	6	0	8	3	0
5	A	517	0	0	9	0
5	B	517	0	0	6	0
All	All	8445	0	7367	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:LEU:HD12	1:A:37:GLY:N	1.86	0.91
1:A:36:LEU:C	1:A:36:LEU:HD12	1.95	0.88
1:B:52:HIS:HD2	1:B:60:PHE:H	1.21	0.86
1:A:347:GLU:O	1:A:351:LYS:HD2	1.75	0.85
1:A:52:HIS:HD2	1:A:60:PHE:H	1.21	0.84
1:A:106:ASN:H	1:A:113:ASN:HD21	1.26	0.84
1:B:89:ASP:HB2	5:B:838:HOH:O	1.77	0.82
1:B:47[A]:LYS:HE2	1:B:371:GLU:OE1	1.87	0.74
1:B:109:GLU:O	1:B:112:LEU:HB3	1.88	0.73
1:A:402:ASN:HA	1:A:411:ASN:HD21	1.54	0.73
4:B:460:GOL:H32	5:B:1033:HOH:O	1.89	0.71
1:B:402:ASN:HA	1:B:411:ASN:HD21	1.57	0.70
1:B:129:LYS:HG2	4:B:460:GOL:H2	1.73	0.69
1:A:249:ASN:HB3	1:A:252:LEU:HD22	1.76	0.68
1:A:106:ASN:HB3	1:A:109:GLU:HG2	1.76	0.67
1:A:234:LEU:HD13	1:A:238:LEU:HD22	1.77	0.67
1:B:57:SER:O	1:B:58:TYR:HB3	1.95	0.67
1:A:39:SER:HB3	1:A:59:GLU:OE2	1.95	0.66
1:A:36:LEU:HD13	1:A:139:TYR:CE2	2.33	0.64
1:A:423:PRO:HG2	1:A:428:GLU:HB3	1.80	0.62
1:B:112:LEU:HB2	5:B:930:HOH:O	1.98	0.61
1:A:35:VAL:HG13	1:A:143:PHE:HE2	1.66	0.60
1:A:197:LEU:HD21	1:A:224:LYS:HG3	1.82	0.59
1:B:271:VAL:HG12	3:B:459:BME:H12	1.85	0.59
1:B:110:PRO:HA	1:B:113:ASN:HB2	1.84	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:249:ASN:HB3	1:B:252:LEU:HD22	1.85	0.58
1:A:36:LEU:CD1	1:A:36:LEU:C	2.68	0.57
1:B:88:ILE:HD11	1:B:102:LEU:HD22	1.88	0.56
1:B:47[B]:LYS:HE2	1:B:166:SER:HB3	1.86	0.56
1:A:43:ASN:ND2	5:A:952:HOH:O	2.38	0.56
1:B:12:LYS:HD2	1:B:90:LEU:HB3	1.87	0.55
1:A:35:VAL:HG12	1:A:144:MSE:SE	2.56	0.55
1:A:293:ARG:HD3	5:A:652:HOH:O	2.07	0.55
1:B:181:GLN:HE22	1:B:424:LEU:HD22	1.72	0.54
1:B:193:LYS:HG3	1:B:234:LEU:HD23	1.90	0.53
1:A:95:GLU:HB2	5:A:726:HOH:O	2.08	0.53
1:A:235:SER:OG	1:A:250:PHE:CZ	2.62	0.52
1:B:347:GLU:HG2	1:B:351:LYS:HE2	1.92	0.52
1:B:71:LYS:HE2	1:B:101:GLU:OE2	2.10	0.51
1:A:88:ILE:HD11	1:A:102:LEU:HD22	1.92	0.51
1:A:244:PRO:O	1:A:245:ASN:HB2	2.10	0.51
1:A:36:LEU:HD13	1:A:139:TYR:CD2	2.47	0.50
1:B:141:GLY:O	1:B:145:THR:HG23	2.12	0.50
1:B:106:ASN:H	1:B:113:ASN:HD21	1.57	0.50
1:B:12:LYS:HE3	1:B:90:LEU:O	2.12	0.50
1:B:426:GLU:HG2	5:B:929:HOH:O	2.11	0.50
1:A:205:ALA:HA	1:A:210:LEU:HD13	1.95	0.49
1:A:204:ARG:NH1	5:A:763:HOH:O	2.46	0.48
1:B:235:SER:HG	1:B:248:PHE:HE1	1.62	0.48
1:B:253:LEU:HD13	1:B:294:TYR:HD1	1.79	0.47
1:A:334:PRO:HG2	1:A:390:ARG:NH1	2.29	0.47
1:A:187[B]:SER:HB3	1:A:299[B]:SER:OG	2.14	0.47
1:A:306:LEU:HB3	1:A:309:PRO:HB3	1.97	0.46
1:A:256:THR:HG21	1:A:372:PHE:CD2	2.51	0.46
1:A:161:ASN:ND2	5:A:453:HOH:O	2.49	0.46
1:B:57:SER:O	1:B:58:TYR:CB	2.63	0.46
1:A:232:LYS:NZ	5:A:620:HOH:O	2.39	0.46
1:A:346:ILE:O	1:A:350[A]:GLU:HG3	2.16	0.45
1:B:71:LYS:HD3	1:B:73[B]:ILE:HD11	1.98	0.45
1:A:61:PRO:HD2	1:A:80:LYS:HE2	1.98	0.45
1:B:306:LEU:HB3	1:B:309:PRO:HB3	1.97	0.45
1:B:8:PHE:HZ	1:B:83:HIS:O	1.99	0.45
1:A:231:LEU:HB3	1:A:250:PHE:CE1	2.52	0.45
1:A:228:VAL:CG1	1:A:232:LYS:HE3	2.46	0.45
1:B:93:ASP:HA	1:B:94:PRO:HD3	1.71	0.45
1:A:106:ASN:HB3	1:A:109:GLU:CG	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:365:GLY:HA3	1:B:395:ASN:O	2.16	0.44
1:B:180:GLU:HG2	1:B:297:TYR:OH	2.18	0.44
1:A:109:GLU:HA	1:A:110:PRO:HD2	1.91	0.44
1:B:105:ARG:HG2	1:B:113:ASN:ND2	2.33	0.44
1:B:61:PRO:O	1:B:80:LYS:NZ	2.49	0.43
1:A:201:GLU:HG2	5:A:816:HOH:O	2.17	0.43
1:A:274:ARG:NH1	5:A:583:HOH:O	2.51	0.43
1:A:408:SER:O	1:A:409:ALA:HB3	2.18	0.43
1:A:263:SER:HB3	1:A:286[B]:ILE:HG12	2.01	0.43
1:A:52:HIS:CD2	1:A:60:PHE:H	2.13	0.43
1:B:351:LYS:HG3	5:B:698:HOH:O	2.18	0.43
1:A:254[B]:ASP:HB2	5:A:679:HOH:O	2.18	0.42
1:A:338:PRO:HB3	1:A:379:LEU:HA	2.00	0.42
1:B:193:LYS:HG3	1:B:234:LEU:CD2	2.50	0.42
1:B:346:ILE:O	1:B:350[A]:GLU:HG3	2.20	0.41
1:B:129:LYS:HE2	4:B:460:GOL:H11	2.02	0.41
1:A:228:VAL:HG13	1:B:223:ASN:HD21	1.84	0.41
1:B:404:LYS:HD2	1:B:409:ALA:O	2.19	0.41
1:A:350[A]:GLU:HG2	1:A:357:VAL:HG21	2.02	0.41
1:A:244:PRO:O	1:A:245:ASN:CB	2.68	0.41
1:B:347:GLU:O	1:B:351:LYS:HG3	2.21	0.40
1:B:75:ILE:HG12	1:B:103:TRP:HB2	2.03	0.40
1:B:226[B]:ASP:CG	5:B:958:HOH:O	2.60	0.40
1:A:139:TYR:O	1:A:144:MSE:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	462/453 (102%)	453 (98%)	9 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	447/453 (99%)	438 (98%)	7 (2%)	2 (0%)	34 24
All	All	909/906 (100%)	891 (98%)	16 (2%)	2 (0%)	47 38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	TYR
1	B	93	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	426/409 (104%)	409 (96%)	17 (4%)	31 22
1	B	415/409 (102%)	398 (96%)	17 (4%)	30 21
All	All	841/818 (103%)	807 (96%)	34 (4%)	30 22

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	18	LEU
1	A	22	TYR
1	A	36	LEU
1	A	52	HIS
1	A	86	LYS
1	A	97	LYS
1	A	100	LEU
1	A	102	LEU
1	A	108	LYS
1	A	210	LEU
1	A	229	LYS
1	A	252	LEU
1	A	351	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	362	LYS
1	A	377	PHE
1	A	424	LEU
1	B	18	LEU
1	B	24	GLU
1	B	36	LEU
1	B	43	ASN
1	B	71	LYS
1	B	89	ASP
1	B	100	LEU
1	B	102	LEU
1	B	105	ARG
1	B	112	LEU
1	B	115	LYS
1	B	234	LEU
1	B	252	LEU
1	B	253	LEU
1	B	377	PHE
1	B	404	LYS
1	B	424	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	52	HIS
1	A	113	ASN
1	A	148	ASN
1	A	214	ASN
1	A	315	ASN
1	A	330	ASN
1	A	389	GLN
1	A	411	ASN
1	B	43	ASN
1	B	52	HIS
1	B	83	HIS
1	B	113	ASN
1	B	148	ASN
1	B	181	GLN
1	B	223	ASN
1	B	315	ASN
1	B	316	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	330	ASN
1	B	411	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BME	B	459	-	3,3,3	0.38	0	1,2,2	0.99	0
3	BME	B	458	-	3,3,3	0.47	0	1,2,2	0.19	0
3	BME	A	456	-	3,3,3	0.33	0	1,2,2	0.33	0
4	GOL	B	460	-	5,5,5	0.70	0	5,5,5	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BME	B	459	-	-	0/1/1/1	-
3	BME	B	458	-	-	1/1/1/1	-
3	BME	A	456	-	-	1/1/1/1	-
4	GOL	B	460	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	458	BME	O1-C1-C2-S2
3	A	456	BME	O1-C1-C2-S2
4	B	460	GOL	C1-C2-C3-O3
4	B	460	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	459	BME	1	0
4	B	460	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/453 (97%)	-0.25	6 (1%) 75 77	8, 19, 40, 84	2 (0%)
1	B	434/453 (95%)	-0.27	5 (1%) 79 81	8, 18, 50, 77	5 (1%)
All	All	876/906 (96%)	-0.26	11 (1%) 77 79	8, 18, 47, 84	7 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	112	LEU	5.7
1	A	41	ALA	3.3
1	A	447	PHE	3.1
1	B	106	ASN	3.1
1	B	83	HIS	2.5
1	A	107	ASN	2.5
1	A	110	PRO	2.5
1	B	142	LYS	2.3
1	A	96	SER	2.3
1	A	112	LEU	2.2
1	B	6	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	B	460	6/6	0.82	0.27	28,31,31,33	0
3	BME	A	456	4/4	0.85	0.17	52,53,53,55	0
3	BME	B	458	4/4	0.88	0.12	23,25,29,35	0
3	BME	B	459	4/4	0.91	0.10	29,34,34,42	0
2	CL	B	457	1/1	0.99	0.06	22,22,22,22	0
2	CL	A	454	1/1	0.99	0.03	18,18,18,18	0
2	CL	A	455	1/1	1.00	0.03	23,23,23,23	0
2	CL	B	456	1/1	1.00	0.02	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.