



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 12:04 am BST

PDB ID : 6BMU
Title : Non-receptor Protein Tyrosine Phosphatase SHP2 in Complex with Allosteric Inhibitors SHP099 and SHP244
Authors : Stams, T.; Fodor, M.
Deposited on : 2017-11-15
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>9%</div> </div> </div>
1	B	526	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	604	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	480	Total	C	N	O	S	0	1	0
			3889	2456	690	724	19			
1	B	474	Total	C	N	O	S	0	2	0
			3856	2436	687	714	19			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 6-(4-azanyl-4-methyl-piperidin-1-yl)-3-[2,3-bis(chloranyl)phenyl]pyrazin-2-amine (three-letter code: 5OD) (formula: C₁₆H₁₉Cl₂N₅) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	0	0
			23	16	2	5		
2	B	1	Total	C	Cl	N	0	0
			23	16	2	5		

- Molecule 3 is 4-[(2-chlorophenyl)methyl]-1-(2-hydroxy-3-methoxyphenyl)[1,2,4]triazolo[4,3-a]quinazolin-5(4H)-one (three-letter code: DZV) (formula: C₂₃H₁₇ClN₄O₃) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			31	23	1	4	3		
3	B	1	Total	C	Cl	N	O	0	0
			31	23	1	4	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	241	Total	O	0	2
			243	243		
6	B	246	Total	O	0	0
			246	246		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.13 Å 213.45 Å 55.94 Å 90.00° 96.75° 90.00°	Depositor
Resolution (Å)	55.55 – 2.12 55.55 – 2.12	Depositor EDS
% Data completeness (in resolution range)	97.3 (55.55-2.12) 97.3 (55.55-2.12)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.12 Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.181 , 0.225 0.183 , 0.226	Depositor DCC
R_{free} test set	2895 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8370	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, 5OD, DZV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/3971	0.65	0/5351
1	B	0.51	0/3940	0.66	0/5309
All	All	0.51	0/7911	0.65	0/10660

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3889	0	3836	19	0
1	B	3856	0	3815	12	0
2	A	23	0	0	0	0
2	B	23	0	0	0	0
3	A	31	0	0	0	0
3	B	31	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	6	0	8	4	0
5	B	12	0	16	1	0
6	A	243	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	246	0	0	1	0
All	All	8370	0	7675	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:GLN:HE22	5:A:604:GOL:H31	1.43	0.83
1:A:294:ASP:HB2	1:A:343:ARG:HE	1.58	0.69
1:A:272:GLU:CB	1:A:301:VAL:HG11	2.28	0.64
1:A:294:ASP:CB	1:A:343:ARG:HE	2.13	0.61
1:A:133:GLY:HA3	1:A:213:LYS:HB2	1.82	0.60
1:B:407:GLY:HA3	5:B:605:GOL:H2	1.88	0.55
1:B:88:LEU:HG	1:B:96:ILE:HG23	1.89	0.55
1:A:222:ASN:O	1:A:225:GLU:HG2	2.08	0.54
1:B:125:LEU:HB3	1:B:216:LEU:HD21	1.96	0.48
1:B:356:THR:OG1	1:B:459:CYS:HB3	2.13	0.48
5:A:604:GOL:H12	6:A:731:HOH:O	2.14	0.47
1:A:256:GLN:HG3	6:A:810:HOH:O	2.15	0.47
1:A:224:ALA:HB2	1:A:484:VAL:HG13	1.96	0.47
1:A:478:ILE:HG23	1:A:482:LYS:HE3	1.97	0.47
1:A:81:TYR:HE1	1:A:87:GLN:HB3	1.80	0.47
1:B:133:GLY:HA3	1:B:213:LYS:HB2	1.97	0.46
1:A:290:VAL:HG11	1:A:344:MET:HG3	1.97	0.46
1:A:325:LYS:HG3	6:A:725:HOH:O	2.15	0.46
1:A:257:GLN:HE22	5:A:604:GOL:C3	2.21	0.46
1:B:366:LYS:HD2	6:B:770:HOH:O	2.17	0.45
1:A:6:TRP:HB3	1:A:101:PRO:HB3	1.99	0.45
1:B:4:ARG:HD3	1:B:256:GLN:HA	1.99	0.43
1:B:11:ILE:HG13	1:B:15:GLU:HB3	2.01	0.43
1:A:257:GLN:NE2	5:A:604:GOL:H31	2.20	0.43
1:A:125:LEU:HB3	1:A:216:LEU:HD21	1.99	0.43
1:A:41:PHE:HB2	1:A:56:ILE:HB	2.01	0.43
1:B:362:ARG:HD3	1:B:425:ASP:HB3	2.01	0.42
1:B:65:LEU:HD21	1:B:88:LEU:HD13	2.02	0.41
1:A:107:PRO:HG3	1:A:190:LEU:HD12	2.02	0.41
1:A:54:ILE:HD11	1:A:96:ILE:HD13	2.02	0.41
1:B:432:PRO:HG3	1:B:516:MET:HG2	2.03	0.41
1:B:309:ILE:HD13	1:B:328:ILE:HG12	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/526 (88%)	448 (97%)	11 (2%)	4 (1%)	17	12
1	B	460/526 (88%)	449 (98%)	10 (2%)	1 (0%)	47	48
All	All	923/1052 (88%)	897 (97%)	21 (2%)	5 (0%)	29	25

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	295	GLY
1	A	115	GLY
1	B	505	VAL
1	A	301	VAL
1	A	505	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	428/468 (92%)	407 (95%)	21 (5%)	25	22
1	B	424/468 (91%)	409 (96%)	15 (4%)	36	37
All	All	852/936 (91%)	816 (96%)	36 (4%)	30	29

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	62	TYR
1	A	87	GLN
1	A	94	ASP
1	A	102	LEU
1	A	129	LYS
1	A	152	ARG
1	A	205	THR
1	A	211	GLN
1	A	213	LYS
1	A	249	GLU
1	A	294	ASP
1	A	324	LYS
1	A	325	LYS
1	A	335	GLN
1	A	410	ASN
1	A	413	ARG
1	A	439	LEU
1	A	481	GLU
1	A	484	VAL
1	A	508	GLU
1	B	47	ARG
1	B	96	ILE
1	B	111	ARG
1	B	152	ARG
1	B	166	LYS
1	B	171	MET
1	B	173	ARG
1	B	205	THR
1	B	211	GLN
1	B	275	ASN
1	B	280	LYS
1	B	303	ASP
1	B	325	LYS
1	B	413	ARG
1	B	508	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
1	A	114	HIS
1	A	116	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	256	GLN
1	B	114	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DZV	B	602	-	30,35,35	1.90	6 (20%)	36,51,51	1.13	4 (11%)
5	GOL	A	604	-	5,5,5	0.08	0	5,5,5	0.35	0
4	PO4	A	603	-	4,4,4	2.50	1 (25%)	6,6,6	0.57	0
4	PO4	B	603	-	4,4,4	2.45	1 (25%)	6,6,6	0.53	0
2	5OD	A	601	-	23,25,25	1.36	2 (8%)	28,37,37	1.66	8 (28%)
5	GOL	B	605	-	5,5,5	0.07	0	5,5,5	0.18	0
2	5OD	B	601	-	23,25,25	1.19	2 (8%)	28,37,37	2.00	10 (35%)
3	DZV	A	602	-	30,35,35	1.85	5 (16%)	36,51,51	1.50	6 (16%)
5	GOL	B	604	-	5,5,5	0.10	0	5,5,5	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DZV	B	602	-	-	3/10/10/10	0/5/5/5
5	GOL	A	604	-	-	2/4/4/4	-
2	5OD	A	601	-	-	0/8/20/20	0/3/3/3
5	GOL	B	605	-	-	0/4/4/4	-
2	5OD	B	601	-	-	0/8/20/20	0/3/3/3
3	DZV	A	602	-	-	3/10/10/10	0/5/5/5
5	GOL	B	604	-	-	2/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	DZV	C9-C10	6.47	1.53	1.41
3	A	602	DZV	C9-C10	6.12	1.52	1.41
4	A	603	PO4	P-O1	4.14	1.60	1.50
4	B	603	PO4	P-O1	4.05	1.60	1.50
2	A	601	5OD	C1-C6	-3.51	1.41	1.45
3	A	602	DZV	C10-C11	3.41	1.44	1.41
3	B	602	DZV	C10-C11	3.30	1.44	1.41
2	A	601	5OD	C23-C12	3.09	1.56	1.53
2	B	601	5OD	C23-C12	3.07	1.56	1.53
2	B	601	5OD	C1-C6	-2.65	1.42	1.45
3	B	602	DZV	C19-C18	2.46	1.43	1.40
3	A	602	DZV	C14-C15	2.32	1.42	1.36
3	A	602	DZV	C13-C12	2.31	1.42	1.36
3	B	602	DZV	C13-C12	2.22	1.41	1.36
3	A	602	DZV	C21-C19	2.21	1.44	1.39
3	B	602	DZV	C14-C15	2.17	1.41	1.36
3	B	602	DZV	C4-C3	2.12	1.42	1.39

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	5OD	C1-C6-N7	5.39	125.60	121.11
3	A	602	DZV	C18-C17-C16	-4.05	119.10	121.94
2	A	601	5OD	C15-C16-CL2	-3.74	116.88	120.52
3	A	602	DZV	C15-C10-C11	3.49	119.54	117.96
2	A	601	5OD	N5-C4-N9	3.35	120.31	116.55
2	B	601	5OD	N5-C4-N9	3.09	120.02	116.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	5OD	C3-C4-N5	-3.08	117.14	121.32
2	B	601	5OD	C15-C16-CL2	-3.08	117.52	120.52
3	B	602	DZV	C3-C2-N2	2.89	118.04	113.26
3	A	602	DZV	C3-C2-N2	2.85	117.96	113.26
2	A	601	5OD	C3-C4-N5	-2.85	117.47	121.32
2	B	601	5OD	C8-C1-C6	2.83	125.84	122.34
3	A	602	DZV	C23-C17-C18	2.65	120.59	117.96
3	B	602	DZV	C20-O2-C19	2.57	121.40	117.53
3	A	602	DZV	C17-C16-N4	-2.50	120.15	123.58
2	A	601	5OD	C8-C15-CL1	2.46	121.80	119.66
2	A	601	5OD	C1-C6-N5	-2.41	118.53	120.09
2	B	601	5OD	C11-C10-N9	-2.39	106.12	110.34
2	B	601	5OD	C16-C15-CL1	-2.32	117.03	120.02
2	A	601	5OD	C10-N9-C14	2.29	116.57	111.52
2	B	601	5OD	C8-C15-CL1	2.24	121.61	119.66
3	B	602	DZV	C17-C16-N4	-2.20	120.56	123.58
2	A	601	5OD	C10-C11-C12	2.20	117.06	113.22
2	B	601	5OD	C1-C6-N5	-2.19	118.67	120.09
2	A	601	5OD	C3-N2-C1	2.16	121.87	117.95
2	B	601	5OD	C10-N9-C14	2.12	116.19	111.52
3	A	602	DZV	C8-C3-C4	2.09	120.35	116.91
3	B	602	DZV	C15-C10-C11	2.07	118.90	117.96

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	604	GOL	O1-C1-C2-C3
5	B	604	GOL	O1-C1-C2-C3
5	A	604	GOL	O1-C1-C2-O2
5	B	604	GOL	O1-C1-C2-O2
3	B	602	DZV	N4-C16-C17-C18
3	B	602	DZV	N4-C16-C17-C23
3	A	602	DZV	N4-C16-C17-C18
3	A	602	DZV	N4-C16-C17-C23
3	B	602	DZV	C3-C2-N2-C1
3	A	602	DZV	C3-C2-N2-C1

There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	604	GOL	4	0
5	B	605	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 5OD A 601



Ligand 5OD B 601





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	480/526 (91%)	0.15	11 (2%) 60 65	21, 40, 67, 82	0
1	B	474/526 (90%)	0.16	8 (1%) 70 74	22, 41, 66, 81	0
All	All	954/1052 (90%)	0.15	19 (1%) 65 69	21, 40, 66, 82	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	LEU	6.1
1	B	426	HIS	4.2
1	A	116	HIS	4.1
1	A	37	ASN	3.4
1	B	205	THR	3.4
1	A	205	THR	3.3
1	A	177	LEU	3.3
1	B	171	MET	2.9
1	B	11	ILE	2.8
1	A	95	VAL	2.5
1	A	182	GLY	2.4
1	A	525	LEU	2.3
1	B	179	TYR	2.3
1	A	469	PHE	2.2
1	A	127	THR	2.2
1	B	116	HIS	2.2
1	B	96	ILE	2.1
1	A	47	ARG	2.1
1	A	85	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	604	6/6	0.69	0.19	70,70,71,71	0
5	GOL	B	604	6/6	0.77	0.18	58,63,64,65	0
5	GOL	B	605	6/6	0.84	0.16	52,61,63,64	0
4	PO4	A	603	5/5	0.85	0.14	79,79,80,81	0
3	DZV	B	602	31/31	0.94	0.15	36,40,46,49	0
2	5OD	B	601	23/23	0.95	0.14	31,38,43,45	0
3	DZV	A	602	31/31	0.95	0.14	38,41,45,46	0
2	5OD	A	601	23/23	0.95	0.11	30,36,42,43	0
4	PO4	B	603	5/5	0.96	0.14	50,52,53,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DZV B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 5OD B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DZV A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 5OD A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.