



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 05:05 AM BST

PDB ID : 5BO8  
Title : Structure of human sialyltransferase ST8SiaIII  
Authors : Volkers, G.; Worrall, L.; Strynadka, N.C.J.  
Deposited on : 2015-05-27  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>8%</div> <div>19%</div> </div> </div>
1	B	343	<div> <div>%</div> <div> <div></div> <div>72%</div> <div>8%</div> <div>20%</div> </div> </div>
2	C	2	<div> <div></div> <div>100%</div> </div>
2	D	2	<div> <div></div> <div>100%</div> </div>
2	E	2	<div> <div></div> <div>100%</div> </div>
2	H	2	<div> <div></div> <div>100%</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	I	2	 100%
3	F	4	 25%50%25%
4	G	4	 100%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sia-alpha-2,3-Gal-beta-1,4-GlcNAc-R:alpha 2,8-sialyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	0	0
			2270	1468	392	400	10			
1	B	276	Total	C	N	O	S	0	0	0
			2278	1474	395	399	10			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	ALA	-	expression tag	UNP O43173
A	39	PRO	-	expression tag	UNP O43173
A	40	GLU	-	expression tag	UNP O43173
A	41	HIS	-	expression tag	UNP O43173
A	42	HIS	-	expression tag	UNP O43173
A	43	HIS	-	expression tag	UNP O43173
A	44	HIS	-	expression tag	UNP O43173
A	45	HIS	-	expression tag	UNP O43173
A	46	HIS	-	expression tag	UNP O43173
A	47	ASP	-	expression tag	UNP O43173
A	48	TYR	-	expression tag	UNP O43173
A	49	ASP	-	expression tag	UNP O43173
A	50	ILE	-	expression tag	UNP O43173
A	51	PRO	-	expression tag	UNP O43173
A	52	THR	-	expression tag	UNP O43173
A	53	THR	-	expression tag	UNP O43173
A	54	GLU	-	expression tag	UNP O43173
A	55	ASN	-	expression tag	UNP O43173
A	56	LEU	-	expression tag	UNP O43173
A	57	TYR	-	expression tag	UNP O43173
A	58	PHE	-	expression tag	UNP O43173
A	59	GLN	-	expression tag	UNP O43173
A	60	GLY	-	expression tag	UNP O43173
B	38	ALA	-	expression tag	UNP O43173

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	39	PRO	-	expression tag	UNP O43173
B	40	GLU	-	expression tag	UNP O43173
B	41	HIS	-	expression tag	UNP O43173
B	42	HIS	-	expression tag	UNP O43173
B	43	HIS	-	expression tag	UNP O43173
B	44	HIS	-	expression tag	UNP O43173
B	45	HIS	-	expression tag	UNP O43173
B	46	HIS	-	expression tag	UNP O43173
B	47	ASP	-	expression tag	UNP O43173
B	48	TYR	-	expression tag	UNP O43173
B	49	ASP	-	expression tag	UNP O43173
B	50	ILE	-	expression tag	UNP O43173
B	51	PRO	-	expression tag	UNP O43173
B	52	THR	-	expression tag	UNP O43173
B	53	THR	-	expression tag	UNP O43173
B	54	GLU	-	expression tag	UNP O43173
B	55	ASN	-	expression tag	UNP O43173
B	56	LEU	-	expression tag	UNP O43173
B	57	TYR	-	expression tag	UNP O43173
B	58	PHE	-	expression tag	UNP O43173
B	59	GLN	-	expression tag	UNP O43173
B	60	GLY	-	expression tag	UNP O43173

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula:  $C_6H_8O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			13	6	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	35	Total	O	0	0
			35	35		
7	B	25	Total	O	0	0
			25	25		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sia-alpha-2,3-Gal-beta-1,4-GlcNAc-R:alpha 2,8-sialyltransferase



- Molecule 1: Sia-alpha-2,3-Gal-beta-1,4-GlcNAc-R:alpha 2,8-sialyltransferase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain D:  100%


NAG1  
NAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%


NAG1  
NAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

NAG1  
NAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

NAG1  
NAG2

- Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  25% 50% 25%

NAG1  
NAG2  
BMA3  
MAN4

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

NAG1  
NAG2  
BMA3  
FUC4

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.17Å 94.22Å 126.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.43 – 2.70 51.31 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.5 (52.43-2.70) 96.5 (51.31-2.70)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.42 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.193 , 0.246 0.201 , 0.251	Depositor DCC
$R_{free}$ test set	1450 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.7	Xtriage
Anisotropy	0.507	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 36.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4874	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, CIT, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	0/2336	0.86	5/3168 (0.2%)
1	B	0.68	0/2344	0.82	3/3176 (0.1%)
All	All	0.69	0/4680	0.84	8/6344 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	188	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	B	188	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	A	122	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	A	199	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	A	366	MET	CG-SD-CE	-5.26	91.78	100.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	295	SER	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	B	295	SER	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2270	0	2193	7	0
1	B	2278	0	2222	8	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
2	E	28	0	25	0	0
2	H	28	0	25	0	0
2	I	28	0	25	0	0
3	F	50	0	43	2	0
4	G	49	0	43	0	0
5	B	14	0	13	0	0
6	B	13	0	5	0	0
7	A	35	0	0	0	0
7	B	25	0	0	0	0
All	All	4874	0	4644	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:212:PRO:O	1:B:256:VAL:HG11	1.89	0.73
1:B:188:ARG:NH1	1:B:204:LYS:O	2.36	0.59
1:A:188:ARG:NH1	1:A:204:LYS:O	2.37	0.58
3:F:4:MAN:O3	3:F:4:MAN:H62	2.04	0.57
1:A:188:ARG:NH2	1:A:201:VAL:O	2.40	0.54

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/343 (80%)	264 (97%)	9 (3%)	0	100	100
1	B	272/343 (79%)	261 (96%)	10 (4%)	1 (0%)	34	60
All	All	545/686 (79%)	525 (96%)	19 (4%)	1 (0%)	47	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	296	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/313 (79%)	235 (95%)	12 (5%)	25	52
1	B	250/313 (80%)	239 (96%)	11 (4%)	28	56
All	All	497/626 (79%)	474 (95%)	23 (5%)	27	54

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	361	GLN
1	B	103	GLU
1	B	295	SER
1	A	372	THR
1	B	113	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	367	HIS
1	B	367	HIS
1	B	291	ASN
1	A	156	ASN
1	B	100	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.71	0	17,19,21	1.82	7 (41%)
2	NAG	C	2	2	14,14,15	0.56	0	17,19,21	1.07	2 (11%)
2	NAG	D	1	1,2	14,14,15	0.64	0	17,19,21	2.00	6 (35%)
2	NAG	D	2	2	14,14,15	0.73	0	17,19,21	1.45	3 (17%)
2	NAG	E	1	1,2	14,14,15	0.81	0	17,19,21	1.21	1 (5%)
2	NAG	E	2	2	14,14,15	0.45	0	17,19,21	1.07	1 (5%)
3	NAG	F	1	1,3	14,14,15	0.69	0	17,19,21	1.45	3 (17%)
3	NAG	F	2	3	14,14,15	0.40	0	17,19,21	0.86	0
3	BMA	F	3	3	11,11,12	0.62	0	15,15,17	1.28	1 (6%)
3	MAN	F	4	3	11,11,12	0.24	0	15,15,17	0.67	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	1	1,4	14,14,15	0.45	0	17,19,21	1.50	3 (17%)
4	NAG	G	2	4	14,14,15	0.52	0	17,19,21	1.31	2 (11%)
4	BMA	G	3	4	11,11,12	0.59	0	15,15,17	1.03	1 (6%)
4	FUC	G	4	4	10,10,11	0.47	0	14,14,16	1.17	1 (7%)
2	NAG	H	1	1,2	14,14,15	0.74	0	17,19,21	1.47	3 (17%)
2	NAG	H	2	2	14,14,15	0.38	0	17,19,21	1.31	2 (11%)
2	NAG	I	1	1,2	14,14,15	0.69	0	17,19,21	2.05	3 (17%)
2	NAG	I	2	2	14,14,15	0.62	0	17,19,21	1.63	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	2/2/19/22	0/1/1/1
3	MAN	F	4	3	-	2/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	2/2/19/22	0/1/1/1
4	FUC	G	4	4	-	-	0/1/1/1
2	NAG	H	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	1	NAG	C1-O5-C5	-6.22	103.76	112.19
2	H	1	NAG	C1-O5-C5	4.20	117.89	112.19
2	I	2	NAG	C3-C4-C5	-4.20	102.75	110.24
2	D	1	NAG	C8-C7-N2	4.08	123.01	116.10
2	I	1	NAG	O5-C1-C2	-3.75	105.36	111.29

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
3	F	3	BMA	O5-C5-C6-O6
3	F	4	MAN	O5-C5-C6-O6
2	I	2	NAG	C4-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	3	BMA	1	0
3	F	4	MAN	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


















## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	CIT	B	410	-	3,12,12	1.51	0	3,17,17	3.25	1 (33%)
5	NAG	B	405	1	14,14,15	1.27	2 (14%)	17,19,21	2.52	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CIT	B	410	-	-	6/6/16/16	-
5	NAG	B	405	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	405	NAG	C1-C2	3.40	1.57	1.52
5	B	405	NAG	C3-C2	2.37	1.57	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	410	CIT	C3-C2-C1	5.25	123.38	114.98
5	B	405	NAG	C1-C2-N2	3.99	117.30	110.49
5	B	405	NAG	C2-N2-C7	3.92	128.49	122.90
5	B	405	NAG	O5-C1-C2	-3.79	105.31	111.29
5	B	405	NAG	O3-C3-C2	3.63	116.98	109.47

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	410	CIT	C1-C2-C3-O7
6	B	410	CIT	C1-C2-C3-C4
6	B	410	CIT	C2-C3-C4-C5
6	B	410	CIT	O7-C3-C4-C5
6	B	410	CIT	C6-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	277/343 (80%)	0.03	9 (3%) 47 48	30, 52, 96, 138	0
1	B	276/343 (80%)	0.04	4 (1%) 75 77	32, 56, 101, 141	0
All	All	553/686 (80%)	0.03	13 (2%) 59 60	30, 54, 101, 141	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	297	LYS	4.3
1	A	293	HIS	3.6
1	B	293	HIS	3.5
1	B	380	ALA	3.5
1	A	287	ARG	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	D	1	14/15	0.77	0.21	83,103,123,135	0
2	NAG	D	2	14/15	0.78	0.33	92,129,142,146	0
3	MAN	F	4	11/12	0.79	0.22	89,109,116,126	0
4	BMA	G	3	11/12	0.82	0.22	111,120,126,128	0
2	NAG	E	2	14/15	0.84	0.44	106,128,138,140	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	E	1	14/15	0.85	0.32	83,99,114,122	0
3	BMA	F	3	11/12	0.89	0.13	107,118,120,126	0
2	NAG	I	2	14/15	0.89	0.23	34,41,48,52	14
3	NAG	F	2	14/15	0.93	0.17	67,82,99,120	0
2	NAG	C	2	14/15	0.93	0.17	66,79,89,92	0
4	NAG	G	2	14/15	0.94	0.17	72,87,100,116	0
2	NAG	H	2	14/15	0.96	0.17	60,68,74,75	0
2	NAG	I	1	14/15	0.96	0.33	27,32,35,36	14
4	FUC	G	4	10/11	0.96	0.13	63,69,73,74	0
2	NAG	C	1	14/15	0.97	0.14	44,51,57,66	0
2	NAG	H	1	14/15	0.97	0.13	41,45,56,57	0
3	NAG	F	1	14/15	0.97	0.16	38,47,59,71	0
4	NAG	G	1	14/15	0.98	0.11	43,52,65,73	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around Chain I:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain F:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	CIT	B	410	13/13	0.65	0.31	83,104,110,112	0
5	NAG	B	405	14/15	0.83	0.27	79,93,100,111	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.