



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 02:41 am BST

PDB ID : 6BOJ
Title : Crystal Structure of the PDE4D Catalytic Domain and UCR2 Regulatory Helix with BPN5004
Authors : Fox III, D.; Fairman, J.W.; Gurney, M.E.
Deposited on : 2017-11-20
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 5%, green 85%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 85% 5% 9% </div> </div>
1	B	370	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 5%, green 86%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 86% • 9% </div> </div>
1	C	370	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 5%, green 86%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 86% 5% 9% </div> </div>
1	D	370	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 5%, green 86%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 86% • 9% </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12021 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	9	0
			2689	1711	455	507	16			
1	B	335	Total	C	N	O	S	0	7	0
			2681	1701	450	515	15			
1	C	338	Total	C	N	O	S	0	14	0
			2710	1724	448	523	15			
1	D	335	Total	C	N	O	S	0	10	0
			2675	1707	449	503	16			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	MET	-	initiating methionine	UNP Q08499
A	579	ALA	SER	engineered mutation	UNP Q07343
A	581	ALA	SER	engineered mutation	UNP Q07343
A	591	SER	CYS	engineered mutation	UNP Q07343
A	607	GLU	-	expression tag	UNP Q08499
A	608	ASN	-	expression tag	UNP Q08499
A	609	LEU	-	expression tag	UNP Q08499
A	610	TYR	-	expression tag	UNP Q08499
A	611	PHE	-	expression tag	UNP Q08499
A	612	GLN	-	expression tag	UNP Q08499
B	243	MET	-	initiating methionine	UNP Q08499
B	579	ALA	SER	engineered mutation	UNP Q07343
B	581	ALA	SER	engineered mutation	UNP Q07343
B	591	SER	CYS	engineered mutation	UNP Q07343
B	607	GLU	-	expression tag	UNP Q08499
B	608	ASN	-	expression tag	UNP Q08499
B	609	LEU	-	expression tag	UNP Q08499
B	610	TYR	-	expression tag	UNP Q08499
B	611	PHE	-	expression tag	UNP Q08499
B	612	GLN	-	expression tag	UNP Q08499
C	243	MET	-	initiating methionine	UNP Q08499

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	579	ALA	SER	engineered mutation	UNP Q07343
C	581	ALA	SER	engineered mutation	UNP Q07343
C	591	SER	CYS	engineered mutation	UNP Q07343
C	607	GLU	-	expression tag	UNP Q08499
C	608	ASN	-	expression tag	UNP Q08499
C	609	LEU	-	expression tag	UNP Q08499
C	610	TYR	-	expression tag	UNP Q08499
C	611	PHE	-	expression tag	UNP Q08499
C	612	GLN	-	expression tag	UNP Q08499
D	243	MET	-	initiating methionine	UNP Q08499
D	579	ALA	SER	engineered mutation	UNP Q07343
D	581	ALA	SER	engineered mutation	UNP Q07343
D	591	SER	CYS	engineered mutation	UNP Q07343
D	607	GLU	-	expression tag	UNP Q08499
D	608	ASN	-	expression tag	UNP Q08499
D	609	LEU	-	expression tag	UNP Q08499
D	610	TYR	-	expression tag	UNP Q08499
D	611	PHE	-	expression tag	UNP Q08499
D	612	GLN	-	expression tag	UNP Q08499

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	2	Total Zn 2 2	0	0
2	D	2	Total Zn 2 2	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 2-(4-{[2-(3-chlorophenyl)-6-ethylpyrimidin-4-yl]methyl}phenyl)acetamide (three-letter code: E31) (formula: C₂₁H₂₀ClN₃O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			26	21	1	3	1		
4	B	1	Total	C	Cl	N	O	0	0
			26	21	1	3	1		
4	C	1	Total	C	Cl	N	O	0	0
			26	21	1	3	1		
4	D	1	Total	C	Cl	N	O	0	0
			26	21	1	3	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		
5	D	1	Total	Cl	0	0
			1	1		
5	C	1	Total	Cl	0	0
			1	1		


- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			8	6	2		
6	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	301	Total	O	0	0
			301	301		
7	B	266	Total	O	0	0
			266	266		
7	C	317	Total	O	0	0
			317	317		
7	D	248	Total	O	0	0
			248	248		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.78Å 81.89Å 116.79Å 90.00° 110.41° 90.00°	Depositor
Resolution (Å)	45.50 – 1.70 45.50 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (45.50-1.70) 97.4 (45.50-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.161 , 0.186 0.172 , 0.198	Depositor DCC
R_{free} test set	7761 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12021	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2329e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MPD, MG, E31, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/2767	0.76	6/3768 (0.2%)
1	B	0.66	0/2753	0.75	6/3752 (0.2%)
1	C	0.71	0/2807	0.74	2/3828 (0.1%)
1	D	0.64	0/2759	0.73	4/3761 (0.1%)
All	All	0.68	0/11086	0.74	18/15109 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	423[A]	ARG	NE-CZ-NH2	7.13	123.87	120.30
1	B	423[B]	ARG	NE-CZ-NH2	7.13	123.87	120.30
1	A	484	ASP	CB-CG-OD2	-6.88	112.11	118.30
1	D	423	ARG	NE-CZ-NH1	6.79	123.69	120.30
1	A	423	ARG	NE-CZ-NH1	6.50	123.55	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	593	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2689	0	2608	12	0
1	B	2681	0	2573	8	0
1	C	2710	0	2614	9	0
1	D	2675	0	2594	15	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	26	0	0	0	0
4	B	26	0	0	0	0
4	C	26	0	0	0	0
4	D	26	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	B	8	0	14	1	0
6	C	8	0	14	1	0
7	A	301	0	0	5	0
7	B	266	0	0	1	0
7	C	317	0	0	3	0
7	D	248	0	0	0	0
All	All	12021	0	10417	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:472[B]:ARG:CG	1:D:472[B]:ARG:HH11	1.80	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:472[B]:ARG:CB	1:D:472[B]:ARG:HH11	1.81	0.92
1:D:344[B]:THR:HG22	1:D:557:ASP:OD2	1.69	0.90
1:D:472[B]:ARG:HB2	1:D:472[B]:ARG:HH11	1.32	0.89
1:D:472[B]:ARG:HG3	1:D:472[B]:ARG:HH11	1.46	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/370 (91%)	330 (98%)	8 (2%)	0	100	100
1	B	336/370 (91%)	331 (98%)	5 (2%)	0	100	100
1	C	348/370 (94%)	343 (99%)	5 (1%)	0	100	100
1	D	339/370 (92%)	333 (98%)	6 (2%)	0	100	100
All	All	1361/1480 (92%)	1337 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/338 (88%)	297 (100%)	1 (0%)	92	89
1	B	297/338 (88%)	296 (100%)	1 (0%)	92	89

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	303/338 (90%)	302 (100%)	1 (0%)	92	89
1	D	295/338 (87%)	295 (100%)	0	100	100
All	All	1193/1352 (88%)	1190 (100%)	3 (0%)	92	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	502	ILE
1	B	502	ILE
1	C	502	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 14 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	E31	D	703	-	28,28,28	0.77	1 (3%)	35,38,38	1.33	5 (14%)
6	MPD	C	705	-	7,7,7	0.33	0	9,10,10	1.23	2 (22%)
4	E31	B	703	-	28,28,28	0.76	0	35,38,38	1.34	3 (8%)
4	E31	C	703	-	28,28,28	0.75	0	35,38,38	1.20	1 (2%)
4	E31	A	703	-	28,28,28	0.61	0	35,38,38	1.27	3 (8%)
6	MPD	B	705	-	7,7,7	0.66	0	9,10,10	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	E31	D	703	-	-	4/14/14/14	0/3/3/3
6	MPD	C	705	-	-	3/5/5/5	-
4	E31	B	703	-	-	4/14/14/14	0/3/3/3
4	E31	C	703	-	-	4/14/14/14	0/3/3/3
4	E31	A	703	-	-	4/14/14/14	0/3/3/3
6	MPD	B	705	-	-	3/5/5/5	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	703	E31	C20-C2	2.04	1.42	1.38

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	703	E31	C20-C2-N	-5.28	117.30	122.90
4	C	703	E31	C20-C2-N	-4.36	118.27	122.90
4	A	703	E31	C20-C2-N	-3.83	118.84	122.90
4	B	703	E31	C3-N-C2	3.25	121.15	116.58
4	B	703	E31	C14-C13-C18	2.84	122.18	118.16

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	705	MPD	O2-C2-C3-C4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	C	705	MPD	CM-C2-C3-C4
6	B	705	MPD	C1-C2-C3-C4
6	B	705	MPD	O2-C2-C3-C4
6	B	705	MPD	C2-C3-C4-O4


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	705	MPD	1	0
6	B	705	MPD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand E31 D 703




Ligand E31 B 703



Ligand E31 C 703





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/370 (90%)	-0.35	4 (1%) 79 82	8, 18, 36, 52	0
1	B	335/370 (90%)	-0.29	6 (1%) 68 72	10, 21, 41, 57	0
1	C	338/370 (91%)	-0.33	3 (0%) 84 87	9, 18, 33, 56	0
1	D	335/370 (90%)	-0.27	5 (1%) 73 77	10, 21, 43, 63	0
All	All	1343/1480 (90%)	-0.31	18 (1%) 77 81	8, 19, 38, 63	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	593	GLY	7.6
1	C	593	GLY	4.5
1	A	593	GLY	4.2
1	B	593	GLY	4.0
1	B	577	PRO	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MPD	B	705	8/8	0.80	0.15	27,34,36,38	0
6	MPD	C	705	8/8	0.81	0.17	22,26,27,28	0
5	CL	C	704	1/1	0.90	0.11	29,29,29,29	0
4	E31	B	703	26/26	0.96	0.07	15,17,20,22	0
4	E31	D	703	26/26	0.96	0.07	17,19,21,23	0
4	E31	C	703	26/26	0.97	0.07	13,15,18,18	0
4	E31	A	703	26/26	0.97	0.06	14,15,18,18	0
5	CL	D	705	1/1	0.97	0.04	31,31,31,31	0
5	CL	A	705	1/1	0.98	0.08	25,25,25,25	0
5	CL	B	704	1/1	0.99	0.05	30,30,30,30	0
2	ZN	B	701	1/1	1.00	0.05	13,13,13,13	0
2	ZN	D	704	1/1	1.00	0.03	16,16,16,16	0
2	ZN	D	701	1/1	1.00	0.05	14,14,14,14	0
2	ZN	A	701	1/1	1.00	0.07	11,11,11,11	0
3	MG	C	702	1/1	1.00	0.10	10,10,10,10	0
3	MG	A	702	1/1	1.00	0.08	10,10,10,10	0
3	MG	D	702	1/1	1.00	0.06	11,11,11,11	0
3	MG	B	702	1/1	1.00	0.05	10,10,10,10	0
2	ZN	C	701	1/1	1.00	0.05	11,11,11,11	0
2	ZN	A	704	1/1	1.00	0.04	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around E31 B 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E31 D 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E31 C 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around E31 A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.