



Full wwPDB X-ray Structure Validation Report ⓘ

May 19, 2020 – 01:46 am BST

PDB ID : 6BRU
Title : Crystal Structure of the Human vaccinia-related kinase bound to a (S)-2-phe
nylaminopteridinone inhibitor
Authors : Counago, R.M.; dos Reis, C.V.; de Souza, G.P.; Aquino, B.; Azevedo, A.;
Guimaraes, C.; Mascarello, A.; Gama, F.; Ferreira, M.; Massirer, K.B.; Ar-
ruda, P.; Edwards, A.M.; Elkins, J.M.; Structural Genomics Consortium
(SGC)
Deposited on : 2017-12-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>12%</div> </div> </div>
1	B	364	<div> <div>7%</div> <div> <div></div> <div>80%</div> <div>17%</div> </div> </div>
1	C	364	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>13%</div> </div> </div>
1	D	364	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>15%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase VRK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	1	0
			2546	1626	434	473	13			
1	B	301	Total	C	N	O	S	0	1	0
			2398	1542	409	436	11			
1	C	315	Total	C	N	O	S	0	0	0
			2488	1592	422	461	13			
1	D	311	Total	C	N	O	S	0	1	0
			2442	1569	415	446	12			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q99986
A	2	MET	-	expression tag	UNP Q99986
A	34	ALA	LYS	engineered mutation	UNP Q99986
A	35	ALA	LYS	engineered mutation	UNP Q99986
A	36	ALA	GLU	engineered mutation	UNP Q99986
A	212	ALA	GLU	engineered mutation	UNP Q99986
A	214	ALA	LYS	engineered mutation	UNP Q99986
A	215	ALA	GLU	engineered mutation	UNP Q99986
A	292	ALA	GLU	engineered mutation	UNP Q99986
A	293	ALA	LYS	engineered mutation	UNP Q99986
A	295	ALA	LYS	engineered mutation	UNP Q99986
A	359	ALA	LYS	engineered mutation	UNP Q99986
A	360	ALA	LYS	engineered mutation	UNP Q99986
B	1	SER	-	expression tag	UNP Q99986
B	2	MET	-	expression tag	UNP Q99986
B	34	ALA	LYS	engineered mutation	UNP Q99986
B	35	ALA	LYS	engineered mutation	UNP Q99986
B	36	ALA	GLU	engineered mutation	UNP Q99986
B	212	ALA	GLU	engineered mutation	UNP Q99986
B	214	ALA	LYS	engineered mutation	UNP Q99986
B	215	ALA	GLU	engineered mutation	UNP Q99986

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	292	ALA	GLU	engineered mutation	UNP Q99986
B	293	ALA	LYS	engineered mutation	UNP Q99986
B	295	ALA	LYS	engineered mutation	UNP Q99986
B	359	ALA	LYS	engineered mutation	UNP Q99986
B	360	ALA	LYS	engineered mutation	UNP Q99986
C	1	SER	-	expression tag	UNP Q99986
C	2	MET	-	expression tag	UNP Q99986
C	34	ALA	LYS	engineered mutation	UNP Q99986
C	35	ALA	LYS	engineered mutation	UNP Q99986
C	36	ALA	GLU	engineered mutation	UNP Q99986
C	212	ALA	GLU	engineered mutation	UNP Q99986
C	214	ALA	LYS	engineered mutation	UNP Q99986
C	215	ALA	GLU	engineered mutation	UNP Q99986
C	292	ALA	GLU	engineered mutation	UNP Q99986
C	293	ALA	LYS	engineered mutation	UNP Q99986
C	295	ALA	LYS	engineered mutation	UNP Q99986
C	359	ALA	LYS	engineered mutation	UNP Q99986
C	360	ALA	LYS	engineered mutation	UNP Q99986
D	1	SER	-	expression tag	UNP Q99986
D	2	MET	-	expression tag	UNP Q99986
D	34	ALA	LYS	engineered mutation	UNP Q99986
D	35	ALA	LYS	engineered mutation	UNP Q99986
D	36	ALA	GLU	engineered mutation	UNP Q99986
D	212	ALA	GLU	engineered mutation	UNP Q99986
D	214	ALA	LYS	engineered mutation	UNP Q99986
D	215	ALA	GLU	engineered mutation	UNP Q99986
D	292	ALA	GLU	engineered mutation	UNP Q99986
D	293	ALA	LYS	engineered mutation	UNP Q99986
D	295	ALA	LYS	engineered mutation	UNP Q99986
D	359	ALA	LYS	engineered mutation	UNP Q99986
D	360	ALA	LYS	engineered mutation	UNP Q99986

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is (7S)-2-[(3,5-difluoro-4-hydroxyphenyl)amino]-5,7,8-trimethyl-7,8-dihydropterin-6(5H)-one (three-letter code: E5M) (formula: C₁₅H₁₅F₂N₅O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	F	N	O	0	0
			24	15	2	5	2		
3	D	1	Total	C	F	N	O	0	0
			24	15	2	5	2		


- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	283	Total 283	O 283	0	0
5	B	209	Total 209	O 209	0	0
5	C	273	Total 273	O 273	0	0
5	D	248	Total 248	O 248	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.75Å 97.14Å 193.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.91 – 1.80 19.91 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.91-1.80) 100.0 (19.91-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.195 , 0.214 0.203 , 0.224	Depositor DCC
R_{free} test set	8143 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10984	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: E5M, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2607	0.79	0/3533
1	B	0.59	0/2457	0.76	0/3326
1	C	0.64	0/2547	0.77	0/3454
1	D	0.65	0/2503	0.74	0/3396
All	All	0.63	0/10114	0.76	0/13709

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2546	0	2468	4	0
1	B	2398	0	2331	4	0
1	C	2488	0	2412	7	0
1	D	2442	0	2351	4	0
2	A	10	0	0	0	0
2	B	15	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
3	B	24	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	24	0	0	1	0
4	D	4	0	6	0	0
5	A	283	0	0	1	0
5	B	209	0	0	1	0
5	C	273	0	0	0	0
5	D	248	0	0	0	0
All	All	10984	0	9568	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:LYS:NZ	5:A:501:HOH:O	2.23	0.68
1:B:89:ARG:HG2	5:B:602:HOH:O	2.00	0.59
1:C:294:ASN:O	1:C:296:PRO:HD3	2.08	0.53
1:B:264:ASN:HB3	1:B:270:TYR:CD2	2.44	0.52
1:D:264:ASN:HB3	1:D:270:TYR:CD2	2.44	0.51
1:D:73:GLU:OE2	1:D:77:ASN:CB	2.59	0.50
1:C:264:ASN:HB3	1:C:270:TYR:CD2	2.46	0.50
1:A:267:ASP:OD1	1:A:269:LYS:HG2	2.12	0.50
1:A:204:TYR:CE2	1:A:205:CYS:HB2	2.50	0.47
1:A:200:LEU:HD13	1:A:221:HIS:CG	2.50	0.45
1:C:200:LEU:HD13	1:C:221:HIS:CG	2.51	0.45
1:C:267:ASP:OD1	1:C:269:LYS:HG2	2.17	0.44
1:D:204:TYR:CE2	1:D:205:CYS:HB2	2.53	0.44
1:B:204:TYR:CE2	1:B:205:CYS:HB2	2.54	0.42
1:C:291:PRO:O	1:C:294:ASN:HB2	2.18	0.42
1:C:204:TYR:CE2	1:C:205:CYS:HB2	2.54	0.42
3:D:401:E5M:N3	3:D:401:E5M:C15	2.82	0.42
1:C:87:TYR:HB3	1:C:113:TYR:HB2	2.00	0.42
1:D:183:LEU:C	1:D:184:LEU:HD12	2.41	0.41
1:B:183:LEU:C	1:B:184:LEU:HD12	2.41	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/364 (88%)	316 (98%)	5 (2%)	0	100	100
1	B	292/364 (80%)	287 (98%)	5 (2%)	0	100	100
1	C	311/364 (85%)	307 (99%)	4 (1%)	0	100	100
1	D	306/364 (84%)	302 (99%)	4 (1%)	0	100	100
All	All	1230/1456 (84%)	1212 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/304 (86%)	258 (98%)	4 (2%)	65	56
1	B	244/304 (80%)	239 (98%)	5 (2%)	55	44
1	C	257/304 (84%)	255 (99%)	2 (1%)	81	78
1	D	245/304 (81%)	243 (99%)	2 (1%)	81	78
All	All	1008/1216 (83%)	995 (99%)	13 (1%)	73	62

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ILE
1	A	103[A]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	103[B]	ARG
1	A	133	ARG
1	B	93	PRO
1	B	103[A]	ARG
1	B	103[B]	ARG
1	B	133	ARG
1	B	341	LEU
1	C	103	ARG
1	C	133	ARG
1	D	93	PRO
1	D	133	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	C	401	-	4,4,4	0.17	0	6,6,6	0.23	0
2	SO4	A	402	-	4,4,4	0.40	0	6,6,6	0.18	0
2	SO4	B	402	-	4,4,4	0.38	0	6,6,6	0.25	0
3	E5M	D	401	-	26,26,26	0.81	1 (3%)	34,39,39	1.12	2 (5%)
2	SO4	D	402	-	4,4,4	0.44	0	6,6,6	0.39	0
3	E5M	B	401	-	26,26,26	0.68	1 (3%)	34,39,39	1.18	3 (8%)
2	SO4	D	403	-	4,4,4	0.13	0	6,6,6	0.19	0
2	SO4	B	403	-	4,4,4	0.45	0	6,6,6	0.26	0
4	EDO	D	404	-	3,3,3	0.29	0	2,2,2	0.53	0
2	SO4	C	402	-	4,4,4	0.18	0	6,6,6	0.19	0
2	SO4	B	404	-	4,4,4	0.27	0	6,6,6	0.29	0
2	SO4	A	401	-	4,4,4	0.25	0	6,6,6	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	E5M	D	401	-	-	0/4/24/24	0/3/3/3
3	E5M	B	401	-	-	0/4/24/24	0/3/3/3
4	EDO	D	404	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	401	E5M	C7-N5	3.03	1.49	1.46
3	B	401	E5M	C7-N5	2.31	1.48	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	E5M	C11-C12-C13	-2.48	121.85	123.79
3	D	401	E5M	C15-C14-C13	-2.37	121.93	123.79
3	B	401	E5M	F2-C12-C13	2.29	118.83	117.13
3	D	401	E5M	F2-C12-C13	2.23	118.78	117.13
3	B	401	E5M	C15-C14-C13	-2.12	122.13	123.79

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	404	EDO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	E5M	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/364 (88%)	0.05	10 (3%) 49 43	17, 27, 44, 80	0
1	B	301/364 (82%)	0.27	24 (7%) 12 9	17, 30, 64, 87	0
1	C	315/364 (86%)	0.03	10 (3%) 47 41	16, 28, 48, 61	0
1	D	311/364 (85%)	0.27	22 (7%) 16 12	15, 28, 65, 99	0
All	All	1249/1456 (85%)	0.15	66 (5%) 26 21	15, 28, 57, 99	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	PHE	8.9
1	C	220	CYS	6.8
1	B	126	TYR	5.1
1	A	220	CYS	5.0
1	D	62	VAL	5.0
1	B	293	ALA	5.0
1	D	78	GLY	5.0
1	D	123	GLY	4.9
1	D	77	ASN	4.6
1	B	34	ALA	4.5
1	B	78	GLY	4.5
1	C	217	PRO	4.2
1	B	24	ALA	4.1
1	B	57	ASN	4.0
1	A	217	PRO	4.0
1	D	122	ASN	3.8
1	A	46	GLY	3.7
1	C	123	GLY	3.7
1	D	61	SER	3.4
1	B	120	ASP	3.4
1	B	65	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	50	CYS	3.3
1	D	76	ASP	3.3
1	D	65	ASP	3.2
1	D	35	ALA	3.1
1	B	23	PHE	3.1
1	B	58	SER	3.1
1	D	63	GLY	3.0
1	B	35	ALA	3.0
1	D	293	ALA	3.0
1	B	42	PRO	3.0
1	D	66	ALA	2.9
1	A	20	ALA	2.9
1	B	64	SER	2.8
1	C	59	SER	2.8
1	B	187	TYR	2.8
1	D	126	TYR	2.8
1	B	74	PRO	2.8
1	B	292	ALA	2.8
1	D	125	SER	2.7
1	C	146	ALA	2.7
1	A	123	GLY	2.7
1	A	146	ALA	2.6
1	D	64	SER	2.6
1	A	47	GLY	2.6
1	B	66	ALA	2.6
1	B	56	MET	2.5
1	B	50	CYS	2.5
1	D	23	PHE	2.5
1	D	28	ILE	2.4
1	B	67	PRO	2.4
1	D	57	ASN	2.4
1	A	294	ASN	2.4
1	B	36	ALA	2.4
1	C	187	TYR	2.3
1	B	26	GLY	2.2
1	C	332	GLY	2.2
1	D	36	ALA	2.2
1	B	121	LYS	2.2
1	D	294	ASN	2.2
1	B	43	ILE	2.1
1	D	146	ALA	2.1
1	C	65	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	126	TYR	2.1
1	A	65	ASP	2.0
1	C	64	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	403	5/5	0.73	0.35	73,84,90,90	0
2	SO4	A	401	5/5	0.82	0.28	66,69,72,81	0
2	SO4	A	402	5/5	0.84	0.34	72,89,92,93	0
4	EDO	D	404	4/4	0.89	0.09	43,46,46,48	0
3	E5M	D	401	24/24	0.90	0.10	26,34,39,44	0
3	E5M	B	401	24/24	0.92	0.10	33,37,44,47	0
2	SO4	C	402	5/5	0.93	0.25	61,63,66,71	0
2	SO4	B	404	5/5	0.94	0.27	69,73,75,77	0
2	SO4	D	403	5/5	0.95	0.20	50,51,53,57	0
2	SO4	B	402	5/5	0.96	0.22	50,56,58,59	0
2	SO4	D	402	5/5	0.98	0.20	40,44,47,52	0
2	SO4	C	401	5/5	0.98	0.12	40,44,47,52	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around E5M D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around E5M B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.