



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 10:46 am BST

PDB ID : 6BTW
Title : Crystal Structure of the Human vaccinia-related kinase bound to a phenyl-pteridinone inhibitor
Authors : Counago, R.M.; dos Reis, C.V.; de Souza, G.P.; Azevedo, A.; Guimaraes, C.; Mascarello, A.; Gama, F.; Ferreira, M.; Massirer, K.B.; Arruda, P.; Edwards, A.M.; Elkins, J.M.; Structural Genomics Consortium (SGC)
Deposited on : 2017-12-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>13%</div> </div> </div>
1	B	364	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>17%</div> </div> </div>
1	C	364	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>13%</div> </div> </div>
1	D	364	<div> <div>8%</div> <div> <div></div> <div>82%</div> <div>16%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase VRK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	2	0
			2515	1605	427	469	14			
1	B	301	Total	C	N	O	S	0	1	0
			2391	1537	410	433	11			
1	C	315	Total	C	N	O	S	0	2	0
			2504	1602	426	462	14			
1	D	307	Total	C	N	O	S	0	1	0
			2425	1560	410	442	13			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q99986
A	2	MET	-	expression tag	UNP Q99986
A	34	ALA	LYS	engineered mutation	UNP Q99986
A	35	ALA	LYS	engineered mutation	UNP Q99986
A	36	ALA	GLU	engineered mutation	UNP Q99986
A	212	ALA	GLU	engineered mutation	UNP Q99986
A	214	ALA	LYS	engineered mutation	UNP Q99986
A	215	ALA	GLU	engineered mutation	UNP Q99986
A	292	ALA	GLU	engineered mutation	UNP Q99986
A	293	ALA	LYS	engineered mutation	UNP Q99986
A	295	ALA	LYS	engineered mutation	UNP Q99986
A	359	ALA	LYS	engineered mutation	UNP Q99986
A	360	ALA	LYS	engineered mutation	UNP Q99986
B	1	SER	-	expression tag	UNP Q99986
B	2	MET	-	expression tag	UNP Q99986
B	34	ALA	LYS	engineered mutation	UNP Q99986
B	35	ALA	LYS	engineered mutation	UNP Q99986
B	36	ALA	GLU	engineered mutation	UNP Q99986
B	212	ALA	GLU	engineered mutation	UNP Q99986
B	214	ALA	LYS	engineered mutation	UNP Q99986
B	215	ALA	GLU	engineered mutation	UNP Q99986

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	292	ALA	GLU	engineered mutation	UNP Q99986
B	293	ALA	LYS	engineered mutation	UNP Q99986
B	295	ALA	LYS	engineered mutation	UNP Q99986
B	359	ALA	LYS	engineered mutation	UNP Q99986
B	360	ALA	LYS	engineered mutation	UNP Q99986
C	1	SER	-	expression tag	UNP Q99986
C	2	MET	-	expression tag	UNP Q99986
C	34	ALA	LYS	engineered mutation	UNP Q99986
C	35	ALA	LYS	engineered mutation	UNP Q99986
C	36	ALA	GLU	engineered mutation	UNP Q99986
C	212	ALA	GLU	engineered mutation	UNP Q99986
C	214	ALA	LYS	engineered mutation	UNP Q99986
C	215	ALA	GLU	engineered mutation	UNP Q99986
C	292	ALA	GLU	engineered mutation	UNP Q99986
C	293	ALA	LYS	engineered mutation	UNP Q99986
C	295	ALA	LYS	engineered mutation	UNP Q99986
C	359	ALA	LYS	engineered mutation	UNP Q99986
C	360	ALA	LYS	engineered mutation	UNP Q99986
D	1	SER	-	expression tag	UNP Q99986
D	2	MET	-	expression tag	UNP Q99986
D	34	ALA	LYS	engineered mutation	UNP Q99986
D	35	ALA	LYS	engineered mutation	UNP Q99986
D	36	ALA	GLU	engineered mutation	UNP Q99986
D	212	ALA	GLU	engineered mutation	UNP Q99986
D	214	ALA	LYS	engineered mutation	UNP Q99986
D	215	ALA	GLU	engineered mutation	UNP Q99986
D	292	ALA	GLU	engineered mutation	UNP Q99986
D	293	ALA	LYS	engineered mutation	UNP Q99986
D	295	ALA	LYS	engineered mutation	UNP Q99986
D	359	ALA	LYS	engineered mutation	UNP Q99986
D	360	ALA	LYS	engineered mutation	UNP Q99986

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	3	Total Cl 3 3	0	0
2	A	2	Total Cl 2 2	0	0
2	D	3	Total Cl 3 3	0	0
2	C	1	Total Cl 1 1	0	0

- Molecule 3 is 2-[(3,5-difluoro-4-hydroxyphenyl)amino]-8-phenyl-7,8-dihydropteridin-6(5H)-one (three-letter code: E8D) (formula: $C_{18}H_{13}F_2N_5O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			27	18	2	5	2		
3	B	1	Total	C	F	N	O	0	0
			27	18	2	5	2		
3	D	1	Total	C	F	N	O	0	0
			27	18	2	5	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total O S 5 4 1	0	0


- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	261	Total O 261 261	0	0
6	B	191	Total O 191 191	0	0
6	C	276	Total O 276 276	0	0
6	D	251	Total O 251 251	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.58 Å 97.56 Å 193.28 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.89 – 1.90 19.89 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.89-1.90) 99.8 (19.89-1.90)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.90 Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.182 , 0.215 0.192 , 0.222	Depositor DCC
R_{free} test set	6723 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.547	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10921	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, E8D, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2577	0.77	0/3493
1	B	0.59	0/2450	0.73	0/3317
1	C	0.67	0/2566	0.75	0/3478
1	D	0.63	0/2485	0.72	0/3369
All	All	0.64	0/10078	0.74	0/13657

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2515	0	2432	4	0
1	B	2391	0	2320	1	0
1	C	2504	0	2433	3	0
1	D	2425	0	2347	4	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	1	0	0	0	0
2	D	3	0	0	0	0
3	A	27	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	27	0	0	0	0
3	D	27	0	0	1	0
4	C	5	0	0	0	0
5	C	12	0	16	1	0
6	A	261	0	0	1	0
6	B	191	0	0	0	0
6	C	276	0	0	0	0
6	D	251	0	0	0	0
All	All	10921	0	9548	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:LYS:NZ	6:A:501:HOH:O	2.27	0.62
1:C:207:GLU:OE2	5:C:403:GOL:O2	2.18	0.61
1:D:134:PHE:HB2	1:D:184:LEU:HD23	1.87	0.55
1:B:178:ILE:HD12	1:B:178:ILE:H	1.76	0.50
1:A:87:TYR:CZ	1:A:131:MET:HE2	2.50	0.46
1:D:333:SER:OG	1:D:334:LYS:N	2.47	0.46
1:D:51:ILE:HD12	3:D:404:E8D:C15	2.46	0.46
1:A:178:ILE:HD12	1:A:178:ILE:H	1.81	0.44
1:A:204:TYR:CE2	1:A:205:CYS:HB2	2.53	0.43
1:D:204:TYR:CE2	1:D:205:CYS:HB2	2.55	0.42
1:C:200:LEU:HD13	1:C:221:HIS:CG	2.56	0.41
1:C:204:TYR:CE2	1:C:205:CYS:HB2	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/364 (86%)	309 (98%)	6 (2%)	0	100	100
1	B	292/364 (80%)	286 (98%)	6 (2%)	0	100	100
1	C	313/364 (86%)	306 (98%)	7 (2%)	0	100	100
1	D	300/364 (82%)	292 (97%)	8 (3%)	0	100	100
All	All	1220/1456 (84%)	1193 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/304 (86%)	259 (100%)	1 (0%)	91	91
1	B	242/304 (80%)	241 (100%)	1 (0%)	91	91
1	C	259/304 (85%)	258 (100%)	1 (0%)	91	91
1	D	246/304 (81%)	245 (100%)	1 (0%)	91	91
All	All	1007/1216 (83%)	1003 (100%)	4 (0%)	91	91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	ARG
1	B	133	ARG
1	C	133	ARG
1	D	133	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 9 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	GOL	C	404	-	5,5,5	0.85	0	5,5,5	1.30	0
3	E8D	A	403	-	29,30,30	0.64	0	35,43,43	1.26	4 (11%)
3	E8D	B	404	-	29,30,30	0.52	0	35,43,43	1.16	4 (11%)
4	SO4	C	401	-	4,4,4	0.31	0	6,6,6	0.67	0
5	GOL	C	403	-	5,5,5	0.18	0	5,5,5	0.66	0
3	E8D	D	404	-	29,30,30	0.59	0	35,43,43	1.15	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	403	-	-	4/4/4/4	-
3	E8D	A	403	-	-	0/8/20/20	0/4/4/4
3	E8D	D	404	-	-	0/8/20/20	0/4/4/4
5	GOL	C	404	-	-	0/4/4/4	-
3	E8D	B	404	-	-	0/8/20/20	0/4/4/4

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	E8D	F2-C10-C11	3.59	119.79	117.13
3	D	404	E8D	F1-C12-C11	3.25	119.55	117.13
3	A	403	E8D	N2-C2-N3	-3.23	123.49	126.55
3	B	404	E8D	F2-C10-C11	2.79	119.20	117.13
3	B	404	E8D	C9-C10-C11	-2.62	121.74	123.79
3	B	404	E8D	C13-C12-C11	-2.53	121.81	123.79
3	D	404	E8D	C9-C10-C11	-2.39	121.92	123.79
3	A	403	E8D	C3-C4-N4	2.16	123.14	119.99
3	D	404	E8D	F2-C10-C11	2.12	118.70	117.13
3	B	404	E8D	F1-C12-C11	2.09	118.69	117.13
3	A	403	E8D	F1-C12-C11	2.03	118.64	117.13

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	403	GOL	O1-C1-C2-C3
5	C	403	GOL	O1-C1-C2-O2
5	C	403	GOL	O2-C2-C3-O3
5	C	403	GOL	C1-C2-C3-O3


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	403	GOL	1	0
3	D	404	E8D	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand E8D A 403



Ligand E8D B 404





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/364 (87%)	0.10	14 (4%) 34 37	17, 26, 43, 66	0
1	B	301/364 (82%)	0.33	23 (7%) 13 15	15, 30, 59, 72	0
1	C	315/364 (86%)	0.19	14 (4%) 34 37	16, 28, 50, 60	0
1	D	307/364 (84%)	0.31	28 (9%) 9 10	14, 27, 62, 79	0
All	All	1240/1456 (85%)	0.23	79 (6%) 19 22	14, 28, 54, 79	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	126	TYR	5.7
1	C	220	CYS	5.7
1	D	122	ASN	5.2
1	B	34	ALA	5.0
1	C	217	PRO	4.8
1	D	62	VAL	4.7
1	D	293	ALA	4.3
1	A	220	CYS	4.3
1	A	217	PRO	4.2
1	D	126	TYR	4.0
1	D	50	CYS	4.0
1	D	123	GLY	4.0
1	B	58	SER	3.7
1	C	187	TYR	3.7
1	B	187	TYR	3.6
1	A	49	GLY	3.5
1	C	122	ASN	3.5
1	D	65	ASP	3.4
1	B	74	PRO	3.3
1	C	59	SER	3.3
1	B	120	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	42	PRO	3.2
1	C	146	ALA	3.2
1	C	64	SER	3.2
1	B	66	ALA	3.2
1	A	123	GLY	3.1
1	C	332	GLY	3.1
1	B	57	ASN	3.1
1	D	61	SER	3.0
1	D	145	ASN	3.0
1	A	20	ALA	2.9
1	B	78	GLY	2.8
1	B	263	ASP	2.7
1	D	56	MET	2.7
1	D	64	SER	2.7
1	C	123	GLY	2.6
1	A	59	SER	2.6
1	D	263	ASP	2.6
1	D	43	ILE	2.6
1	D	125	SER	2.5
1	D	41	LEU	2.5
1	B	35	ALA	2.4
1	B	36	ALA	2.4
1	B	144	ALA	2.4
1	A	293	ALA	2.4
1	D	146	ALA	2.4
1	D	25	VAL	2.4
1	A	60	GLU	2.4
1	B	65	ASP	2.4
1	B	64	SER	2.4
1	D	63	GLY	2.4
1	B	24	ALA	2.4
1	A	64	SER	2.4
1	D	120	ASP	2.4
1	D	294	ASN	2.4
1	B	25	VAL	2.3
1	D	57	ASN	2.3
1	B	79	PRO	2.3
1	C	65	ASP	2.3
1	C	215	ALA	2.3
1	C	50	CYS	2.2
1	B	81	PHE	2.2
1	D	28	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	ALA	2.1
1	A	146	ALA	2.1
1	C	120	ASP	2.1
1	D	33	ALA	2.1
1	A	291	PRO	2.1
1	B	33	ALA	2.1
1	B	294	ASN	2.1
1	D	66	ALA	2.1
1	D	124	LYS	2.1
1	C	25	VAL	2.1
1	A	65	ASP	2.1
1	D	42	PRO	2.1
1	B	23	PHE	2.1
1	A	215	ALA	2.1
1	D	144	ALA	2.1
1	D	130	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	D	403	1/1	0.71	0.08	72,72,72,72	0
3	E8D	A	403	27/27	0.71	0.24	36,52,73,75	0
5	GOL	C	404	6/6	0.78	0.20	39,44,45,48	0
2	CL	B	403	1/1	0.86	0.06	58,58,58,58	0
2	CL	B	401	1/1	0.88	0.08	63,63,63,63	0
5	GOL	C	403	6/6	0.88	0.14	30,40,43,43	0
2	CL	A	402	1/1	0.89	0.13	65,65,65,65	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	C	402	1/1	0.90	0.12	66,66,66,66	0
3	E8D	B	404	27/27	0.91	0.12	36,43,50,56	0
2	CL	A	401	1/1	0.91	0.10	62,62,62,62	0
2	CL	D	401	1/1	0.92	0.14	48,48,48,48	0
3	E8D	D	404	27/27	0.92	0.12	24,35,48,50	0
2	CL	B	402	1/1	0.93	0.07	58,58,58,58	0
4	SO4	C	401	5/5	0.97	0.12	45,45,52,54	0
2	CL	D	402	1/1	0.98	0.11	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around E8D A 403:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E8D B 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around E8D D 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.