



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 06:04 AM BST

PDB ID : 6BUR
Title : Crystal structures of cyanuric acid hydrolase from *Moorella thermoacetica* complexed with barbituric acid
Authors : Shi, K.; Aihara, H.
Deposited on : 2017-12-11
Resolution : 2.18 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	<div> <div>10%</div> <div>90%</div> <div>10%</div> </div>
1	B	363	<div> <div>89%</div> <div>10%</div> <div>•</div> </div>
1	C	363	<div> <div>88%</div> <div>11%</div> <div>•</div> </div>
1	D	363	<div> <div>10%</div> <div>73%</div> <div>25%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR8	A	401[A]	-	X	-	-
2	BR8	B	401	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11266 atoms, of which 18 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyanuric acid amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	2	0
			2697	1668	481	533	15			
1	B	363	Total	C	N	O	S	0	5	0
			2736	1692	490	539	15			
1	C	362	Total	C	N	O	S	0	0	0
			2681	1659	479	529	14			
1	D	362	Total	C	N	O	S	0	0	0
			2681	1659	479	529	14			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q2RGM7
A	103	ALA	GLN	engineered mutation	UNP Q2RGM7
A	104	ALA	GLU	engineered mutation	UNP Q2RGM7
A	107	ALA	LYS	engineered mutation	UNP Q2RGM7
A	279	ILE	LEU	engineered mutation	UNP Q2RGM7
A	280	ARG	LYS	engineered mutation	UNP Q2RGM7
A	281	SER	PHE	engineered mutation	UNP Q2RGM7
A	?	-	CYS	deletion	UNP Q2RGM7
A	?	-	CYS	deletion	UNP Q2RGM7
A	?	-	PRO	deletion	UNP Q2RGM7
A	?	-	PRO	deletion	UNP Q2RGM7
A	?	-	ALA	deletion	UNP Q2RGM7
A	283	ASP	GLU	engineered mutation	UNP Q2RGM7
A	290	MET	LEU	engineered mutation	UNP Q2RGM7
A	291	ASP	ALA	engineered mutation	UNP Q2RGM7
A	292	ARG	LYS	engineered mutation	UNP Q2RGM7
B	0	HIS	-	expression tag	UNP Q2RGM7
B	103	ALA	GLN	engineered mutation	UNP Q2RGM7
B	104	ALA	GLU	engineered mutation	UNP Q2RGM7
B	107	ALA	LYS	engineered mutation	UNP Q2RGM7
B	279	ILE	LEU	engineered mutation	UNP Q2RGM7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	280	ARG	LYS	engineered mutation	UNP Q2RGM7
B	281	SER	PHE	engineered mutation	UNP Q2RGM7
B	?	-	CYS	deletion	UNP Q2RGM7
B	?	-	CYS	deletion	UNP Q2RGM7
B	?	-	PRO	deletion	UNP Q2RGM7
B	?	-	PRO	deletion	UNP Q2RGM7
B	?	-	ALA	deletion	UNP Q2RGM7
B	283	ASP	GLU	engineered mutation	UNP Q2RGM7
B	290	MET	LEU	engineered mutation	UNP Q2RGM7
B	291	ASP	ALA	engineered mutation	UNP Q2RGM7
B	292	ARG	LYS	engineered mutation	UNP Q2RGM7
C	0	HIS	-	expression tag	UNP Q2RGM7
C	103	ALA	GLN	engineered mutation	UNP Q2RGM7
C	104	ALA	GLU	engineered mutation	UNP Q2RGM7
C	107	ALA	LYS	engineered mutation	UNP Q2RGM7
C	279	ILE	LEU	engineered mutation	UNP Q2RGM7
C	280	ARG	LYS	engineered mutation	UNP Q2RGM7
C	281	SER	PHE	engineered mutation	UNP Q2RGM7
C	?	-	CYS	deletion	UNP Q2RGM7
C	?	-	CYS	deletion	UNP Q2RGM7
C	?	-	PRO	deletion	UNP Q2RGM7
C	?	-	PRO	deletion	UNP Q2RGM7
C	?	-	ALA	deletion	UNP Q2RGM7
C	283	ASP	GLU	engineered mutation	UNP Q2RGM7
C	290	MET	LEU	engineered mutation	UNP Q2RGM7
C	291	ASP	ALA	engineered mutation	UNP Q2RGM7
C	292	ARG	LYS	engineered mutation	UNP Q2RGM7
D	0	HIS	-	expression tag	UNP Q2RGM7
D	103	ALA	GLN	engineered mutation	UNP Q2RGM7
D	104	ALA	GLU	engineered mutation	UNP Q2RGM7
D	107	ALA	LYS	engineered mutation	UNP Q2RGM7
D	279	ILE	LEU	engineered mutation	UNP Q2RGM7
D	280	ARG	LYS	engineered mutation	UNP Q2RGM7
D	281	SER	PHE	engineered mutation	UNP Q2RGM7
D	?	-	CYS	deletion	UNP Q2RGM7
D	?	-	CYS	deletion	UNP Q2RGM7
D	?	-	PRO	deletion	UNP Q2RGM7
D	?	-	PRO	deletion	UNP Q2RGM7
D	?	-	ALA	deletion	UNP Q2RGM7
D	283	ASP	GLU	engineered mutation	UNP Q2RGM7
D	290	MET	LEU	engineered mutation	UNP Q2RGM7
D	291	ASP	ALA	engineered mutation	UNP Q2RGM7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	292	ARG	LYS	engineered mutation	UNP Q2RGM7

- Molecule 2 is BARBITURIC ACID (three-letter code: BR8) (formula: $C_4H_4N_2O_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	1
			13	4	4	2	3		
2	B	1	Total	C	H	N	O	0	0
			13	4	4	2	3		
2	C	1	Total	C	H	N	O	0	0
			13	4	4	2	3		
2	D	1	Total	C	H	N	O	0	0
			13	4	4	2	3		

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	1
			9	3	2	4		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	131	Total	O	0	0
			131	131		
5	B	116	Total	O	0	0
			116	116		
5	C	108	Total	O	0	0
			108	108		
5	D	51	Total	O	0	0
			51	51		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyanuric acid amidohydrolase




- Molecule 1: Cyanuric acid amidohydrolase




- Molecule 1: Cyanuric acid amidohydrolase



- Molecule 1: Cyanuric acid amidohydrolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.08Å 89.12Å 199.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.72 – 2.18 81.35 – 2.18	Depositor EDS
% Data completeness (in resolution range)	99.1 (39.72-2.18) 99.1 (81.35-2.18)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.18Å)	Xtriage
Refinement program	PHENIX (1.13rc2_2981: ???)	Depositor
R, R_{free}	0.163 , 0.212 0.165 , 0.210	Depositor DCC
R_{free} test set	3795 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.165	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11266	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MLI, BR8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2729	0.54	0/3693
1	B	0.43	0/2769	0.56	0/3746
1	C	0.38	0/2713	0.54	0/3672
1	D	0.32	0/2713	0.49	0/3672
All	All	0.40	0/10924	0.53	0/14783

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2697	0	2716	28	0
1	B	2736	0	2753	39	0
1	C	2681	0	2705	30	0
1	D	2681	0	2705	80	0
2	A	9	4	4	1	0
2	B	9	4	4	1	0
2	C	9	4	4	0	0
2	D	9	4	4	0	0
3	A	7	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	131	0	0	5	0
5	B	116	0	0	0	0
5	C	108	0	0	0	0
5	D	51	0	0	1	0
All	All	11248	18	10897	169	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 169 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:127:LEU:HD11	1:D:163:PRO:HG2	1.48	0.94
1:C:171:GLU:OE1	1:C:175:ARG:NH2	2.04	0.90
1:A:71:ARG:NH2	1:B:65:GLU:OE2	2.10	0.84
1:D:275:LYS:HE3	1:D:280:ARG:HA	1.59	0.83
1:D:275:LYS:HG3	1:D:280:ARG:HG3	1.61	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/363 (100%)	348 (96%)	14 (4%)	0	100	100
1	B	366/363 (101%)	353 (96%)	13 (4%)	0	100	100
1	C	360/363 (99%)	347 (96%)	13 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	360/363 (99%)	346 (96%)	14 (4%)	0	100	100
All	All	1448/1452 (100%)	1394 (96%)	54 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/286 (100%)	284 (99%)	3 (1%)	76	85
1	B	291/286 (102%)	285 (98%)	6 (2%)	53	64
1	C	285/286 (100%)	282 (99%)	3 (1%)	73	83
1	D	285/286 (100%)	276 (97%)	9 (3%)	39	47
All	All	1148/1144 (100%)	1127 (98%)	21 (2%)	59	70

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	76	LYS
1	C	239	ASP
1	D	210	ASP
1	B	290	MET
1	D	237	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	249	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	MLI	A	402[B]	-	0,6,6	0.00	-	0,7,7	0.00	-
2	BR8	C	401	-	9,9,9	3.55	5 (55%)	12,12,12	3.21	4 (33%)
2	BR8	D	401	-	9,9,9	3.60	6 (66%)	12,12,12	3.59	6 (50%)
2	BR8	B	401	-	9,9,9	3.63	7 (77%)	12,12,12	3.93	7 (58%)
2	BR8	A	401[A]	-	9,9,9	4.11	7 (77%)	12,12,12	3.88	7 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	A	402[B]	-	-	0/0/4/4	-
2	BR8	C	401	-	-	-	0/1/1/1
2	BR8	D	401	-	-	-	0/1/1/1
2	BR8	B	401	-	-	-	0/1/1/1
2	BR8	A	401[A]	-	-	-	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[A]	BR8	C2-N1	7.00	1.49	1.37
2	A	401[A]	BR8	C2-N3	6.31	1.48	1.37
2	B	401	BR8	C2-N3	6.14	1.48	1.37
2	C	401	BR8	C2-N1	6.02	1.47	1.37
2	D	401	BR8	C2-N3	5.96	1.47	1.37

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[A]	BR8	C4-N3-C2	-8.13	119.20	125.73
2	B	401	BR8	C6-N1-C2	-8.01	119.29	125.73
2	D	401	BR8	C6-N1-C2	-7.98	119.32	125.73
2	A	401[A]	BR8	C6-N1-C2	-7.62	119.60	125.73
2	B	401	BR8	C4-N3-C2	-7.44	119.75	125.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	BR8	1	0
2	A	401[A]	BR8	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/363 (99%)	-0.24	4 (1%) 80 80	26, 40, 70, 142	0
1	B	363/363 (100%)	-0.33	0 100 100	27, 42, 68, 92	0
1	C	362/363 (99%)	-0.23	2 (0%) 89 89	29, 50, 81, 151	0
1	D	362/363 (99%)	0.49	36 (9%) 7 7	33, 74, 129, 164	0
All	All	1449/1452 (99%)	-0.08	42 (2%) 51 52	26, 47, 98, 164	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	178	LYS	7.9
1	D	177	GLY	7.4
1	D	242	ILE	5.6
1	D	106	ALA	5.6
1	A	107	ALA	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BR8	A	401[A]	9/9	0.95	0.10	32,35,42,42	13
4	CA	D	402	1/1	0.97	0.05	66,66,66,66	0
2	BR8	B	401	9/9	0.97	0.10	29,31,38,39	0
2	BR8	C	401	9/9	0.97	0.10	32,35,41,43	0
2	BR8	D	401	9/9	0.98	0.09	42,45,53,55	0
3	MLI	A	402[B]	7/7	0.98	0.10	30,35,42,42	9
4	CA	C	402	1/1	0.99	0.07	37,37,37,37	1
4	CA	A	403	1/1	1.00	0.12	37,37,37,37	1
4	CA	B	402	1/1	1.00	0.07	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.