



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 03:12 PM BST

PDB ID : 6BUY  
Title : Crystal structure of porcine aminopeptidase-N with Glycine  
Authors : Chen, L.; Lin, Y.-L.; Li, F.  
Deposited on : 2017-12-11  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	902	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div></div> </div> </div>
2	B	3	<div> <div>67%</div> <div>33%</div> </div>
2	C	3	<div> <div>33%</div> <div>67%</div> </div>
2	F	3	<div> <div>67%</div> <div>33%</div> </div>
3	D	2	<div> <div>100%</div> </div>
3	E	2	<div> <div>100%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	2	 100%
3	H	2	 100%
3	I	2	 50%50%
3	J	2	 50%50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	3	-	-	-	X
2	NAG	F	3	-	-	-	X
4	NAG	A	1020	-	-	-	X
6	SO4	A	1025	-	X	-	-
6	SO4	A	1026	-	X	-	-
6	SO4	A	1028	-	X	-	-
6	SO4	A	1033	-	X	-	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	902	Total	C	N	O	S	0	0	0
			7241	4622	1210	1379	30			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	PHE	LEU	conflict	UNP P15145
A	964	SER	-	expression tag	UNP P15145

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			42	24	3	15			
2	C	3	Total	C	N	O	0	0	0
			42	24	3	15			
2	F	3	Total	C	N	O	0	0	0
			42	24	3	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is GLYCINE (three-letter code: GLY) (formula:  $C_2H_5NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 6 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	932	Total	O	0	0
			932	932		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminopeptidase N



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose







- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	260.61Å 62.73Å 81.64Å 90.00° 100.26° 90.00°	Depositor
Resolution (Å)	128.22 – 2.10 47.59 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (128.22-2.10) 99.8 (47.59-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.11 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.197 , 0.248 0.202 , 0.253	Depositor DCC
$R_{free}$ test set	3817 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.9	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 35.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8551	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	17/7429 (0.2%)	0.74	6/10124 (0.1%)

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	111	GLU	CD-OE2	9.75	1.36	1.25
1	A	766	GLU	CD-OE1	-7.26	1.17	1.25
1	A	455	GLU	CD-OE2	7.06	1.33	1.25
1	A	375	GLU	CD-OE1	-6.83	1.18	1.25
1	A	455	GLU	CD-OE1	6.68	1.32	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	637	ARG	NE-CZ-NH2	7.18	123.89	120.30
1	A	533	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	A	437	ARG	NE-CZ-NH1	-5.80	117.40	120.30
1	A	147	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	A	296	ARG	NE-CZ-NH2	-5.22	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7241	0	6994	101	0
2	B	42	0	37	3	0
2	C	42	0	37	1	0
2	F	42	0	37	1	0
3	D	28	0	25	0	0
3	E	28	0	25	0	0
3	G	28	0	25	0	0
3	H	28	0	25	0	0
3	I	28	0	25	4	0
3	J	28	0	25	1	0
4	A	28	0	26	0	0
5	A	5	0	2	2	0
6	A	50	0	0	2	1
7	A	1	0	0	0	0
8	A	932	0	0	24	4
All	All	8551	0	7283	112	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:820:ASN:HB3	8:A:1617:HOH:O	1.41	1.19
1:A:556:ASN:HD22	3:I:1:NAG:H83	1.23	1.03
1:A:380:VAL:HG13	8:A:1187:HOH:O	1.63	0.98
1:A:779:ASP:OD2	8:A:1101:HOH:O	1.88	0.92
1:A:69:ARG:NH1	6:A:1034:SO4:O2	2.01	0.92

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1697:HOH:O	8:A:2032:HOH:O[4_557]	1.26	0.94
8:A:1874:HOH:O	8:A:2031:HOH:O[4_557]	1.41	0.79
6:A:1029:SO4:O1	8:A:1352:HOH:O[2_556]	2.10	0.10
8:A:1641:HOH:O	8:A:1949:HOH:O[2_556]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	900/902 (100%)	873 (97%)	21 (2%)	6 (1%)	22 18

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	93	ALA
1	A	882	LYS
1	A	892	SER
1	A	893	PHE
1	A	923	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	797/797 (100%)	772 (97%)	25 (3%)	40 43

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	644	GLN
1	A	738	THR
1	A	932	ARG
1	A	723	VAL
1	A	764	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	728	GLN
1	A	913	GLN
1	A	897	ASN
1	A	448	HIS
1	A	782	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	B	1	1,2	14,14,15	1.04	2 (14%)	17,19,21	1.07	1 (5%)
2	NAG	B	2	2	14,14,15	1.10	1 (7%)	17,19,21	1.58	3 (17%)
2	NAG	B	3	2	14,14,15	1.19	1 (7%)	17,19,21	1.62	3 (17%)
2	NAG	C	1	1,2	14,14,15	0.67	0	17,19,21	1.34	2 (11%)
2	NAG	C	2	2	14,14,15	1.20	2 (14%)	17,19,21	2.07	6 (35%)
2	NAG	C	3	2	14,14,15	1.29	1 (7%)	17,19,21	1.45	3 (17%)
3	NAG	D	1	1,3	14,14,15	0.94	0	17,19,21	1.24	1 (5%)
3	NAG	D	2	3	14,14,15	0.96	0	17,19,21	1.55	4 (23%)
3	NAG	E	1	1,3	14,14,15	0.77	0	17,19,21	1.07	1 (5%)
3	NAG	E	2	3	14,14,15	0.79	0	17,19,21	1.51	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	F	1	1,2	14,14,15	0.62	0	17,19,21	1.38	2 (11%)
2	NAG	F	2	2	14,14,15	0.94	0	17,19,21	1.66	5 (29%)
2	NAG	F	3	2	14,14,15	1.51	1 (7%)	17,19,21	1.56	4 (23%)
3	NAG	G	1	1,3	14,14,15	1.29	2 (14%)	17,19,21	0.97	0
3	NAG	G	2	3	14,14,15	0.97	1 (7%)	17,19,21	1.21	1 (5%)
3	NAG	H	1	1,3	14,14,15	1.03	1 (7%)	17,19,21	1.20	1 (5%)
3	NAG	H	2	3	14,14,15	0.76	0	17,19,21	2.27	5 (29%)
3	NAG	I	1	1,3	14,14,15	1.21	1 (7%)	17,19,21	1.62	6 (35%)
3	NAG	I	2	3	14,14,15	1.23	1 (7%)	17,19,21	1.29	2 (11%)
3	NAG	J	1	1,3	14,14,15	0.73	0	17,19,21	1.20	1 (5%)
3	NAG	J	2	3	14,14,15	1.40	2 (14%)	17,19,21	1.68	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	4/6/23/26	0/1/1/1
2	NAG	B	3	2	-	4/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	3	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	NAG	F	3	2	-	3/6/23/26	0/1/1/1
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	NAG	H	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	4/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	I	2	3	-	3/6/23/26	0/1/1/1
3	NAG	J	1	1,3	-	0/6/23/26	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	J	2	3	-	0/6/23/26	0/1/1/1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	3	NAG	O7-C7	3.81	1.31	1.23
3	I	1	NAG	C8-C7	3.35	1.57	1.50
3	G	1	NAG	O5-C1	3.18	1.48	1.43
3	J	2	NAG	O5-C5	3.08	1.49	1.43
2	C	3	NAG	O7-C7	2.82	1.29	1.23

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	2	NAG	O5-C5-C6	5.34	115.58	107.20
2	C	2	NAG	C4-C3-C2	4.82	118.08	111.02
2	B	3	NAG	C4-C3-C2	4.40	117.47	111.02
3	H	2	NAG	C2-N2-C7	-4.16	116.99	122.90
3	J	2	NAG	O5-C1-C2	-4.05	104.89	111.29

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	3	NAG	C3-C2-N2-C7
2	B	3	NAG	C3-C2-N2-C7
3	G	2	NAG	C4-C5-C6-O6
2	F	3	NAG	C4-C5-C6-O6
2	C	3	NAG	O5-C5-C6-O6


There are no ring outliers.


6 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	1	0
2	C	3	NAG	1	0
2	B	3	NAG	3	0
3	J	2	NAG	1	0
2	F	2	NAG	1	0
3	I	1	NAG	4	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

























## 5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	1020	1	14,14,15	1.31	1 (7%)	17,19,21	1.47	2 (11%)
6	SO4	A	1034	-	4,4,4	2.13	2 (50%)	6,6,6	1.06	0
6	SO4	A	1026	-	4,4,4	1.95	2 (50%)	6,6,6	2.24	3 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	SO4	A	1030	-	4,4,4	1.46	1 (25%)	6,6,6	0.61	0
6	SO4	A	1029	-	4,4,4	1.88	1 (25%)	6,6,6	0.78	0
6	SO4	A	1033	-	4,4,4	2.32	3 (75%)	6,6,6	1.73	2 (33%)
6	SO4	A	1027	-	4,4,4	2.57	3 (75%)	6,6,6	1.28	0
6	SO4	A	1032	-	4,4,4	2.44	3 (75%)	6,6,6	1.32	0
6	SO4	A	1025	-	4,4,4	2.73	2 (50%)	6,6,6	3.59	4 (66%)
6	SO4	A	1028	-	4,4,4	2.86	3 (75%)	6,6,6	1.55	1 (16%)
4	NAG	A	1023	1	14,14,15	1.24	0	17,19,21	1.41	3 (17%)
6	SO4	A	1031	-	4,4,4	2.51	2 (50%)	6,6,6	1.68	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1020	1	-	3/6/23/26	0/1/1/1
4	NAG	A	1023	1	-	1/6/23/26	0/1/1/1

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1025	SO4	O2-S	4.79	1.72	1.46
6	A	1028	SO4	O1-S	4.34	1.69	1.46
6	A	1031	SO4	O2-S	3.70	1.66	1.46
6	A	1027	SO4	O2-S	3.61	1.65	1.46
6	A	1032	SO4	O1-S	3.50	1.65	1.46

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1025	SO4	O4-S-O3	5.89	134.21	109.06
6	A	1025	SO4	O4-S-O2	-4.94	83.53	109.31
4	A	1020	NAG	C1-O5-C5	-3.37	107.62	112.19
6	A	1031	SO4	O4-S-O2	3.33	126.70	109.31
6	A	1025	SO4	O3-S-O1	-3.28	92.20	109.31

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1020	NAG	O5-C5-C6-O6
4	A	1020	NAG	C4-C5-C6-O6
4	A	1020	NAG	C3-C2-N2-C7
4	A	1023	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1034	SO4	1	0
6	A	1026	SO4	1	0
6	A	1029	SO4	0	1

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	902/902 (100%)	0.27	41 (4%) 33 38	24, 32, 41, 45	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	889	GLY	9.2
1	A	885	PHE	9.1
1	A	888	TYR	8.4
1	A	925	VAL	7.6
1	A	890	GLY	7.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	3	14/15	0.52	0.51	33,35,39,41	0
2	NAG	F	3	14/15	0.61	0.51	32,35,39,39	0
2	NAG	F	2	14/15	0.67	0.28	30,34,37,37	0
3	NAG	G	1	14/15	0.82	0.26	32,35,36,36	0
3	NAG	G	2	14/15	0.82	0.36	31,34,40,41	0
2	NAG	C	2	14/15	0.84	0.42	32,34,39,39	0
2	NAG	B	3	14/15	0.84	0.32	33,34,39,40	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	E	2	14/15	0.84	0.31	31,34,40,41	0
3	NAG	J	2	14/15	0.85	0.27	33,35,37,40	0
3	NAG	H	2	14/15	0.85	0.26	31,33,37,39	0
3	NAG	I	2	14/15	0.86	0.32	32,34,38,40	0
2	NAG	B	2	14/15	0.90	0.24	31,34,38,38	0
3	NAG	E	1	14/15	0.91	0.18	31,33,38,38	0
3	NAG	D	2	14/15	0.91	0.23	31,32,35,38	0
3	NAG	I	1	14/15	0.91	0.22	31,34,38,42	0
2	NAG	C	1	14/15	0.92	0.16	30,32,35,37	0
2	NAG	F	1	14/15	0.92	0.20	30,33,34,35	0
3	NAG	H	1	14/15	0.92	0.12	28,31,32,33	0
3	NAG	J	1	14/15	0.93	0.09	31,33,37,38	0
3	NAG	D	1	14/15	0.96	0.10	28,31,33,33	0
2	NAG	B	1	14/15	0.96	0.10	28,31,32,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain D:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around Chain E:**




$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain H:**


$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	1020	14/15	0.75	0.54	33,35,40,41	0
6	SO4	A	1028	5/5	0.81	0.38	34,36,37,39	0
5	GLY	A	1024	5/5	0.83	0.21	29,30,34,36	0
6	SO4	A	1027	5/5	0.84	0.41	34,37,39,40	0
4	NAG	A	1023	14/15	0.84	0.41	31,34,41,42	0
6	SO4	A	1032	5/5	0.88	0.42	33,38,38,38	0
6	SO4	A	1025	5/5	0.88	0.27	30,32,35,38	0
6	SO4	A	1034	5/5	0.89	0.52	36,37,38,40	0
6	SO4	A	1033	5/5	0.90	0.45	32,32,39,41	0
6	SO4	A	1031	5/5	0.93	0.42	33,34,38,38	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	SO4	A	1026	5/5	0.97	0.49	30,35,36,37	0
6	SO4	A	1029	5/5	0.97	0.29	35,36,38,38	0
6	SO4	A	1030	5/5	0.98	0.16	33,33,35,36	0
7	ZN	A	1035	1/1	1.00	0.15	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.