



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 09:21 PM BST

PDB ID : 4BWC  
Title : X-ray structure of a phospholipase B like protein 1 from bovine kidneys  
Authors : Repo, H.; Kuokkanen, E.; Oksanen, E.; Goldman, A.; Heikinheimo, P.  
Deposited on : 2013-07-01  
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>9%</div> <div> <div></div> <div>85%</div> <div>14%</div> </div> <div>•</div> </div>
2	B	321	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>13%</div> </div> <div>•</div> </div>
3	C	2	<div> <div>50%</div> <div>50%</div> </div>
3	D	2	<div> <div>100%</div> </div>
3	E	2	<div> <div>100%</div> </div>

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOLIPASE B-LIKE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	0	0	0
			1376	893	226	250	7			

- Molecule 2 is a protein called PHOSPHOLIPASE B-LIKE 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	321	Total	C	N	O	S	0	2	0
			2631	1699	431	487	14			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 1-ETHOXY-2-(2-ETHOXYETHOXY)ETHANE (three-letter code: P4G) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	8	3		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula:  $C_{12}H_{26}O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			19	12	7		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	50	Total	O	0	0
			50	50		
8	B	134	Total	O	0	0
			134	134		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PHOSPHOLIPASE B-LIKE 1




#### • Molecule 2: PHOSPHOLIPASE B-LIKE 1



#### • Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



#### • Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



#### • Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

100%

EM61  
EM62

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.37Å 97.37Å 140.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.98 – 1.89 46.98 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.98-1.89) 99.8 (46.98-1.89)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.188 , 0.231 0.187 , 0.230	Depositor DCC
$R_{free}$ test set	3127 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.7	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4334	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P4G, OCS, NAG, P6G, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.01	0/1412	0.91	1/1909 (0.1%)
2	B	1.27	6/2705 (0.2%)	1.10	5/3673 (0.1%)
All	All	1.19	6/4117 (0.1%)	1.04	6/5582 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	320	TRP	CD2-CE2	7.95	1.50	1.41
2	B	409	TRP	CD2-CE2	6.44	1.49	1.41
2	B	425	TYR	CE2-CZ	5.66	1.46	1.38
2	B	519	TRP	CD2-CE2	5.64	1.48	1.41
2	B	482	SER	CB-OG	-5.27	1.35	1.42
2	B	336	TRP	CD2-CE2	5.02	1.47	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	443	ASP	CB-CG-OD1	9.29	126.67	118.30
2	B	322	ARG	NE-CZ-NH2	-6.57	117.02	120.30
2	B	435	ARG	NE-CZ-NH1	6.37	123.49	120.30
1	A	122	LEU	CB-CG-CD2	-5.95	100.89	111.00
2	B	328	MET	CG-SD-CE	-5.86	90.82	100.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	246	TYR	CB-CG-CD2	-5.36	117.79	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	76	TRP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1376	0	1356	22	0
2	B	2631	0	2551	21	0
3	C	28	0	25	1	0
3	D	28	0	25	0	0
3	E	28	0	25	1	0
4	A	11	0	18	0	0
5	B	28	0	26	1	0
6	B	19	0	26	3	0
7	B	1	0	0	0	0
8	A	50	0	0	2	0
8	B	134	0	0	2	0
All	All	4334	0	4052	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:LYS:H	1:A:128:MET:CE	1.90	0.85
2:B:304:TYR:CD2	2:B:415:LEU:HD11	2.17	0.79
2:B:225:OCS:OD1	2:B:225:OCS:N	2.15	0.78
2:B:498:SER:HA	6:B:1546:P6G:H122	1.68	0.75

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:SER:HB3	8:A:2031:HOH:O	1.88	0.72
1:A:124:LYS:H	1:A:128:MET:HE1	1.54	0.70
1:A:124:LYS:H	1:A:128:MET:HE3	1.57	0.69
1:A:125:LYS:H	1:A:128:MET:CE	2.04	0.69
2:B:304:TYR:CD2	2:B:415:LEU:CD1	2.78	0.67
2:B:445:GLU:OE2	8:B:2107:HOH:O	2.14	0.65
2:B:304:TYR:CG	2:B:415:LEU:HD11	2.31	0.65
1:A:93:GLU:H	1:A:93:GLU:CD	2.02	0.64
2:B:495:TYR:CE1	6:B:1546:P6G:H21	2.37	0.60
1:A:125:LYS:H	1:A:128:MET:HE3	1.67	0.59
2:B:523:ASN:HD22	5:B:1503:NAG:H83	1.70	0.56
1:A:125:LYS:N	1:A:128:MET:CE	2.70	0.55
3:E:1:NAG:H62	3:E:2:NAG:C1	2.38	0.54
1:A:85:TYR:O	8:A:2002:HOH:O	2.19	0.53
1:A:124:LYS:N	1:A:128:MET:HE3	2.22	0.52
2:B:252:LEU:O	2:B:278:PRO:HD3	2.11	0.50
1:A:124:LYS:N	1:A:128:MET:CE	2.68	0.50
1:A:96:MET:HG3	1:A:161:TYR:HB3	1.95	0.48
1:A:48:GLU:H	1:A:48:GLU:CD	2.17	0.48
1:A:68:ASN:HD22	3:C:1:NAG:C7	2.26	0.48
1:A:46:PRO:HG3	1:A:88:GLN:OE1	2.15	0.46
2:B:501:LYS:HG2	2:B:542:LYS:HB3	1.97	0.46
2:B:293:GLY:HA2	6:B:1546:P6G:H111	1.98	0.45
2:B:241:ALA:CB	2:B:451:MET:HG3	2.46	0.45
2:B:271:ARG:HB3	2:B:291:SER:HB3	1.99	0.44
1:A:48:GLU:OE1	1:A:48:GLU:N	2.43	0.44
1:A:202:LEU:O	1:A:205:SER:HB2	2.18	0.43
1:A:101:PHE:HA	1:A:168:GLY:O	2.19	0.43
2:B:412:TYR:N	2:B:413:PRO:CD	2.82	0.43
1:A:47:THR:HB	1:A:48:GLU:OE1	2.19	0.42
2:B:488:ASP:C	2:B:488:ASP:OD1	2.58	0.42
2:B:401:PRO:HG2	2:B:407:TYR:HA	2.01	0.42
2:B:415:LEU:HD22	2:B:423:TYR:CE2	2.54	0.42
1:A:125:LYS:N	1:A:128:MET:HE2	2.33	0.42
1:A:153:ASP:OD2	1:A:155:PHE:HB3	2.20	0.42
2:B:465:ASP:HB3	2:B:468:ASN:HB2	2.03	0.41
2:B:348:TYR:HD1	2:B:348:TYR:HA	1.81	0.41
2:B:456:TYR:CG	2:B:466:PRO:HA	2.55	0.40
2:B:359:LYS:HE2	8:B:2044:HOH:O	2.21	0.40
1:A:200:LEU:HA	1:A:200:LEU:HD23	1.76	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/170 (99%)	166 (99%)	2 (1%)	0	100	100
2	B	321/321 (100%)	303 (94%)	18 (6%)	0	100	100
All	All	489/491 (100%)	469 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/144 (100%)	140 (97%)	4 (3%)	43	36
2	B	289/287 (101%)	281 (97%)	8 (3%)	43	36
All	All	433/431 (100%)	421 (97%)	12 (3%)	43	36

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	SER
1	A	82	LYS
1	A	124	LYS
1	A	179	LEU
2	B	240	PHE
2	B	247	THR
2	B	298	GLN
2	B	306	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	348	TYR
2	B	415	LEU
2	B	480	SER
2	B	487	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	OCS	B	225	2	7,8,9	2.49	2 (28%)	6,11,13	3.44	4 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OCS	B	225	2	-	4/4/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	225	OCS	OD2-SG	5.68	1.67	1.47
2	B	225	OCS	CB-SG	-2.67	1.67	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	225	OCS	OD2-SG-CB	4.78	113.35	105.74
2	B	225	OCS	OD2-SG-OD3	4.40	122.03	111.27
2	B	225	OCS	OD2-SG-OD1	-4.30	100.78	111.27
2	B	225	OCS	OD3-SG-OD1	-2.93	103.80	113.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	225	OCS	N-CA-CB-SG
2	B	225	OCS	CA-CB-SG-OD1
2	B	225	OCS	CA-CB-SG-OD2
2	B	225	OCS	CA-CB-SG-OD3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	225	OCS	1	0

## 5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	C	1	1,3	14,14,15	1.28	1 (7%)	17,19,21	1.78	5 (29%)
3	NAG	C	2	3	14,14,15	0.75	1 (7%)	17,19,21	2.16	6 (35%)
3	NAG	D	1	3,2	14,14,15	0.67	0	17,19,21	1.97	3 (17%)
3	NAG	D	2	3	14,14,15	0.78	1 (7%)	17,19,21	1.83	4 (23%)
3	NAG	E	1	3,2	14,14,15	0.75	0	17,19,21	2.08	7 (41%)
3	NAG	E	2	3	14,14,15	0.66	0	17,19,21	1.70	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	NAG	D	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	NAG	E	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	3/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	NAG	O5-C1	-3.57	1.38	1.43
3	D	2	NAG	C1-C2	2.16	1.55	1.52
3	C	2	NAG	C1-C2	2.13	1.55	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-C2-N2	-5.59	100.94	110.49
3	C	2	NAG	C3-C4-C5	-5.01	101.31	110.24
3	D	2	NAG	C4-C3-C2	4.35	117.40	111.02
3	E	1	NAG	C6-C5-C4	-4.08	103.46	113.00
3	C	1	NAG	O7-C7-C8	-3.93	114.77	122.06
3	C	2	NAG	C2-N2-C7	3.85	128.38	122.90
3	D	2	NAG	O5-C5-C6	3.40	112.53	107.20
3	E	2	NAG	O5-C5-C6	3.25	112.31	107.20
3	C	2	NAG	C1-C2-N2	3.12	115.81	110.49
3	E	1	NAG	O4-C4-C3	-3.12	103.14	110.35
3	D	1	NAG	O3-C3-C4	-3.03	103.35	110.35
3	C	2	NAG	O5-C5-C6	2.99	111.90	107.20
3	C	1	NAG	O3-C3-C4	-2.99	103.45	110.35
3	E	1	NAG	O7-C7-C8	-2.94	116.59	122.06
3	E	1	NAG	O6-C6-C5	-2.81	101.64	111.29
3	E	1	NAG	C8-C7-N2	2.81	120.86	116.10
3	E	2	NAG	O5-C1-C2	2.77	115.66	111.29
3	C	1	NAG	O7-C7-N2	2.55	126.63	121.95
3	E	2	NAG	C2-N2-C7	2.44	126.38	122.90
3	C	1	NAG	O4-C4-C5	2.31	115.03	109.30
3	E	2	NAG	C4-C3-C2	2.29	114.37	111.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	NAG	C1-C2-N2	-2.27	106.62	110.49
3	D	2	NAG	O5-C5-C4	-2.26	105.33	110.83
3	C	2	NAG	O4-C4-C5	2.20	114.75	109.30
3	C	1	NAG	O5-C5-C6	-2.18	103.79	107.20
3	E	1	NAG	C1-O5-C5	2.16	115.12	112.19
3	D	1	NAG	O4-C4-C5	2.16	114.66	109.30
3	C	2	NAG	O3-C3-C2	2.12	113.85	109.47
3	E	1	NAG	O5-C5-C4	2.08	115.89	110.83
3	D	2	NAG	O3-C3-C4	-2.06	105.58	110.35

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
3	E	2	NAG	C1-C2-N2-C7
3	C	1	NAG	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0
3	E	1	NAG	1	0
3	E	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	B	1388	2	14,14,15	0.97	1 (7%)	17,19,21	2.39	4 (23%)
4	P4G	A	1206	-	10,10,10	0.78	0	9,9,9	0.94	0
6	P6G	B	1546	-	18,18,18	0.71	0	17,17,17	0.87	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	B	1503	2	14,14,15	0.69	0	17,19,21	2.13	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	1388	2	-	0/6/23/26	0/1/1/1
4	P4G	A	1206	-	-	3/8/8/8	-
6	P6G	B	1546	-	-	8/16/16/16	-
5	NAG	B	1503	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1388	NAG	C1-C2	2.94	1.56	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1388	NAG	C3-C4-C5	-6.51	98.62	110.24
5	B	1388	NAG	O5-C5-C6	4.69	114.55	107.20
5	B	1503	NAG	C1-O5-C5	4.41	118.17	112.19
5	B	1503	NAG	C1-C2-N2	3.97	117.28	110.49
5	B	1503	NAG	C8-C7-N2	3.20	121.53	116.10
5	B	1503	NAG	O5-C1-C2	-3.19	106.25	111.29
5	B	1503	NAG	C4-C3-C2	-3.06	106.53	111.02
5	B	1388	NAG	C2-N2-C7	2.54	126.53	122.90
5	B	1388	NAG	O6-C6-C5	-2.16	103.87	111.29
5	B	1503	NAG	O7-C7-C8	-2.06	118.24	122.06

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	1546	P6G	O4-C5-C6-O7
6	B	1546	P6G	O10-C11-C12-O13
5	B	1503	NAG	C8-C7-N2-C2
5	B	1503	NAG	O7-C7-N2-C2

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
6	B	1546	P6G	O13-C14-C15-O16
6	B	1546	P6G	O16-C17-C18-O19
4	A	1206	P4G	O3-C5-C6-O4
6	B	1546	P6G	O7-C8-C9-O10
4	A	1206	P4G	O2-C3-C4-O3
6	B	1546	P6G	C12-C11-O10-C9
6	B	1546	P6G	C15-C14-O13-C12
4	A	1206	P4G	C1-C2-O2-C3
6	B	1546	P6G	C11-C12-O13-C14

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1546	P6G	3	0
5	B	1503	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	170/170 (100%)	0.22	16 (9%) <b>8</b> <b>9</b>	24, 38, 63, 81	0
2	B	320/321 (99%)	-0.26	2 (0%) <b>89</b> <b>90</b>	18, 27, 44, 61	0
All	All	490/491 (99%)	-0.09	18 (3%) <b>41</b> <b>44</b>	18, 31, 53, 81	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	LEU	4.2
1	A	47	THR	4.1
1	A	149	TYR	3.3
1	A	56	VAL	3.2
1	A	89	SER	2.9
1	A	46	PRO	2.8
1	A	53	VAL	2.7
1	A	181	GLY	2.7
2	B	479	HIS	2.6
1	A	151	LYS	2.6
1	A	50	THR	2.6
1	A	183	LYS	2.5
1	A	48	GLU	2.4
1	A	86	GLY	2.2
2	B	478	SER	2.2
1	A	88	GLN	2.1
1	A	152	SER	2.0
1	A	45	MET	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OCS	B	225	9/10	0.97	0.09	22,27,46,50	0

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	E	2	14/15	0.80	0.45	64,100,111,115	0
3	NAG	D	2	14/15	0.85	0.43	64,78,81,94	0
3	NAG	C	2	14/15	0.85	0.30	71,83,96,99	0
3	NAG	D	1	14/15	0.93	0.19	44,52,59,65	0
3	NAG	E	1	14/15	0.94	0.10	39,47,57,75	0
3	NAG	C	1	14/15	0.96	0.11	42,51,57,65	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain D:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	B	1503	14/15	0.64	0.40	75,86,102,104	0
4	P4G	A	1206	11/11	0.75	0.32	57,65,79,90	0
5	NAG	B	1388	14/15	0.88	0.17	43,52,60,72	0
6	P6G	B	1546	19/19	0.89	0.13	45,53,72,75	0
7	CL	B	1547	1/1	1.00	0.07	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.