



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 03:28 am BST

PDB ID : 1BXC  
Title : XYLOSE ISOMERASE FROM THERMUS CALDOPHILUS  
Authors : Chang, C.; Park, B.C.; Lee, D.-S.; Suh, S.W.  
Deposited on : 1998-10-02  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	<div> <div>%</div> <div> <div></div> <div>72%</div> <div>25%</div> <div>.</div> </div> </div>
1	B	387	<div> <div>%</div> <div> <div></div> <div>69%</div> <div>27%</div> <div>.</div> </div> </div>
1	C	387	<div> <div></div> <div> <div>71%</div> <div>26%</div> <div>.</div> </div> </div>
1	D	387	<div> <div>%</div> <div> <div></div> <div>70%</div> <div>27%</div> <div>.</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called XYLOSE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			3100	1973	550	567	10			
1	B	387	Total	C	N	O	S	0	0	0
			3100	1973	550	567	10			
1	C	387	Total	C	N	O	S	0	0	0
			3100	1973	550	567	10			
1	D	387	Total	C	N	O	S	0	0	0
			3100	1973	550	567	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	GLN	ALA	CONFLICT	UNP P56681
B	65	GLN	ALA	CONFLICT	UNP P56681
C	65	GLN	ALA	CONFLICT	UNP P56681
D	65	GLN	ALA	CONFLICT	UNP P56681


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	203	Total	O	0	0
			203	203		
2	B	204	Total	O	0	0
			204	204		
2	C	201	Total	O	0	0
			201	201		
2	D	204	Total	O	0	0
			204	204		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: XYLOSE ISOMERASE




#### • Molecule 1: XYLOSE ISOMERASE




#### • Molecule 1: XYLOSE ISOMERASE





• Molecule 1: XYLOSE ISOMERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.35Å 123.60Å 140.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.30 56.55 – 1.72	Depositor EDS
% Data completeness (in resolution range)	76.3 (8.00-2.30) 44.3 (56.55-1.72)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.20 (at 1.72Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.178 , 0.264 0.175 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtriage
Anisotropy	0.588	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 81.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13212	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/3181	0.55	0/4306
1	B	0.45	0/3181	0.56	0/4306
1	C	0.42	0/3181	0.55	0/4306
1	D	0.44	0/3181	0.55	0/4306
All	All	0.44	0/12724	0.56	0/17224

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	C	0	3
1	D	0	2
All	All	0	10

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	PHE	Sidechain
1	A	258	ARG	Sidechain
1	B	115	PHE	Sidechain
1	B	191	TYR	Sidechain
1	B	258	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3100	0	3009	107	0
1	B	3100	0	3009	114	2
1	C	3100	0	3009	114	0
1	D	3100	0	3009	114	0
2	A	203	0	0	13	2
2	B	204	0	0	11	0
2	C	201	0	0	13	0
2	D	204	0	0	15	0
All	All	13212	0	12036	386	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 386 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ARG:HH22	1:B:147:THR:HG21	1.23	1.02
1:D:106:ARG:HH22	1:D:147:THR:HG21	1.25	1.00
1:A:106:ARG:HH22	1:A:147:THR:HG21	1.25	0.99
1:C:106:ARG:HH22	1:C:147:THR:HG21	1.23	0.98
1:C:290:LEU:HD22	1:D:143:GLU:HG3	1.54	0.90

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:ARG:N	2:A:568:HOH:O[1_455]	2.11	0.09
1:B:363:ALA:O	2:A:568:HOH:O[1_455]	2.11	0.09



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/387 (100%)	369 (96%)	14 (4%)	2 (0%)	29	35
1	B	385/387 (100%)	368 (96%)	14 (4%)	3 (1%)	19	23
1	C	385/387 (100%)	368 (96%)	15 (4%)	2 (0%)	29	35
1	D	385/387 (100%)	370 (96%)	13 (3%)	2 (0%)	29	35
All	All	1540/1548 (100%)	1475 (96%)	56 (4%)	9 (1%)	25	31

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	344	ASP
1	B	344	ASP
1	C	344	ASP
1	D	344	ASP
1	A	185	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/313 (100%)	288 (92%)	24 (8%)	13	16
1	B	312/313 (100%)	288 (92%)	24 (8%)	13	16
1	C	312/313 (100%)	288 (92%)	24 (8%)	13	16
1	D	312/313 (100%)	287 (92%)	25 (8%)	12	15
All	All	1248/1252 (100%)	1151 (92%)	97 (8%)	12	16

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	343	MET
1	C	115	PHE
1	D	315	ARG
1	B	350	LYS
1	C	38	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	376	GLN
1	C	219	HIS
1	D	246	ASN
1	C	69	GLN
1	C	229	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/387 (100%)	-0.74	2 (0%) 91 94	6, 20, 50, 69	0
1	B	387/387 (100%)	-0.68	4 (1%) 82 86	6, 19, 51, 71	0
1	C	387/387 (100%)	-0.70	1 (0%) 94 96	6, 21, 51, 69	0
1	D	387/387 (100%)	-0.74	2 (0%) 91 94	5, 20, 51, 63	0
All	All	1548/1548 (100%)	-0.71	9 (0%) 89 92	5, 20, 51, 71	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	344	ASP	4.3
1	C	344	ASP	3.0
1	B	76	ARG	3.0
1	A	344	ASP	2.9
1	B	1	MET	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.