



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 10:03 pm BST

PDB ID : 1BXN
Title : THE CRYSTAL STRUCTURE OF RUBISCO FROM ALCALIGENES EUTROPHUS TO 2.7 ANGSTROMS.
Authors : Hansen, S.; Vollan, V.B.; Hough, E.; Andersen, K.
Deposited on : 1998-10-06
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	<div> <div>8%</div> <div> <div></div> <div>65%</div> <div>24%</div> <div>• 8%</div> </div> </div>
1	C	485	<div> <div>9%</div> <div> <div></div> <div>65%</div> <div>24%</div> <div>• 8%</div> </div> </div>
1	E	485	<div> <div>10%</div> <div> <div></div> <div>65%</div> <div>24%</div> <div>• 8%</div> </div> </div>
1	G	485	<div> <div>8%</div> <div> <div></div> <div>66%</div> <div>23%</div> <div>• 8%</div> </div> </div>
2	I	139	<div> <div>10%</div> <div> <div></div> <div>55%</div> <div>33%</div> <div>5% 7%</div> </div> </div>
2	J	139	<div> <div>12%</div> <div> <div></div> <div>53%</div> <div>35%</div> <div>5% 7%</div> </div> </div>
2	K	139	<div> <div>10%</div> <div> <div></div> <div>53%</div> <div>35%</div> <div>• 7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	139	<div><div></div><div>12%</div><div>50%</div><div>37%</div><div>5%</div><div>7%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	0	0
			3470	2201	612	636	21			
1	C	445	Total	C	N	O	S	0	0	0
			3470	2201	612	636	21			
1	E	445	Total	C	N	O	S	0	0	0
			3470	2201	612	636	21			
1	G	445	Total	C	N	O	S	0	0	0
			3470	2201	612	636	21			

- Molecule 2 is a protein called PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	129	Total	C	N	O	S	0	0	0
			1056	670	180	200	6			
2	J	129	Total	C	N	O	S	0	0	0
			1056	670	180	200	6			
2	K	129	Total	C	N	O	S	0	0	0
			1056	670	180	200	6			
2	L	129	Total	C	N	O	S	0	0	0
			1056	670	180	200	6			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	E	1	Total	O	P	0	0
			5	4	1		
3	E	1	Total	O	P	0	0
			5	4	1		
3	G	1	Total	O	P	0	0
			5	4	1		
3	G	1	Total	O	P	0	0
			5	4	1		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN)




• Molecule 1: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN)






• Molecule 1: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN)




• Molecule 1: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN)




• Molecule 2: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN)







● Molecule 2: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN)



● Molecule 2: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN)



● Molecule 2: PROTEIN (RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	112.00Å 112.00Å 402.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 24.60 – 2.65	Depositor EDS
% Data completeness (in resolution range)	90.0 (15.00-2.70) 86.9 (24.60-2.65)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.64Å)	Xtriage
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.266 , 0.322 0.280 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	45.5	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	18144	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/3552	0.57	0/4816
1	C	0.33	0/3552	0.57	0/4816
1	E	0.33	0/3552	0.57	0/4816
1	G	0.34	0/3552	0.57	0/4816
2	I	0.37	0/1084	0.61	0/1474
2	J	0.36	0/1084	0.61	0/1474
2	K	0.36	0/1084	0.61	0/1474
2	L	0.36	0/1084	0.61	0/1474
All	All	0.34	0/18544	0.58	0/25160

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3470	0	3402	89	0
1	C	3470	0	3402	88	0
1	E	3470	0	3402	86	0
1	G	3470	0	3402	84	0
2	I	1056	0	1007	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	J	1056	0	1007	39	0
2	K	1056	0	1007	39	0
2	L	1056	0	1007	42	0
3	A	10	0	0	0	0
3	C	10	0	0	0	0
3	E	10	0	0	0	0
3	G	10	0	0	1	0
All	All	18144	0	17636	467	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 467 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:THR:HG23	2:I:10:PHE:HE2	1.48	0.79
1:G:433:ARG:HH21	1:G:434:ASN:HD21	1.31	0.78
1:C:420:THR:HG23	2:J:10:PHE:HE2	1.50	0.77
1:C:433:ARG:HH21	1:C:434:ASN:HD21	1.32	0.77
1:E:433:ARG:HH21	1:E:434:ASN:HD21	1.29	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/485 (91%)	390 (88%)	41 (9%)	12 (3%)	5	12
1	C	443/485 (91%)	389 (88%)	43 (10%)	11 (2%)	5	14
1	E	443/485 (91%)	390 (88%)	42 (10%)	11 (2%)	5	14
1	G	443/485 (91%)	389 (88%)	44 (10%)	10 (2%)	6	16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	127/139 (91%)	104 (82%)	18 (14%)	5 (4%)	3	6
2	J	127/139 (91%)	104 (82%)	18 (14%)	5 (4%)	3	6
2	K	127/139 (91%)	104 (82%)	19 (15%)	4 (3%)	4	9
2	L	127/139 (91%)	104 (82%)	18 (14%)	5 (4%)	3	6
All	All	2280/2496 (91%)	1974 (87%)	243 (11%)	63 (3%)	5	11

5 of 63 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	207	GLU
2	I	103	PRO
2	I	105	ASP
1	C	207	GLU
2	J	103	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/391 (92%)	343 (96%)	15 (4%)	30	58
1	C	358/391 (92%)	343 (96%)	15 (4%)	30	58
1	E	358/391 (92%)	343 (96%)	15 (4%)	30	58
1	G	358/391 (92%)	343 (96%)	15 (4%)	30	58
2	I	115/122 (94%)	105 (91%)	10 (9%)	10	23
2	J	115/122 (94%)	105 (91%)	10 (9%)	10	23
2	K	115/122 (94%)	105 (91%)	10 (9%)	10	23
2	L	115/122 (94%)	105 (91%)	10 (9%)	10	23
All	All	1892/2052 (92%)	1792 (95%)	100 (5%)	22	48

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	96	MET
1	E	119	LEU
2	L	49	GLU
2	J	99	ILE
1	E	25	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	45	ASN
1	E	244	ASN
2	L	45	ASN
2	J	70	ASN
1	E	210	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	A	502	-	4,4,4	1.58	1 (25%)	6,6,6	1.07	0
3	PO4	C	504	-	4,4,4	1.65	1 (25%)	6,6,6	1.04	0
3	PO4	G	507	-	4,4,4	1.77	1 (25%)	6,6,6	1.12	0
3	PO4	A	501	-	4,4,4	1.71	1 (25%)	6,6,6	1.10	0
3	PO4	G	508	-	4,4,4	1.56	1 (25%)	6,6,6	1.13	0
3	PO4	C	503	-	4,4,4	1.73	1 (25%)	6,6,6	1.08	0
3	PO4	E	506	-	4,4,4	1.45	1 (25%)	6,6,6	1.02	0
3	PO4	E	505	-	4,4,4	1.58	1 (25%)	6,6,6	1.11	0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	PO4	P-O3	-3.17	1.45	1.54
3	C	503	PO4	P-O3	-3.17	1.45	1.54
3	G	507	PO4	P-O3	-3.13	1.45	1.54
3	A	502	PO4	P-O3	-2.93	1.45	1.54
3	C	504	PO4	P-O3	-2.92	1.45	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	507	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/485 (91%)	0.20	38 (8%) 10 9	2, 29, 99, 100	0
1	C	445/485 (91%)	0.20	43 (9%) 7 6	2, 28, 98, 100	0
1	E	445/485 (91%)	0.38	49 (11%) 5 4	2, 31, 100, 100	0
1	G	445/485 (91%)	0.18	38 (8%) 10 9	2, 26, 98, 100	0
2	I	129/139 (92%)	0.46	14 (10%) 5 4	4, 35, 99, 100	0
2	J	129/139 (92%)	0.55	17 (13%) 3 2	2, 40, 100, 100	0
2	K	129/139 (92%)	0.59	14 (10%) 5 4	3, 41, 98, 100	0
2	L	129/139 (92%)	0.47	16 (12%) 4 3	2, 36, 100, 100	0
All	All	2296/2496 (91%)	0.30	229 (9%) 7 5	2, 31, 100, 100	0

The worst 5 of 229 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	72	VAL	12.4
1	G	72	VAL	11.9
1	E	337	LEU	10.1
1	C	72	VAL	9.8
1	A	74	THR	9.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PO4	G	508	5/5	0.92	0.24	18,23,40,42	0
3	PO4	C	504	5/5	0.94	0.25	18,23,40,42	0
3	PO4	G	507	5/5	0.94	0.18	18,23,40,42	0
3	PO4	E	506	5/5	0.94	0.33	18,23,40,42	0
3	PO4	A	501	5/5	0.95	0.22	18,23,40,42	0
3	PO4	C	503	5/5	0.95	0.22	18,23,40,42	0
3	PO4	A	502	5/5	0.96	0.25	18,23,40,42	0
3	PO4	E	505	5/5	0.97	0.20	18,23,40,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.