



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 08:22 am BST

PDB ID : 3BZN  
Title : Crystal Structure of Open form of Menaquinone-Specific Isochorismate Synthase, MenF  
Authors : Parsons, J.F.; Shi, K.M.; Ladner, J.E.  
Deposited on : 2008-01-18  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	504	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Menaquinone-specific isochorismate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	11	0
			3482	2197	634	637	14			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	220	Total	O		0	0
			220	220			

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Menaquinone-specific isochorismate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.28Å 70.49Å 90.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 29.62 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.00) 98.7 (29.62-1.99)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.04 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.202 , 0.284 0.204 , 0.287	Depositor DCC
$R_{free}$ test set	1463 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.0	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	2/3583 (0.1%)	1.06	19/4861 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	381	GLU	CD-OE2	6.71	1.33	1.25
1	A	90	ALA	CA-CB	-5.29	1.41	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	283	ARG	NE-CZ-NH2	-11.53	114.54	120.30
1	A	283	ARG	NE-CZ-NH1	8.76	124.68	120.30
1	A	72	ARG	NE-CZ-NH2	-8.14	116.23	120.30
1	A	359	ARG	NE-CZ-NH1	7.36	123.98	120.30
1	A	359	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	A	72	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	A	11	LEU	CA-CB-CG	6.37	129.96	115.30
1	A	259	VAL	CB-CA-C	-6.26	99.50	111.40
1	A	255	LEU	CA-CB-CG	6.06	129.24	115.30
1	A	295	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	A	81	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	A	168	ASP	CB-CG-OD1	5.88	123.59	118.30
1	A	241	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	235	LEU	CB-CG-CD2	-5.73	101.27	111.00
1	A	241	ARG	NE-CZ-NH1	5.71	123.15	120.30
1	A	386	LEU	CB-CG-CD1	5.29	120.00	111.00
1	A	306	LEU	CA-CB-CG	-5.24	103.24	115.30
1	A	81	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	A	336	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3482	0	3487	62	0
2	A	1	0	0	0	0
3	A	15	0	0	3	0
4	A	220	0	0	8	0
All	All	3718	0	3487	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:ILE:HD11	1:A:137:ALA:HB2	1.55	0.86
1:A:15:LEU:O	1:A:19:ILE:HG12	1.83	0.79
1:A:143:PHE:O	1:A:146:THR:HB	1.89	0.72
3:A:504:SO4:O2	4:A:603:HOH:O	2.13	0.67
1:A:66:VAL:HB	1:A:110:PRO:HB3	1.76	0.66
1:A:19:ILE:HD11	1:A:137:ALA:CB	2.27	0.65
1:A:19:ILE:HG13	1:A:134:GLN:HA	1.83	0.60
1:A:70:ILE:HG21	1:A:128:PHE:HB3	1.84	0.60
1:A:98:ASN:HB2	1:A:371[B]:SER:OG	2.02	0.59
1:A:25:ILE:HA	1:A:127:LEU:O	2.01	0.59
1:A:126:THR:HG23	4:A:549:HOH:O	2.01	0.59
1:A:72:ARG:NH2	4:A:636:HOH:O	2.35	0.58
1:A:201:PHE:HE1	1:A:396:VAL:HG23	1.68	0.58
1:A:50:TYR:HB3	1:A:51:PRO:HA	1.87	0.57
1:A:141:LYS:HE2	4:A:678:HOH:O	2.03	0.57
1:A:81:ARG:HG3	4:A:641:HOH:O	2.04	0.57
1:A:57[A]:GLN:OE1	1:A:57[A]:GLN:HA	2.05	0.56
1:A:162:ARG:HB3	1:A:200:HIS:HB2	1.88	0.56
1:A:242:LEU:HD12	1:A:371[A]:SER:OG	2.06	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:SER:O	1:A:135:HIS:HB3	2.05	0.56
1:A:259:VAL:HG22	1:A:275:LEU:HD22	1.87	0.56
1:A:340:GLN:OE1	1:A:371[A]:SER:HB2	2.07	0.55
1:A:65:VAL:HG22	1:A:114:TRP:HB3	1.88	0.54
1:A:283:ARG:NH2	1:A:405[B]:VAL:HG13	2.22	0.54
1:A:57[A]:GLN:HB2	1:A:62:GLU:HB3	1.91	0.53
1:A:308:PRO:HA	1:A:321:ARG:HG2	1.90	0.53
1:A:8:LEU:CD2	1:A:30:ILE:HD11	2.39	0.53
1:A:242:LEU:HD11	1:A:340:GLN:HB2	1.93	0.51
1:A:49:THR:OG1	1:A:146:THR:HG21	2.10	0.51
1:A:23:PRO:HD3	1:A:131:SER:O	2.12	0.50
1:A:200:HIS:CD2	1:A:395:VAL:HG22	2.47	0.50
1:A:378:GLN:HG2	1:A:379:GLN:HG2	1.94	0.49
1:A:386:LEU:HG	3:A:504:SO4:O2	2.14	0.48
1:A:39:ASP:OD2	1:A:42:SER:OG	2.25	0.48
1:A:43:TRP:HD1	1:A:150:ILE:HG13	1.79	0.48
1:A:80:GLN:O	1:A:84:ARG:HG3	2.13	0.47
1:A:420:LYS:HD3	3:A:504:SO4:O3	2.14	0.46
1:A:19:ILE:CD1	1:A:137:ALA:HB2	2.36	0.46
1:A:304[B]:ASP:OD1	4:A:689:HOH:O	2.21	0.46
1:A:150:ILE:O	1:A:151:LYS:HE3	2.15	0.46
1:A:14:HIS:ND1	1:A:14:HIS:N	2.64	0.45
1:A:162:ARG:CB	1:A:200:HIS:HB2	2.46	0.45
1:A:258:THR:HA	1:A:317:GLN:O	2.17	0.45
1:A:348:LEU:HA	1:A:349:PRO:C	2.37	0.45
1:A:345:VAL:HG22	1:A:386:LEU:HD11	1.99	0.44
1:A:199:LEU:O	1:A:395:VAL:HA	2.18	0.44
1:A:83:LEU:HB3	1:A:377:LEU:HD13	1.99	0.44
1:A:8:LEU:HD21	1:A:30:ILE:HD11	1.99	0.44
1:A:8:LEU:HD23	1:A:8:LEU:HA	1.92	0.43
1:A:43:TRP:HD1	1:A:150:ILE:CG1	2.31	0.43
1:A:163:GLU:HG2	1:A:199:LEU:HD12	2.01	0.43
1:A:165:HIS:HB3	1:A:167:PRO:O	2.18	0.43
1:A:346:ALA:HB3	4:A:507:HOH:O	2.18	0.42
1:A:54:TYR:HB3	1:A:225:TYR:HB3	2.00	0.42
1:A:201:PHE:CE1	1:A:396:VAL:HG23	2.51	0.42
1:A:126:THR:HG21	4:A:548:HOH:O	2.18	0.42
1:A:240:GLU:O	1:A:383:CYS:HB2	2.20	0.42
1:A:331:ASP:HB3	1:A:334:ILE:HD12	2.01	0.42
1:A:54:TYR:HD1	1:A:65:VAL:HG12	1.85	0.41
1:A:345:VAL:HG21	1:A:384:VAL:HG11	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57[A]:GLN:OE1	1:A:57[A]:GLN:CA	2.67	0.41
1:A:409:ASP:OD2	1:A:409:ASP:C	2.59	0.40
1:A:95:TRP:CE3	1:A:374:TYR:HB3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	437/431 (101%)	420 (96%)	15 (3%)	2 (0%)	29 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247[A]	ASP
1	A	247[B]	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	372/363 (102%)	324 (87%)	48 (13%)	4 2

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	11	LEU
1	A	14	HIS
1	A	18	GLU
1	A	41	LEU
1	A	44	LEU
1	A	104	GLN
1	A	108	LEU
1	A	109	LEU
1	A	115	ARG
1	A	126	THR
1	A	129	SER
1	A	138	ILE
1	A	146	THR
1	A	148	VAL
1	A	151	LYS
1	A	158	LEU
1	A	161	THR
1	A	181	LYS
1	A	199	LEU
1	A	241	ARG
1	A	242	LEU
1	A	250	LEU
1	A	255	LEU
1	A	259	VAL
1	A	269	GLN
1	A	287	LEU
1	A	297	GLN
1	A	302	THR
1	A	303	LEU
1	A	314	ARG
1	A	316	VAL
1	A	326	SER
1	A	336	LEU
1	A	345	VAL
1	A	359	ARG
1	A	371[A]	SER
1	A	371[B]	SER
1	A	377	LEU
1	A	378	GLN
1	A	386	LEU
1	A	390[A]	LYS
1	A	390[B]	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	411	GLU
1	A	424	LEU
1	A	425	ARG
1	A	430[A]	MET
1	A	430[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	502	2	4,4,4	0.20	0	6,6,6	0.58	0
3	SO4	A	504	-	4,4,4	0.38	0	6,6,6	0.79	0
3	SO4	A	503	-	4,4,4	0.34	0	6,6,6	0.84	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	504	SO4	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	430/431 (99%)	-0.12	8 (1%) 66 65	15, 27, 46, 61	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	ALA	3.9
1	A	14	HIS	3.1
1	A	430[A]	MET	2.9
1	A	18	GLU	2.6
1	A	19	ILE	2.6
1	A	20	PRO	2.5
1	A	135	HIS	2.5
1	A	34	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	504	5/5	0.96	0.19	10,19,23,25	5
2	MG	A	501	1/1	0.96	0.05	25,25,25,25	0
3	SO4	A	503	5/5	0.97	0.07	26,26,34,36	0
3	SO4	A	502	5/5	0.99	0.07	18,22,25,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.