



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 06:26 pm BST

PDB ID : 2C0R
Title : CRYSTAL STRUCTURE OF PHOSPHOSERINE AMINOTRANSFERASE
FROM BACILLUS CIRCULANS VAR. ALKALOPHILUS AT pH 8.5
Authors : Kapetanidou, E.G.; Papageorgiou, A.C.
Deposited on : 2005-09-07
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>3%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	B	362	<div> <div>3%</div> <div>85%</div> <div>13%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHOSERINE AMINOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	1	0
			2784	1759	475	536	14			
1	B	361	Total	C	N	O	S	0	3	0
			2790	1763	475	538	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	GLU	LYS	engineered mutation	UNP Q59196
B	3	GLU	LYS	engineered mutation	UNP Q59196

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	397	Total	O	0	0
			397	397		
3	B	412	Total	O	0	0
			412	412		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHOSPHOSERINE AMINOTRANSFERASE



• Molecule 1: PHOSPHOSERINE AMINOTRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.78 Å 90.54 Å 157.82 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 1.20 14.99 – 1.20	Depositor EDS
% Data completeness (in resolution range)	85.6 (12.00-1.20) 82.5 (14.99-1.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.20 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.154 , 0.212 0.147 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	13.0	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 60.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	6413	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2845	1.31	26/3849 (0.7%)
1	B	0.61	0/2861	1.39	26/3869 (0.7%)
All	All	0.61	0/5706	1.35	52/7718 (0.7%)

There are no bond length outliers.

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	287	ARG	NE-CZ-NH2	-29.34	105.63	120.30
1	B	287	ARG	NE-CZ-NH1	15.66	128.13	120.30
1	B	258[A]	ARG	CD-NE-CZ	11.15	139.20	123.60
1	B	258[B]	ARG	CD-NE-CZ	11.15	139.20	123.60
1	A	258	ARG	NE-CZ-NH1	10.62	125.61	120.30
1	A	129	MET	CG-SD-CE	-10.55	83.32	100.20
1	A	258	ARG	CD-NE-CZ	10.04	137.65	123.60
1	A	303	ARG	CD-NE-CZ	9.74	137.24	123.60
1	A	43	ARG	NE-CZ-NH1	9.62	125.11	120.30
1	A	258	ARG	NE-CZ-NH2	-9.32	115.64	120.30
1	B	295	ARG	NE-CZ-NH2	-9.22	115.69	120.30
1	A	226	ARG	NE-CZ-NH2	-9.03	115.78	120.30
1	A	43	ARG	NE-CZ-NH2	-8.60	116.00	120.30
1	A	277	ASP	CB-CG-OD1	7.82	125.34	118.30
1	A	303	ARG	NE-CZ-NH1	7.75	124.17	120.30
1	B	226	ARG	NE-CZ-NH1	-7.71	116.44	120.30
1	B	57	ARG	NE-CZ-NH1	-7.65	116.48	120.30
1	B	335	ARG	NE-CZ-NH2	-7.63	116.49	120.30
1	A	361	ARG	NE-CZ-NH2	-7.59	116.50	120.30
1	B	258[A]	ARG	NE-CZ-NH2	7.42	124.01	120.30
1	B	258[B]	ARG	NE-CZ-NH2	7.42	124.01	120.30
1	A	22	ARG	NE-CZ-NH2	7.38	123.99	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	287	ARG	CD-NE-CZ	6.55	132.77	123.60
1	B	183	PHE	CB-CG-CD2	-6.39	116.33	120.80
1	A	226	ARG	NE-CZ-NH1	6.33	123.46	120.30
1	B	361	ARG	C-N-CA	6.17	135.26	122.30
1	B	287	ARG	NH1-CZ-NH2	6.17	126.19	119.40
1	B	210	ARG	NE-CZ-NH2	-6.05	117.28	120.30
1	A	47	TYR	CB-CG-CD2	5.99	124.59	121.00
1	A	6	TYR	CB-CG-CD1	-5.95	117.43	121.00
1	A	22	ARG	NE-CZ-NH1	-5.81	117.39	120.30
1	A	291	ASP	CB-CG-OD1	5.76	123.49	118.30
1	B	30	TYR	CB-CG-CD1	-5.73	117.56	121.00
1	A	47	TYR	CA-CB-CG	-5.71	102.56	113.40
1	A	88	PHE	CB-CG-CD2	5.56	124.69	120.80
1	B	356	HIS	CA-CB-CG	-5.56	104.15	113.60
1	A	30	TYR	CB-CG-CD1	-5.45	117.73	121.00
1	B	361	ARG	NE-CZ-NH1	-5.45	117.57	120.30
1	A	57	ARG	CD-NE-CZ	5.45	131.23	123.60
1	B	313	PHE	CB-CG-CD2	5.43	124.60	120.80
1	B	210	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	B	57	ARG	NE-CZ-NH2	5.42	123.01	120.30
1	B	47	TYR	CA-CB-CG	-5.40	103.14	113.40
1	A	269	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	30	TYR	CB-CG-CD2	5.20	124.12	121.00
1	A	291	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	B	97	TYR	CB-CG-CD2	-5.16	117.90	121.00
1	A	277	ASP	CB-CG-OD2	-5.15	113.66	118.30
1	B	227	TYR	CB-CG-CD2	-5.12	117.92	121.00
1	A	155	GLU	OE1-CD-OE2	-5.08	117.20	123.30
1	A	2	SER	CA-C-N	5.03	128.26	117.20
1	B	286	TYR	CB-CG-CD1	-5.02	117.99	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2784	0	2706	39	0
1	B	2790	0	2716	35	0
2	A	15	0	6	1	0
2	B	15	0	6	1	0
3	A	397	0	0	7	0
3	B	412	0	0	7	0
All	All	6413	0	5434	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:ASN:HD21	1:A:234:ASN:HD21	1.19	0.90
1:A:96:ASN:HD21	1:A:140:GLN:H	1.19	0.89
1:B:52:ASN:HD21	1:B:234:ASN:HD21	1.20	0.85
1:A:58:LEU:HD13	1:A:179:LEU:HD11	1.57	0.85
1:A:140:GLN:HG2	1:A:142:ASN:OD1	1.83	0.78
1:A:258:ARG:HG2	1:A:258:ARG:HH11	1.50	0.77
1:A:139:LEU:HD21	1:A:146:LEU:HD13	1.66	0.76
1:A:232:LYS:HE2	3:A:2262:HOH:O	1.84	0.75
1:A:225:LEU:HD11	1:B:225:LEU:HD11	1.69	0.73
1:B:181:ARG:HD2	3:B:2257:HOH:O	1.93	0.69
1:B:151:ASN:HD21	1:B:299:ASN:HD22	1.41	0.69
1:B:352:GLN:HG3	3:B:2407:HOH:O	1.90	0.68
1:B:127:ASN:HB2	3:B:2106:HOH:O	1.93	0.67
1:A:197:LYS:HE3	3:B:2303:HOH:O	1.94	0.67
1:A:210:ARG:HH21	1:A:213:LEU:HD21	1.61	0.66
1:A:307:GLU:HG2	3:A:2346:HOH:O	1.95	0.66
1:A:96:ASN:ND2	1:A:140:GLN:H	1.91	0.65
1:B:52:ASN:ND2	1:B:234:ASN:HD21	1.96	0.62
1:B:210:ARG:NH2	1:B:213:LEU:HD11	2.15	0.62
1:B:210:ARG:HH21	1:B:213:LEU:HD21	1.65	0.61
1:B:127:ASN:HB2	3:B:2202:HOH:O	2.03	0.58
1:A:57:ARG:HG2	1:A:57:ARG:HH11	1.68	0.58
1:A:52:ASN:ND2	1:A:234:ASN:HD21	1.98	0.58
1:B:57:ARG:NH2	1:B:253:LYS:HE2	2.19	0.58
1:B:52:ASN:HD21	1:B:234:ASN:ND2	1.98	0.57
1:A:289:CYS:SG	1:A:303:ARG:HG3	2.44	0.57
1:A:127:ASN:HB3	1:A:129:MET:CE	2.35	0.56
1:A:52:ASN:HD21	1:A:234:ASN:ND2	1.99	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:312:GLU:HG2	1:B:357:PHE:HZ	1.71	0.56
1:A:129:MET:HG2	1:A:289:CYS:O	2.07	0.54
1:A:225:LEU:CD1	1:B:225:LEU:HD11	2.37	0.54
1:B:7:ASN:HA	1:B:323[B]:VAL:HG22	1.88	0.54
1:A:129:MET:HG2	1:A:290:VAL:HA	1.89	0.53
1:A:139:LEU:CD2	1:A:146:LEU:HD13	2.38	0.52
1:B:225:LEU:HD21	3:B:2073:HOH:O	2.10	0.51
1:B:210:ARG:HG2	1:B:212:ASP:OD1	2.12	0.49
1:B:285:PHE:HB2	1:B:358:LYS:NZ	2.27	0.49
1:B:90:LYS:HB2	1:B:93:GLN:OE1	2.12	0.49
1:A:74:GLN:HB3	1:B:74:GLN:HB3	1.96	0.48
1:A:43:ARG:HD2	1:A:43:ARG:HA	1.63	0.48
1:B:312:GLU:HG2	1:B:357:PHE:CZ	2.48	0.48
1:A:258:ARG:HG2	1:A:258:ARG:NH1	2.19	0.47
1:A:197:LYS:HE2	3:A:2019:HOH:O	2.13	0.47
1:A:53:GLU:OE2	1:A:57:ARG:NH2	2.48	0.46
1:A:197:LYS:NZ	2:A:363:PLP:O3	2.49	0.46
1:B:197:LYS:NZ	2:B:363:PLP:O3	2.49	0.46
1:B:151:ASN:HD21	1:B:299:ASN:ND2	2.10	0.45
1:A:287:ARG:NE	3:A:2324:HOH:O	2.49	0.45
1:B:219:LYS:HE2	3:B:2282:HOH:O	2.17	0.45
1:A:112:LYS:NZ	3:A:2166:HOH:O	2.48	0.45
1:B:139:LEU:HD21	1:B:146:LEU:HD13	1.98	0.45
1:A:57:ARG:NH1	3:A:2106:HOH:O	2.49	0.45
1:A:18:GLU:OE2	1:A:258:ARG:NH2	2.50	0.45
1:B:142:ASN:N	1:B:142:ASN:OD1	2.50	0.44
1:B:7:ASN:HA	1:B:323[A]:VAL:HG13	1.98	0.44
1:A:53:GLU:HG3	3:A:2100:HOH:O	2.17	0.44
1:B:94:THR:HG23	1:B:142:ASN:ND2	2.33	0.44
1:B:43:ARG:HD2	1:B:43:ARG:HA	1.78	0.43
1:A:58:LEU:CD1	1:A:179:LEU:HD11	2.40	0.43
1:B:304:LEU:HB3	1:B:309:LEU:HD23	1.99	0.43
1:B:179:LEU:HD12	1:B:199:LEU:CD2	2.48	0.43
1:A:334:LEU:HD22	1:A:334:LEU:N	2.34	0.43
1:B:2:SER:OG	1:B:3:GLU:N	2.50	0.43
1:A:94:THR:HG21	1:A:140:GLN:HE21	1.83	0.42
1:A:127:ASN:HB3	1:A:129:MET:HE3	2.01	0.42
1:A:195:ALA:CB	1:A:204[B]:VAL:HG13	2.50	0.41
1:A:94:THR:HG21	1:A:140:GLN:NE2	2.34	0.41
1:B:135:GLN:O	1:B:135:GLN:HG3	2.20	0.41
1:A:53:GLU:OE2	1:A:57:ARG:NE	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:ARG:HH21	1:B:253:LYS:HE2	1.86	0.41
1:B:285:PHE:HB2	1:B:358:LYS:HZ2	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/362 (99%)	353 (98%)	7 (2%)	0	100	100
1	B	362/362 (100%)	354 (98%)	7 (2%)	1 (0%)	41	16
All	All	722/724 (100%)	707 (98%)	14 (2%)	1 (0%)	51	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	3	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/297 (98%)	287 (98%)	5 (2%)	60	24
1	B	295/297 (99%)	289 (98%)	6 (2%)	55	17
All	All	587/594 (99%)	576 (98%)	11 (2%)	57	19

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	PRO
1	A	197	LYS
1	A	258	ARG
1	A	311	LYS
1	A	329	ARG
1	B	3	GLU
1	B	287	ARG
1	B	308	GLU
1	B	319	GLN
1	B	360	SER
1	B	361	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	64	ASN
1	A	80	GLN
1	A	96	ASN
1	A	140	GLN
1	A	233	ASN
1	B	52	ASN
1	B	64	ASN
1	B	80	GLN
1	B	127	ASN
1	B	233	ASN
1	B	265	GLN
1	B	266	GLN
1	B	299	ASN
1	B	319	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	363	1	15,15,16	1.49	1 (6%)	20,22,23	1.44	2 (10%)
2	PLP	B	363	1	15,15,16	1.20	1 (6%)	20,22,23	1.68	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	363	1	-	0/6/6/8	0/1/1/1
2	PLP	B	363	1	-	0/6/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	363	PLP	C3-C2	-4.06	1.36	1.40
2	B	363	PLP	C5-C4	-3.11	1.37	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	363	PLP	C4A-C4-C5	-4.03	116.78	120.94
2	A	363	PLP	C6-C5-C4	3.91	121.24	118.16
2	A	363	PLP	C5-C6-N1	-3.50	118.00	123.82
2	B	363	PLP	C6-C5-C4	2.86	120.41	118.16
2	B	363	PLP	C4A-C4-C3	2.62	124.95	120.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	363	PLP	C3-C2-N1	-2.58	117.43	120.77
2	B	363	PLP	C5-C6-N1	-2.06	120.38	123.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	363	PLP	1	0
2	B	363	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	361/362 (99%)	0.03	10 (2%) 53 51	11, 16, 31, 64	1 (0%)
1	B	361/362 (99%)	0.04	10 (2%) 53 51	10, 15, 34, 76	0
All	All	722/724 (99%)	0.04	20 (2%) 53 51	10, 15, 33, 76	1 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	362	GLY	12.3
1	B	2	SER	11.6
1	A	129	MET	5.9
1	A	362	GLY	4.3
1	A	329	ARG	3.9
1	B	360	SER	3.6
1	B	308	GLU	3.5
1	A	330	SER	3.2
1	B	3	GLU	3.2
1	B	361	ARG	3.2
1	A	92	GLY	3.0
1	B	356	HIS	2.8
1	B	141	ASP	2.6
1	A	359	ARG	2.6
1	B	135	GLN	2.3
1	A	141	ASP	2.3
1	A	199	LEU	2.2
1	A	200	GLY	2.1
1	A	289	CYS	2.1
1	B	283	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PLP	A	363	15/16	0.98	0.06	11,12,15,19	0
2	PLP	B	363	15/16	0.98	0.07	11,12,15,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.