



Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2020 – 09:32 am BST

PDB ID : 6C0A
Title : Actinin-1 EF-Hand bound to the Cav1.2 IQ Motif
Authors : Turner, M.L.; Ames, J.B.
Deposited on : 2017-12-28

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 61%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	71	
2	B	23	

2 Ensemble composition and analysis ⓘ

This entry contains 4 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:829-A:889, B:1653-B:1668 (77)	0.25	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1488 atoms, of which 732 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Alpha-actinin-1.

Mol	Chain	Residues	Atoms						Trace
1	A	71	Total	C	H	N	O	S	0
			1065	342	517	85	116	5	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	845	MET	GLU	conflict	UNP Q7TPR4
A	868	THR	ALA	conflict	UNP Q7TPR4

- Molecule 2 is a protein called Voltage-dependent L-type calcium channel subunit alpha-1C.


Mol	Chain	Residues	Atoms					Trace
2	B	23	Total	C	H	N	O	0
			423	139	215	36	33	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Alpha-actinin-1



- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1C




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Alpha-actinin-1




- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1C



4.2.2 Score per residue for model 2

- Molecule 1: Alpha-actinin-1




- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1C



4.2.3 Score per residue for model 3

- Molecule 1: Alpha-actinin-1




- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1C



4.2.4 Score per residue for model 4

- Molecule 1: Alpha-actinin-1



- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1C



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 194 calculated structures, 4 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
HADDOCK	structure calculation	
PALES	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	755
Number of shifts mapped to atoms	755
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	61%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	473	456	456	4±1
2	B	149	158	158	1±0
All	All	2488	2456	2456	18

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:867:TYR:CD1	1:A:867:TYR:N	0.58	2.72	3	3
1:A:860:CYS:O	1:A:864:MET:HG3	0.48	2.08	1	3
1:A:845:MET:HG3	1:A:846:ASP:N	0.48	2.23	4	1
2:B:1662:LYS:O	2:B:1666:GLU:HG3	0.47	2.09	1	1
2:B:1655:GLN:O	2:B:1659:ARG:HD3	0.45	2.11	2	1
1:A:849:ARG:HA	1:A:857:ALA:CB	0.43	2.43	4	1
2:B:1660:LYS:O	2:B:1664:ARG:HB2	0.42	2.14	3	1
1:A:867:TYR:N	1:A:867:TYR:CD1	0.42	2.87	4	1
1:A:867:TYR:O	1:A:878:ASP:HB2	0.41	2.16	3	1
1:A:866:PRO:O	1:A:868:THR:N	0.41	2.54	2	1
2:B:1659:ARG:O	2:B:1663:LYS:HG3	0.41	2.16	4	1
1:A:865:ALA:O	1:A:877:LEU:HG	0.41	2.15	4	1
1:A:833:PHE:HB3	1:A:879:TYR:CE1	0.41	2.50	4	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:867:TYR:HD1	1:A:867:TYR:N	0.40	2.13	3	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	61/71 (86%)	56±1 (91±1%)	3±1 (5±2%)	3±0 (5±1%)	4	28
2	B	15/23 (65%)	15±0 (98±3%)	0±0 (2±3%)	0±0 (0±0%)	100	100
All	All	304/376 (81%)	281 (92%)	12 (4%)	11 (4%)	6	34

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	866	PRO	4
1	A	867	TYR	4
1	A	868	THR	3

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	50/59 (85%)	41±0 (82±0%)	9±0 (18±0%)	4	38
2	B	15/20 (75%)	14±1 (93±5%)	1±1 (7±5%)	20	68
All	All	260/316 (82%)	220 (85%)	40 (15%)	6	43

All 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	844	THR	4
1	A	850	ARG	4
1	A	842	TYR	4
1	A	867	TYR	4
1	A	833	PHE	4
1	A	847	GLU	3
2	B	1660	LYS	3
1	A	889	GLU	3
1	A	848	LEU	3
1	A	877	LEU	2
1	A	883	SER	2
1	A	846	ASP	2
1	A	836	LEU	1
2	B	1659	ARG	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 61% for the well-defined parts and 57% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	755
Number of shifts mapped to atoms	755
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	68	-0.34 ± 0.25	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	63	0.18 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}'$	66	0.00 ± 0.20	None needed (< 0.5 ppm)
^{15}N	64	1.12 ± 0.36	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 61%, i.e. 599 atoms were assigned a chemical shift out of a possible 976. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	290/375 (77%)	116/149 (78%)	118/154 (77%)	56/72 (78%)
Sidechain	297/517 (57%)	186/308 (60%)	111/184 (60%)	0/25 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	12/84 (14%)	12/44 (27%)	0/40 (0%)	0/0 (—%)
Overall	599/976 (61%)	314/501 (63%)	229/378 (61%)	56/97 (58%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 57%, i.e. 667 atoms were assigned a chemical shift out of a possible 1163. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	330/460 (72%)	132/183 (72%)	134/188 (71%)	64/89 (72%)
Sidechain	325/593 (55%)	205/351 (58%)	120/215 (56%)	0/27 (0%)
Aromatic	12/110 (11%)	12/58 (21%)	0/52 (0%)	0/0 (—%)
Overall	667/1163 (57%)	349/592 (59%)	254/455 (56%)	64/116 (55%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

