



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2021 – 12:32 PM JST

PDB ID : 7C0K
Title : Crystal structure of a dinucleotide-binding protein of ABC transporter endogenously bound to uridylyl-3'-5'-phospho-guanosine (Form II)
Authors : Kanaujia, S.P.; Chandravanshi, M.
Deposited on : 2020-05-01
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div></div> <div>91%</div> <div>5%</div> <div>••</div> </div>
1	B	406	<div> <div></div> <div>91%</div> <div>7%</div> <div>•</div> </div>

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	EDO	B	517	-	-	X	-

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 7081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sugar ABC transporter, periplasmic sugar-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	396	Total	C	N	O	S	0	5	0
			3082	1998	520	560	4			
1	B	397	Total	C	N	O	S	0	4	0
			3086	1997	521	563	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP Q5SLB4
A	0	MET	-	expression tag	UNP Q5SLB4
A	399	HIS	-	expression tag	UNP Q5SLB4
A	400	HIS	-	expression tag	UNP Q5SLB4
A	401	HIS	-	expression tag	UNP Q5SLB4
A	402	HIS	-	expression tag	UNP Q5SLB4
A	403	HIS	-	expression tag	UNP Q5SLB4
A	404	HIS	-	expression tag	UNP Q5SLB4
B	-1	MET	-	initiating methionine	UNP Q5SLB4
B	0	MET	-	expression tag	UNP Q5SLB4
B	399	HIS	-	expression tag	UNP Q5SLB4
B	400	HIS	-	expression tag	UNP Q5SLB4
B	401	HIS	-	expression tag	UNP Q5SLB4
B	402	HIS	-	expression tag	UNP Q5SLB4
B	403	HIS	-	expression tag	UNP Q5SLB4
B	404	HIS	-	expression tag	UNP Q5SLB4


- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	5	Total	Cl	0	0
			5	5		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CARBON DIOXIDE (three-letter code: CO2) (formula: CO₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is HYPOPHOSPHITE (three-letter code: PO2) (formula: O₂P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			3	2	1		
5	A	1	Total	O	P	0	0
			3	2	1		
5	A	1	Total	O	P	0	0
			3	2	1		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: $C_3H_8O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			5	3	2		
8	A	1	Total	C	O	0	0
			5	3	2		
8	B	1	Total	C	O	0	0
			5	3	2		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is 2-HYDROXY BUTANE-1,4-DIOL (three-letter code: BGQ) (formula: C₄H₉O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 11 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 12 is [(1S,3R,3aR,6aS)-3-(2-azanyl-6-oxidanylidene-1H-purin-9-yl)-5,5-bis(oxidanyl)-1,3,3a,4,6,6a-hexahydrocyclopenta[c]furan-1-yl]methyl [(2R,3S,4R,5R)-5-[2,4-bis(oxidanylidene)pyrimidin-1-yl]-2-(hydroxymethyl)-4-oxidanyl-oxolan-3-yl] hydrogen phosphate (three-letter code: FGO) (formula: C₂₂H₂₈N₇O₁₃P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	A	1	Total	C	N	O	P	0	0
			43	22	7	13	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	B	1	Total	C	N	O	P	0	0
			43	22	7	13	1		

- Molecule 13 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	B	1	Total	C	O	0	0
			4	1	3		
13	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 14 is PHOSPHITE ION (three-letter code: PO3) (formula: O₃P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	B	1	Total	O	P	0	0
			4	3	1		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	310	Total	O	0	0
			310	310		
15	B	327	Total	O	0	0
			327	327		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sugar ABC transporter, periplasmic sugar-binding protein



- Molecule 1: Sugar ABC transporter, periplasmic sugar-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.17Å 57.52Å 121.14Å 90.00° 94.99° 90.00°	Depositor
Resolution (Å)	46.57 – 1.80 46.57 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.57-1.80) 99.9 (46.57-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.146 , 0.176 0.159 , 0.188	Depositor DCC
R_{free} test set	4082 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.235	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7081	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CO3, PO4, FGO, CO2, CL, PDO, PO2, EDO, PO3, PG4, BGQ, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.01	6/3167 (0.2%)	1.09	11/4296 (0.3%)
1	B	1.00	7/3168 (0.2%)	1.07	7/4295 (0.2%)
All	All	1.00	13/6335 (0.2%)	1.08	18/8591 (0.2%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	297	GLU	CD-OE2	14.43	1.41	1.25
1	B	297	GLU	CD-OE2	8.10	1.34	1.25
1	B	370	GLU	CD-OE2	8.02	1.34	1.25
1	B	80	GLU	CD-OE2	7.70	1.34	1.25
1	B	370	GLU	CD-OE1	7.64	1.34	1.25

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	ARG	NE-CZ-NH2	-8.81	115.89	120.30
1	A	114[A]	ARG	NE-CZ-NH1	7.47	124.03	120.30
1	A	114[B]	ARG	NE-CZ-NH1	7.47	124.03	120.30
1	B	346	TYR	CB-CG-CD1	7.39	125.43	121.00
1	A	327	TYR	CB-CG-CD1	-7.03	116.78	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3082	0	3137	21	0
1	B	3086	0	3130	11	0
2	A	1	0	0	0	0
2	B	5	0	0	1	0
3	A	1	0	0	0	0
4	A	12	0	0	0	0
4	B	6	0	0	0	0
5	A	9	0	0	0	0
6	A	5	0	0	0	0
7	A	16	0	24	5	0
7	B	28	0	41	10	0
8	A	10	0	16	0	0
8	B	5	0	8	0	0
9	A	36	0	48	3	0
9	B	24	0	32	3	0
10	A	7	0	9	2	0
11	A	13	0	18	0	0
12	A	43	0	0	0	0
12	B	43	0	0	0	0
13	B	8	0	0	0	0
14	B	4	0	0	0	0
15	A	310	0	0	14	0
15	B	327	0	0	14	1
All	All	7081	0	6463	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:517:GOL:H2	15:A:672:HOH:O	1.32	1.24
1:A:379:LYS:HD3	15:A:741:HOH:O	1.56	1.02
7:B:516:EDO:H11	15:B:733:HOH:O	1.72	0.87
7:A:511:EDO:H21	15:A:782:HOH:O	1.75	0.85
1:B:171:GLN:OE1	15:B:601:HOH:O	1.95	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
15:B:606:HOH:O	15:B:891:HOH:O[2_656]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/406 (98%)	393 (98%)	6 (2%)	0	100	100
1	B	399/406 (98%)	393 (98%)	6 (2%)	0	100	100
All	All	798/812 (98%)	786 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/319 (98%)	312 (99%)	2 (1%)	86	84
1	B	314/319 (98%)	309 (98%)	5 (2%)	62	54
All	All	628/638 (98%)	621 (99%)	7 (1%)	73	68

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	GLU
1	B	191	ASP
1	B	395	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	226	ASN
1	B	1	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	267	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 48 ligands modelled in this entry, 7 are monoatomic - leaving 41 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	EDO	B	512	-	3,3,3	0.15	0	2,2,2	0.11	0
7	EDO	B	517	-	3,3,3	0.56	0	2,2,2	0.56	0
5	PO2	A	509	-	0,2,2	0.00	-	0,1,1	0.00	-
7	EDO	A	514	-	3,3,3	0.53	0	2,2,2	0.24	0
5	PO2	A	507	-	0,2,2	0.00	-	0,1,1	0.00	-
8	PDO	A	516	-	4,4,4	0.17	0	3,3,3	0.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	PG4	A	524	-	12,12,12	0.25	0	11,11,11	0.30	0
7	EDO	B	514	-	3,3,3	0.34	0	2,2,2	0.43	0
9	GOL	A	520	-	5,5,5	0.17	0	5,5,5	0.38	0
9	GOL	B	520	-	5,5,5	0.05	0	5,5,5	0.23	0
9	GOL	A	522	-	5,5,5	0.17	0	5,5,5	0.60	0
4	CO2	A	506	-	2,2,2	0.04	0	1,1,1	0.98	0
7	EDO	A	511	-	3,3,3	0.22	0	2,2,2	0.50	0
7	EDO	A	513	-	3,3,3	0.58	0	2,2,2	0.96	0
13	CO3	B	508	-	0,3,3	0.00	-	0,3,3	0.00	-
12	FGO	B	523	-	40,48,48	1.90	10 (25%)	39,75,75	2.07	8 (20%)
9	GOL	B	521	-	5,5,5	0.05	0	5,5,5	0.20	0
7	EDO	B	515	-	3,3,3	0.21	0	2,2,2	0.35	0
14	PO3	B	510	-	0,3,3	0.00	-	0,3,3	0.00	-
7	EDO	B	516	-	3,3,3	0.49	0	2,2,2	1.44	0
9	GOL	A	519	-	5,5,5	0.11	0	5,5,5	0.31	0
4	CO2	A	505	-	2,2,2	0.20	0	1,1,1	0.82	0
9	GOL	A	521	-	5,5,5	0.16	0	5,5,5	0.69	0
4	CO2	A	503	-	2,2,2	0.24	0	1,1,1	0.71	0
9	GOL	A	517	-	5,5,5	0.23	0	5,5,5	0.35	0
4	CO2	A	504	-	2,2,2	0.18	0	1,1,1	0.95	0
9	GOL	A	518	-	5,5,5	0.10	0	5,5,5	0.38	0
12	FGO	A	525	-	40,48,48	1.92	10 (25%)	39,75,75	1.44	5 (12%)
4	CO2	B	507	-	2,2,2	0.19	0	1,1,1	0.72	0
7	EDO	B	511	-	3,3,3	0.62	0	2,2,2	0.25	0
8	PDO	A	515	-	4,4,4	0.31	0	3,3,3	0.25	0
6	PO4	A	510	-	4,4,4	0.65	0	6,6,6	0.48	0
9	GOL	B	522	-	5,5,5	0.16	0	5,5,5	0.64	0
7	EDO	B	513	-	3,3,3	0.12	0	2,2,2	0.09	0
13	CO3	B	509	-	0,3,3	0.00	-	0,3,3	0.00	-
10	BGQ	A	523	-	6,6,6	0.46	0	4,6,6	0.73	0
5	PO2	A	508	-	0,2,2	0.00	-	0,1,1	0.00	-
8	PDO	B	518	-	4,4,4	0.21	0	3,3,3	0.48	0
4	CO2	B	506	-	2,2,2	0.37	0	1,1,1	0.67	0
9	GOL	B	519	-	5,5,5	0.06	0	5,5,5	0.22	0
7	EDO	A	512	-	3,3,3	0.15	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	512	-	-	1/1/1/1	-
7	EDO	B	517	-	-	0/1/1/1	-
7	EDO	A	514	-	-	1/1/1/1	-
8	PDO	A	516	-	-	1/2/2/2	-
11	PG4	A	524	-	-	1/10/10/10	-
7	EDO	B	514	-	-	1/1/1/1	-
9	GOL	A	520	-	-	2/4/4/4	-
9	GOL	B	520	-	-	4/4/4/4	-
9	GOL	A	522	-	-	2/4/4/4	-
7	EDO	A	511	-	-	0/1/1/1	-
7	EDO	A	513	-	-	1/1/1/1	-
12	FGO	B	523	-	-	1/15/63/63	0/6/6/6
9	GOL	B	521	-	-	4/4/4/4	-
7	EDO	B	515	-	-	0/1/1/1	-
7	EDO	B	516	-	-	0/1/1/1	-
9	GOL	A	519	-	-	4/4/4/4	-
9	GOL	A	521	-	-	2/4/4/4	-
9	GOL	A	517	-	-	3/4/4/4	-
9	GOL	A	518	-	-	4/4/4/4	-
12	FGO	A	525	-	-	0/15/63/63	0/6/6/6
7	EDO	B	511	-	-	1/1/1/1	-
8	PDO	A	515	-	-	1/2/2/2	-
9	GOL	B	522	-	-	4/4/4/4	-
7	EDO	B	513	-	-	1/1/1/1	-
10	BGQ	A	523	-	-	3/5/5/5	-
8	PDO	B	518	-	-	1/2/2/2	-
9	GOL	B	519	-	-	3/4/4/4	-
7	EDO	A	512	-	-	1/1/1/1	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	A	525	FGO	C8-N	5.70	1.42	1.35
12	A	525	FGO	C19-C18	-5.65	1.44	1.53
12	B	523	FGO	C21-C17	-5.37	1.45	1.53
12	B	523	FGO	C8-N	4.71	1.41	1.35
12	B	523	FGO	C19-C18	-3.90	1.47	1.53

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	B	523	FGO	C11-N2-C12	-5.58	116.84	126.64
12	B	523	FGO	C15-C13-C12	-4.56	116.44	120.80
12	B	523	FGO	C13-C15-N4	-4.54	117.22	123.43
12	A	525	FGO	C11-N2-C12	-4.23	119.21	126.64
12	B	523	FGO	C15-N4-C16	4.05	122.37	115.93

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	515	PDO	O1-C1-C2-C3
9	A	518	GOL	O1-C1-C2-C3
9	A	518	GOL	C1-C2-C3-O3
9	A	519	GOL	O1-C1-C2-C3
9	A	519	GOL	C1-C2-C3-O3


There are no ring outliers.

11 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	517	EDO	4	0
7	B	514	EDO	3	0
9	A	520	GOL	1	0
7	A	511	EDO	3	0
7	A	513	EDO	2	0
9	B	521	GOL	2	0
7	B	516	EDO	3	0
9	A	519	GOL	1	0
9	A	517	GOL	1	0
9	B	522	GOL	1	0
10	A	523	BGQ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	396/406 (97%)	-0.14	5 (1%) 77 74	9, 18, 41, 66	0
1	B	397/406 (97%)	-0.17	7 (1%) 68 64	9, 17, 34, 60	0
All	All	793/812 (97%)	-0.15	12 (1%) 73 70	9, 17, 36, 66	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	ARG	4.4
1	A	67	PHE	4.2
1	B	12	ARG	4.1
1	A	68	ALA	4.1
1	B	67	PHE	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	EDO	B	512	4/4	0.67	0.19	50,50,51,54	0
10	BGQ	A	523	7/7	0.75	0.18	47,49,55,56	0
4	CO2	A	504	3/3	0.76	0.12	45,45,48,51	0
7	EDO	B	514	4/4	0.77	0.18	37,39,42,47	0
7	EDO	A	512	4/4	0.77	0.18	42,47,48,50	0
4	CO2	A	505	3/3	0.80	0.14	53,53,56,57	0
7	EDO	B	515	4/4	0.81	0.19	38,41,44,47	0
8	PDO	B	518	5/5	0.81	0.18	34,41,46,51	0
5	PO2	A	509	3/3	0.81	0.14	52,52,60,65	0
13	CO3	B	509	4/4	0.82	0.20	35,36,39,39	0
14	PO3	B	510	4/4	0.82	0.13	49,54,63,65	0
13	CO3	B	508	4/4	0.83	0.27	42,46,46,50	0
9	GOL	A	517	6/6	0.84	0.24	29,33,33,34	0
11	PG4	A	524	13/13	0.85	0.15	31,36,45,52	0
9	GOL	A	522	6/6	0.86	0.28	30,41,44,47	0
7	EDO	A	511	4/4	0.86	0.12	31,33,37,39	0
9	GOL	B	521	6/6	0.87	0.18	45,47,48,48	0
8	PDO	A	516	5/5	0.87	0.16	39,44,50,52	0
5	PO2	A	507	3/3	0.88	0.14	50,50,52,58	0
8	PDO	A	515	5/5	0.88	0.13	40,40,42,48	0
7	EDO	B	513	4/4	0.88	0.12	49,52,53,59	0
7	EDO	B	511	4/4	0.88	0.18	39,40,44,46	0
9	GOL	A	521	6/6	0.89	0.16	23,30,35,41	0
2	CL	B	504	1/1	0.90	0.09	65,65,65,65	0
9	GOL	A	519	6/6	0.90	0.16	39,42,43,46	0
9	GOL	B	519	6/6	0.90	0.20	30,49,52,59	0
2	CL	A	501	1/1	0.91	0.07	37,37,37,37	0
9	GOL	B	520	6/6	0.91	0.18	22,40,46,54	0
7	EDO	B	516	4/4	0.91	0.14	25,26,27,29	0
4	CO2	A	506	3/3	0.91	0.16	42,42,43,52	0
6	PO4	A	510	5/5	0.92	0.16	49,50,56,58	0
9	GOL	A	518	6/6	0.93	0.18	18,28,39,48	0
7	EDO	A	514	4/4	0.93	0.14	24,29,33,40	0
4	CO2	B	507	3/3	0.94	0.11	28,28,32,40	0
9	GOL	A	520	6/6	0.94	0.12	19,22,24,28	0
2	CL	B	503	1/1	0.95	0.09	31,31,31,31	0
2	CL	B	502	1/1	0.95	0.07	48,48,48,48	0
3	NA	A	502	1/1	0.95	0.15	34,34,34,34	0
7	EDO	A	513	4/4	0.95	0.14	22,26,29,31	0
4	CO2	B	506	3/3	0.96	0.21	27,27,27,38	0
4	CO2	A	503	3/3	0.96	0.10	23,23,24,42	0
12	FGO	A	525	43/43	0.96	0.09	11,15,23,31	0
12	FGO	B	523	43/43	0.96	0.10	9,15,23,28	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	B	505	1/1	0.96	0.05	42,42,42,42	0
5	PO2	A	508	3/3	0.96	0.09	33,33,38,44	0
9	GOL	B	522	6/6	0.96	0.16	22,26,28,29	0
7	EDO	B	517	4/4	0.97	0.20	22,25,25,28	0
2	CL	B	501	1/1	0.99	0.07	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.