



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 01:29 PM BST

PDB ID : 5C1U  
Title : Crystal structure of EV71 3C Proteinase in complex with Compound Xb  
Authors : Zhang, L.; Huang, G.; Cai, Q.; Zhao, C.; Ren, H.; Li, P.; Li, N.; Chen, S.; Li, J.; Lin, T.  
Deposited on : 2015-06-15  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	192	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>
1	B	192	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	1	0
			1407	892	250	257	8			
1	B	181	Total	C	N	O	S	0	2	0
			1417	898	255	256	8			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP A9XG43
A	184	LEU	-	expression tag	UNP A9XG43
A	185	GLU	-	expression tag	UNP A9XG43
A	186	HIS	-	expression tag	UNP A9XG43
A	187	HIS	-	expression tag	UNP A9XG43
A	188	HIS	-	expression tag	UNP A9XG43
A	189	HIS	-	expression tag	UNP A9XG43
A	190	HIS	-	expression tag	UNP A9XG43
A	191	HIS	-	expression tag	UNP A9XG43
B	0	MET	-	expression tag	UNP A9XG43
B	184	LEU	-	expression tag	UNP A9XG43
B	185	GLU	-	expression tag	UNP A9XG43
B	186	HIS	-	expression tag	UNP A9XG43
B	187	HIS	-	expression tag	UNP A9XG43
B	188	HIS	-	expression tag	UNP A9XG43
B	189	HIS	-	expression tag	UNP A9XG43
B	190	HIS	-	expression tag	UNP A9XG43
B	191	HIS	-	expression tag	UNP A9XG43

- Molecule 2 is (2S)-2-[[[(E)-3-[4-(dimethylamino)phenyl]prop-2-enoyl]amino]-N-[(2S)-1-oxidan-3-yl-3-[(3S)-2-oxidanylidene-pyrrolidin-3-yl]propan-2-yl]-3-phenyl-propanamide (three-letter code: GHV) (formula: C<sub>27</sub>H<sub>34</sub>N<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			35	27	4	4		
2	B	1	Total	C	N	O	0	0
			35	27	4	4		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	178	Total	O	0	0
			178	178		
3	B	147	Total	O	0	0
			147	147		



- Molecule 1: 3C proteinase



- |    |    |    |     |     |     |     |     |     |     |     |     |     |      |      |      |      |      |      |      |      |      |      |      |      |      |      |     |     |     |     |     |     |     |     |     |     |     |
|----|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| H0 | D5 | S9 | R12 | R13 | R16 | H24 | R31 | E50 | E51 | R84 | K88 | F89 | S111 | D118 | Q121 | L127 | H133 | Y138 | N139 | V154 | H161 | R167 | Q168 | Y178 | F179 | A180 | SER | GLU | GLN | LEU | GLU | HIS | HIS | HIS | HIS | HIS | HIS |
|----|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.54Å 64.22Å 75.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.75 – 1.49 38.75 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.75-1.49) 99.5 (38.75-1.49)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.19 (at 1.49Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.181 , 0.212 0.178 , 0.209	Depositor DCC
$R_{free}$ test set	2584 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.3	Xtriage
Anisotropy	0.758	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.002 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 67.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.9904e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GHV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.12	2/1437 (0.1%)	1.34	11/1945 (0.6%)
1	B	1.21	3/1450 (0.2%)	1.23	9/1961 (0.5%)
All	All	1.17	5/2887 (0.2%)	1.29	20/3906 (0.5%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	12	ARG	CZ-NH2	-6.13	1.25	1.33
1	B	9	SER	CB-OG	6.08	1.50	1.42
1	B	178	TYR	CE1-CZ	6.00	1.46	1.38
1	B	111	SER	CB-OG	5.11	1.48	1.42
1	A	12	ARG	CZ-NH1	5.09	1.39	1.33

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	12	ARG	NE-CZ-NH2	-20.75	109.93	120.30
1	A	12	ARG	NE-CZ-NH1	14.19	127.40	120.30
1	A	118	ASP	CB-CG-OD1	8.02	125.52	118.30
1	B	167	ARG	CG-CD-NE	7.43	127.40	111.80
1	A	84	ARG	NE-CZ-NH1	-6.97	116.81	120.30
1	B	13[A]	ARG	NE-CZ-NH1	-6.70	116.95	120.30
1	B	13[B]	ARG	NE-CZ-NH1	-6.70	116.95	120.30
1	A	118	ASP	CB-CG-OD2	-6.40	112.54	118.30
1	B	81	GLU	OE1-CD-OE2	5.98	130.48	123.30
1	A	16	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	A	134	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	B	5	ASP	CB-CG-OD2	-5.73	113.14	118.30
1	B	118	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	138	TYR	CB-CG-CD2	-5.43	117.74	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	12	ARG	NE-CZ-NH2	5.34	122.97	120.30
1	A	88	LYS	CD-CE-NZ	-5.27	99.57	111.70
1	B	31	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	A	85	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	176	ARG	NE-CZ-NH1	-5.02	117.79	120.30
1	B	138	TYR	CB-CG-CD2	-5.00	118.00	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1407	0	1428	16	0
1	B	1417	0	1446	17	0
2	A	35	0	0	1	0
2	B	35	0	0	1	0
3	A	178	0	0	5	2
3	B	147	0	0	4	1
All	All	3219	0	2874	33	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:LYS:NZ	1:B:89:PHE:CE2	1.89	1.40
1:B:88:LYS:NZ	1:B:89:PHE:CZ	2.06	1.23
1:B:154:VAL:HB	3:B:366:HOH:O	1.50	1.10
1:B:52:LYS:NZ	3:B:301:HOH:O	1.95	0.97
1:B:138:TYR:OH	1:B:161:HIS:HD2	1.75	0.70
1:A:139:ASN:HD22	1:A:168:GLN:HG2	1.57	0.69
1:B:13[B]:ARG:HH11	1:B:13[B]:ARG:HG2	1.61	0.66
1:A:138:TYR:OH	1:A:161:HIS:HD2	1.80	0.65

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:LEU:CD1	1:A:163:GLY:HA3	2.31	0.60
1:B:13[B]:ARG:HH11	1:B:13[B]:ARG:CG	2.14	0.60
1:B:13[A]:ARG:NE	1:B:84:ARG:HD3	2.17	0.60
1:A:24:HIS:HD2	3:A:364:HOH:O	1.84	0.60
1:B:139:ASN:HD22	1:B:168:GLN:HG2	1.66	0.60
1:B:24:HIS:HD2	3:B:376:HOH:O	1.86	0.59
1:A:127:LEU:HD11	1:A:163:GLY:HA3	1.86	0.56
1:A:139:ASN:HD21	1:A:168:GLN:HE21	1.53	0.55
1:A:24:HIS:HE1	3:A:461:HOH:O	1.88	0.55
1:B:139:ASN:HD21	1:B:168:GLN:HE21	1.56	0.54
1:B:139:ASN:ND2	1:B:168:GLN:HE21	2.09	0.50
1:A:156:LYS:HE3	3:A:368:HOH:O	2.13	0.47
1:A:17:GLN:HB2	1:A:50:GLU:HG2	1.97	0.47
1:A:12:ARG:NH2	3:A:302:HOH:O	2.39	0.46
1:B:154:VAL:CG2	3:B:366:HOH:O	2.61	0.45
1:B:16[B]:ARG:NE	1:B:16[B]:ARG:HA	2.32	0.44
1:B:88:LYS:NZ	1:B:89:PHE:CD2	2.68	0.44
1:A:139:ASN:ND2	1:A:168:GLN:HG2	2.29	0.43
1:A:139:ASN:ND2	1:A:168:GLN:HE21	2.17	0.43
1:A:28:LEU:HD11	1:A:86:ILE:HD11	2.01	0.42
1:A:161:HIS:HE1	2:A:201:GHV:O1	2.02	0.42
1:B:161:HIS:HE1	2:B:201:GHV:O1	2.04	0.41
1:B:138:TYR:OH	1:B:161:HIS:CD2	2.65	0.41
1:A:143:LYS:NZ	3:A:301:HOH:O	2.12	0.40
1:A:19[B]:GLN:HG2	1:A:24:HIS:CD2	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:307:HOH:O	3:A:458:HOH:O[3_544]	1.51	0.69
3:A:354:HOH:O	3:B:401:HOH:O[4_444]	2.18	0.02

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/192 (94%)	175 (97%)	5 (3%)	0	100	100
1	B	181/192 (94%)	177 (98%)	3 (2%)	1 (1%)	25	7
All	All	361/384 (94%)	352 (98%)	8 (2%)	1 (0%)	41	18

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	50	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/166 (94%)	155 (99%)	1 (1%)	86	74
1	B	157/166 (95%)	154 (98%)	3 (2%)	57	27
All	All	313/332 (94%)	309 (99%)	4 (1%)	69	44

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	TYR
1	B	121	GLN
1	B	133	HIS
1	B	138	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	24	HIS
1	A	42	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	51	HIS
1	A	80	ASN
1	A	108	HIS
1	A	139	ASN
1	A	161	HIS
1	B	19	GLN
1	B	22	GLN
1	B	24	HIS
1	B	139	ASN
1	B	161	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GHV	A	201	1	37,37,37	1.38	5 (13%)	46,49,49	1.74	14 (30%)
2	GHV	B	201	1	37,37,37	1.43	4 (10%)	46,49,49	1.63	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GHV	A	201	1	-	6/31/41/41	0/3/3/3
2	GHV	B	201	1	-	5/31/41/41	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	GHV	C34-N33	-4.15	1.29	1.33
2	B	201	GHV	C2-N7	3.20	1.44	1.37
2	A	201	GHV	C2-N7	3.12	1.44	1.37
2	A	201	GHV	C16-C15	-2.72	1.47	1.54
2	A	201	GHV	C31-C32	2.69	1.57	1.53
2	B	201	GHV	C15-C23	-2.61	1.46	1.52
2	A	201	GHV	C34-N33	-2.34	1.31	1.33
2	B	201	GHV	C30-C34	-2.10	1.49	1.52
2	A	201	GHV	C27-C26	2.06	1.55	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	GHV	C9-N7-C2	-5.22	111.59	120.54
2	B	201	GHV	C9-N7-C2	-4.38	113.03	120.54
2	B	201	GHV	C29-C26-C27	-4.20	105.81	111.65
2	B	201	GHV	C8-N7-C2	-3.80	114.03	120.54
2	A	201	GHV	C32-C31-C30	-3.17	100.61	105.75
2	A	201	GHV	C8-N7-C2	-3.03	115.35	120.54
2	B	201	GHV	C32-C31-C30	-2.97	100.94	105.75
2	A	201	GHV	O1-C34-C30	-2.80	122.94	126.23
2	A	201	GHV	C20-C19-C18	-2.74	116.02	120.19
2	A	201	GHV	C16-C15-N13	2.57	116.21	110.79
2	B	201	GHV	C26-N24-C23	-2.54	119.16	123.20
2	B	201	GHV	C16-C15-C23	2.47	116.66	110.25
2	A	201	GHV	C8-N7-C9	-2.36	108.52	116.12
2	A	201	GHV	C10-C11-C12	-2.19	117.27	121.56
2	A	201	GHV	C11-C12-N13	2.18	118.75	114.56
2	A	201	GHV	C21-C22-C17	-2.18	117.29	120.63
2	A	201	GHV	C17-C16-C15	2.10	119.18	113.39
2	B	201	GHV	C20-C21-C22	-2.07	117.03	120.19
2	A	201	GHV	C29-C30-C31	-2.07	109.71	117.31

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	GHV	C8-N7-C9	-2.06	109.47	116.12
2	B	201	GHV	C17-C16-C15	2.04	119.01	113.39
2	A	201	GHV	C5-C10-C11	-2.03	122.26	126.91
2	A	201	GHV	O1-C34-N33	2.01	128.33	125.54

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	201	GHV	C1-C2-N7-C9
2	B	201	GHV	C3-C2-N7-C9
2	A	201	GHV	C1-C2-N7-C9
2	A	201	GHV	C3-C2-N7-C9
2	A	201	GHV	C10-C11-C12-O14
2	A	201	GHV	C10-C11-C12-N13
2	A	201	GHV	C11-C10-C5-C6
2	B	201	GHV	C10-C11-C12-O14
2	B	201	GHV	C10-C11-C12-N13
2	A	201	GHV	C11-C10-C5-C4
2	B	201	GHV	C15-C16-C17-C22

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	GHV	1	0
2	B	201	GHV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/192 (94%)	-0.14	2 (1%) 80 84	9, 13, 25, 46	0
1	B	181/192 (94%)	-0.10	2 (1%) 80 84	9, 14, 26, 34	0
All	All	362/384 (94%)	-0.12	4 (1%) 80 84	9, 14, 26, 46	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	LEU	7.0
1	A	128	SER	3.0
1	B	51	HIS	2.7
1	B	127	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GHV	A	201	35/35	0.92	0.16	13,20,85,89	0


*Continued on next page...*



*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GHV	B	201	35/35	0.95	0.16	13,19,98,103	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around GHV B 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.