



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 04:21 pm BST

PDB ID : 3C3R  
Title : ALIX BRO1 CHMP4C complex  
Authors : McCullough, J.B.; Fisher, R.D.; Whitby, F.G.; Sundquist, W.I.; Hill, C.P.  
Deposited on : 2008-01-28  
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	<div> <div>16%</div> <div>72%</div> <div>19%</div> <div>• 6%</div> </div>
2	B	13	<div> <div>23%</div> <div>54%</div> <div>46%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Programmed cell death 6-interacting protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2803	1792	471	529	11	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q8WUM4
A	-19	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-18	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-17	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-16	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-15	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-14	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-13	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-12	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-11	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-10	HIS	-	EXPRESSION TAG	UNP Q8WUM4
A	-9	SER	-	EXPRESSION TAG	UNP Q8WUM4
A	-8	GLY	-	EXPRESSION TAG	UNP Q8WUM4
A	-7	GLN	-	EXPRESSION TAG	UNP Q8WUM4
A	-6	ASN	-	EXPRESSION TAG	UNP Q8WUM4
A	-5	LEU	-	EXPRESSION TAG	UNP Q8WUM4
A	-4	TYR	-	EXPRESSION TAG	UNP Q8WUM4
A	-3	PHE	-	EXPRESSION TAG	UNP Q8WUM4
A	-2	GLN	-	EXPRESSION TAG	UNP Q8WUM4
A	-1	GLY	-	EXPRESSION TAG	UNP Q8WUM4
A	0	HIS	-	EXPRESSION TAG	UNP Q8WUM4

- Molecule 2 is a protein called Charged multivesicular body protein 4c peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	13	Total	C	N	O	0	0	0
			104	64	16	24			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	55	Total	O	0	0
			55	55		
4	B	1	Total	O	0	0
			1	1		



- Molecule 1: Programmed cell death 6-interacting protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.91Å 62.40Å 76.35Å 90.00° 121.62° 90.00°	Depositor
Resolution (Å)	33.86 – 2.02 33.86 – 2.02	Depositor EDS
% Data completeness (in resolution range)	95.3 (33.86-2.02) 95.3 (33.86-2.02)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.243 , 0.288 0.259 , 0.306	Depositor DCC
$R_{free}$ test set	2168 reflections (7.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.9	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2969	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.75	0/2856	0.88	6/3854 (0.2%)
2	B	0.47	0/105	1.03	2/141 (1.4%)
All	All	0.74	0/2961	0.89	8/3995 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	314	ASP	CB-CG-OD2	6.86	124.47	118.30
1	A	299	ASP	CB-CG-OD2	6.51	124.16	118.30
1	A	302	ASP	CB-CG-OD2	5.87	123.58	118.30
1	A	226	ASP	CB-CG-OD2	5.81	123.53	118.30
2	B	223	ASP	CB-CG-OD2	5.79	123.51	118.30
1	A	210	ASP	CB-CG-OD2	5.68	123.41	118.30
2	B	222	ASP	CB-CG-OD2	5.04	122.83	118.30
1	A	235	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2803	0	2828	44	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	104	0	92	5	0
3	A	6	0	8	0	0
4	A	55	0	0	4	0
4	B	1	0	0	0	0
All	All	2969	0	2928	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:LYS:HB3	1:A:20:PRO:HD3	1.68	0.76
1:A:337:LEU:HD21	2:B:228:LEU:HB3	1.68	0.74
1:A:147:LYS:NZ	2:B:231:TRP:O	2.21	0.69
1:A:78:ILE:HD11	1:A:355:PHE:CD1	2.28	0.69
1:A:129:ALA:O	1:A:133:GLN:HG3	1.94	0.66
1:A:74:GLN:HE21	1:A:78:ILE:HD13	1.61	0.66
1:A:91:LEU:O	1:A:114:ALA:HA	1.96	0.66
1:A:56:ARG:HG2	4:A:3599:HOH:O	1.97	0.64
1:A:337:LEU:CD2	2:B:228:LEU:HB3	2.27	0.64
1:A:163:ILE:O	1:A:167:VAL:HG22	1.98	0.64
1:A:101:LYS:H	1:A:101:LYS:HD3	1.63	0.63
1:A:216:LEU:HD21	1:A:337:LEU:HD13	1.82	0.61
1:A:216:LEU:CD2	1:A:337:LEU:HD13	2.35	0.56
1:A:357:LYS:HA	1:A:357:LYS:HE3	1.89	0.55
1:A:58:LEU:HD21	4:A:3585:HOH:O	2.06	0.54
1:A:71:TYR:CZ	1:A:75:ILE:HD11	2.42	0.54
1:A:284:ILE:HD12	1:A:297:VAL:HG22	1.90	0.54
1:A:139:ASN:HD22	1:A:139:ASN:C	2.11	0.54
1:A:158:GLY:HA2	1:A:340:SER:HB3	1.90	0.54
1:A:284:ILE:HD12	1:A:297:VAL:CG2	2.39	0.53
1:A:219:GLN:HG2	1:A:336:THR:HB	1.93	0.51
1:A:50:ARG:O	1:A:54:VAL:HG22	2.09	0.51
1:A:201:LEU:O	4:A:3557:HOH:O	2.19	0.51
1:A:150:ALA:O	1:A:154:GLN:HG3	2.14	0.48
1:A:59:ASP:HB3	1:A:64:ALA:HB2	1.96	0.47
1:A:162:HIS:O	1:A:166:THR:HB	2.15	0.47
1:A:212:ILE:HD11	2:B:221:GLU:HA	1.96	0.47
1:A:78:ILE:HG22	1:A:116:LEU:HD21	1.97	0.47
1:A:101:LYS:H	1:A:101:LYS:CD	2.28	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:CYS:HB2	1:A:242:PHE:HD1	1.80	0.46
1:A:56:ARG:CG	4:A:3599:HOH:O	2.59	0.45
1:A:292:ASP:OD2	1:A:292:ASP:N	2.48	0.45
1:A:15:VAL:HG12	1:A:17:LEU:HD23	1.99	0.44
1:A:24:PHE:HD2	1:A:82:PHE:HD2	1.66	0.44
1:A:335:ALA:HB1	2:B:225:ILE:HD11	2.00	0.43
1:A:96:LYS:HG3	1:A:110:LYS:HG2	1.99	0.42
1:A:59:ASP:OD1	1:A:61:HIS:HB2	2.19	0.42
1:A:12:THR:HB	1:A:95:TRP:CE3	2.55	0.42
1:A:73:ASP:OD1	1:A:346:PRO:HA	2.20	0.41
1:A:81:LYS:HE3	1:A:81:LYS:HA	2.03	0.41
1:A:69:LEU:HD23	1:A:127:CYS:SG	2.61	0.40
1:A:19:LYS:HB3	1:A:20:PRO:CD	2.46	0.40
1:A:199:PHE:CE1	1:A:338:VAL:HG23	2.56	0.40
1:A:10:LYS:NZ	1:A:96:LYS:O	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/380 (93%)	335 (94%)	18 (5%)	2 (1%)	25	18
2	B	11/13 (85%)	11 (100%)	0	0	100	100
All	All	366/393 (93%)	346 (94%)	18 (5%)	2 (0%)	29	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	GLU
1	A	31	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/320 (93%)	270 (90%)	29 (10%)	8	4
2	B	10/10 (100%)	10 (100%)	0	100	100
All	All	309/330 (94%)	280 (91%)	29 (9%)	8	4

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	28	THR
1	A	56	ARG
1	A	78	ILE
1	A	79	GLU
1	A	81	LYS
1	A	84	PHE
1	A	86	GLU
1	A	87	ASN
1	A	90	CYS
1	A	101	LYS
1	A	139	ASN
1	A	142	ASN
1	A	146	LEU
1	A	165	GLU
1	A	167	VAL
1	A	174	GLU
1	A	176	THR
1	A	182	ASP
1	A	184	VAL
1	A	228	PHE
1	A	296	ASN
1	A	297	VAL
1	A	298	LYS
1	A	337	LEU
1	A	338	VAL
1	A	339	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	356	GLU
1	A	357	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	133	GLN
1	A	139	ASN
1	A	219	GLN
1	A	296	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	3545	-	5,5,5	0.39	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3545	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3545	GOL	O1-C1-C2-C3
3	A	3545	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/380 (93%)	1.01	59 (16%) <b>1</b> <b>1</b>	17, 30, 64, 76	0
2	B	13/13 (100%)	1.44	3 (23%) <b>0</b> <b>0</b>	36, 40, 51, 52	0
All	All	370/393 (94%)	1.03	62 (16%) <b>1</b> <b>1</b>	17, 30, 64, 76	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	85	SER	10.2
1	A	83	PRO	9.2
1	A	105	PHE	8.0
1	A	84	PHE	7.5
1	A	107	GLY	7.0
1	A	106	GLY	6.7
1	A	28	THR	6.4
1	A	86	GLU	6.3
1	A	21	LEU	6.0
1	A	30	PRO	6.0
1	A	90	CYS	5.8
1	A	31	SER	5.7
1	A	25	ILE	5.4
1	A	103	SER	5.3
1	A	34	GLU	5.2
1	A	33	GLY	5.1
1	A	29	TYR	5.0
1	A	88	GLN	4.9
1	A	57	PRO	4.6
1	A	32	GLY	4.5
1	A	24	PHE	4.5
1	A	39	TYR	4.3
1	A	37	ALA	4.1
1	A	89	ILE	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	35	GLU	3.8
1	A	356	GLU	3.8
1	A	108	SER	3.7
1	A	82	PHE	3.6
1	A	58	LEU	3.5
1	A	81	LYS	3.3
1	A	350	LYS	3.3
1	A	357	LYS	3.2
2	B	221	GLU	3.2
1	A	22	VAL	3.1
1	A	20	PRO	3.1
1	A	124	LEU	3.0
1	A	40	CYS	3.0
1	A	27	GLN	2.9
1	A	358	MET	2.8
1	A	192	LEU	2.8
1	A	2	ALA	2.8
1	A	104	LEU	2.8
2	B	233	THR	2.8
1	A	36	GLN	2.7
1	A	172	SER	2.6
2	B	230	ALA	2.6
1	A	60	LYS	2.5
1	A	191	MET	2.4
1	A	123	VAL	2.3
1	A	61	HIS	2.3
1	A	91	LEU	2.3
1	A	193	ALA	2.3
1	A	190	ILE	2.3
1	A	195	ALA	2.2
1	A	102	GLY	2.2
1	A	41	ARG	2.1
1	A	198	VAL	2.1
1	A	125	PHE	2.1
1	A	200	PHE	2.1
1	A	87	ASN	2.0
1	A	217	ALA	2.0
1	A	127	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	3545	6/6	0.70	0.21	85,85,86,86	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.