



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 07:03 am BST

PDB ID : 5C4X
Title : Crystal structure of a transcribing RNA Polymerase II complex reveals a complete transcription bubble
Authors : Barnes, C.O.; Calero, M.; Malik, I.; Spahr, H.; Zhang, Q.; Pullara, F.; Kaplan, C.D.; Calero, G.
Deposited on : 2015-06-18
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)
RSRZ outliers	127900	1028 (4.34-3.66)
RNA backbone	3102	1048 (5.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	
2	B	1224	
3	C	318	
4	D	221	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	
6	F	155	
7	G	179	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	R	9	
14	S	53	
15	U	53	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
17	MG	A	1803	-	-	-	X
17	MG	A	1804	-	-	-	X

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 33495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1436	Total	C	N	O	S	0	0	0
			11280	7105	1972	2141	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1172	Total	C	N	O	S	0	0	0
			9293	5866	1630	1741	56			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	265	Total	C	N	O	S	0	0	0
			2086	1312	347	414	13			

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	178	Total	C	N	O	S	0	0	0
			1417	875	254	286	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1339	861	222	248	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	172	LEU	-	expression tag	UNP P34087
G	173	GLU	-	expression tag	UNP P34087
G	174	HIS	-	expression tag	UNP P34087
G	175	HIS	-	expression tag	UNP P34087
G	176	HIS	-	expression tag	UNP P34087
G	177	HIS	-	expression tag	UNP P34087
G	178	HIS	-	expression tag	UNP P34087
G	179	HIS	-	expression tag	UNP P34087

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	135	Total	C	N	O	S	0	0	0
			1080	679	182	214	5			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	114	Total	C	N	O	S	0	0	0
			921	568	165	178	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	66	Total	C	N	O	S	0	0	0
			540	345	94	95	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	115	Total	C	N	O	S	0	0	0
			924	593	157	172	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	44	Total	C	N	O	S	0	0	0
			352	217	70	61	4			

- Molecule 13 is a RNA chain called RNA (5'-R(P*UP*CP*GP*AP*GP*AP*GP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	R	9	Total	C	N	O	P	0	0	0
			197	88	40	60	9			

- Molecule 14 is a DNA chain called NON-TEMPLATE STRAND DNA (38-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	S	38	Total	C	N	O	P	0	0	0
			789	373	149	229	38			

- Molecule 15 is a DNA chain called TEMPLATE STRAND DNA (40-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	U	40	Total	C	N	O	P	0	0	0
			809	384	150	235	40			

- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	J	1	Total	Zn	0	0
			1	1		
16	B	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	C	2	Total	Zn	0	0
			2	2		
16	A	2	Total	Zn	0	0
			2	2		
16	L	1	Total	Zn	0	0
			1	1		


- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	2	Total	Mg	0	0
			2	2		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase II subunit RPB1






Chain C: 



- Molecule 4: DNA-directed RNA polymerase II subunit RPB4

Chain D: 




- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1

Chain E: 




- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F: 





- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G: 





- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H: 



- Molecule 9: DNA-directed RNA polymerase II subunit RPB9

Chain I: 




- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5

Chain J: 





- Molecule 11: DNA-directed RNA polymerase II subunit RPB11

Chain K: 




- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4

Chain L: 




- Molecule 13: RNA (5'-R(P*UP*CP*GP*AP*GP*AP*GP*GP*A)-3')

Chain R: 



- Molecule 14: NON-TEMPLATE STRAND DNA (38-MER)

Chain S: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	220.79Å 393.36Å 281.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 4.00 39.65 – 4.00	Depositor EDS
% Data completeness (in resolution range)	96.5 (40.00-4.00) 96.6 (39.65-4.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 4.00Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.210 , 0.227 0.233 , 0.252	Depositor DCC
R_{free} test set	2990 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	137.8	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 155.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.045 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.044 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	33495	wwPDB-VP
Average B, all atoms (Å ²)	197.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/11485	0.69	1/15536 (0.0%)
2	B	0.51	0/9470	0.68	2/12770 (0.0%)
3	C	0.51	0/2124	0.62	0/2879
4	D	0.53	0/1427	0.64	0/1911
5	E	0.51	0/1788	0.61	0/2406
6	F	0.50	0/717	0.65	0/967
7	G	0.48	0/1367	0.66	0/1844
8	H	0.47	0/1097	0.63	0/1484
9	I	0.47	0/939	0.61	0/1266
10	J	0.53	0/549	0.69	0/738
11	K	0.48	0/942	0.61	0/1272
12	L	0.52	0/354	0.69	0/468
13	R	0.84	0/221	0.82	0/343
14	S	1.16	0/885	1.05	0/1364
15	U	1.15	0/906	1.52	7/1392 (0.5%)
All	All	0.57	0/34271	0.72	10/46640 (0.0%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	U	35	DA	O5'-P-OP1	-23.98	81.92	110.70
15	U	35	DA	O5'-P-OP2	-22.60	83.58	110.70
15	U	35	DA	OP1-P-OP2	17.52	145.87	119.60
15	U	34	DG	OP1-P-O3'	-11.59	79.69	105.20
15	U	34	DG	OP2-P-O3'	-11.15	80.66	105.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11280	0	11325	161	0
2	B	9293	0	9282	131	0
3	C	2086	0	2045	42	0
4	D	1417	0	1428	6	0
5	E	1752	0	1776	13	0
6	F	705	0	731	11	0
7	G	1339	0	1357	21	0
8	H	1080	0	1049	12	0
9	I	921	0	870	7	0
10	J	540	0	554	9	0
11	K	924	0	934	9	0
12	L	352	0	378	4	0
13	R	197	0	97	1	0
14	S	789	0	429	8	0
15	U	809	0	447	11	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	C	2	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	2	0	0	0	0
All	All	33495	0	32702	384	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 384 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:746:MET:HE1	2:B:1015:HIS:HA	1.34	1.06
1:A:697:ALA:HA	1:A:702:LEU:HD11	1.42	1.01
14:S:25:DG:H4'	14:S:26:DT:H5'	1.43	1.00
3:C:148:ARG:NH2	10:J:65:PRO:HD3	1.76	0.99
3:C:39:ALA:HA	3:C:164:ALA:HB3	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1430/1733 (82%)	1225 (86%)	144 (10%)	61 (4%)	2	25
2	B	1158/1224 (95%)	963 (83%)	142 (12%)	53 (5%)	2	24
3	C	263/318 (83%)	231 (88%)	25 (10%)	7 (3%)	5	34
4	D	174/221 (79%)	149 (86%)	15 (9%)	10 (6%)	1	19
5	E	212/215 (99%)	196 (92%)	9 (4%)	7 (3%)	4	30
6	F	85/155 (55%)	75 (88%)	8 (9%)	2 (2%)	6	36
7	G	169/179 (94%)	142 (84%)	21 (12%)	6 (4%)	3	28
8	H	129/146 (88%)	108 (84%)	13 (10%)	8 (6%)	1	18
9	I	112/122 (92%)	100 (89%)	9 (8%)	3 (3%)	5	34
10	J	64/70 (91%)	54 (84%)	9 (14%)	1 (2%)	9	44
11	K	113/120 (94%)	106 (94%)	6 (5%)	1 (1%)	17	55
12	L	42/70 (60%)	30 (71%)	9 (21%)	3 (7%)	1	16
All	All	3951/4573 (86%)	3379 (86%)	410 (10%)	162 (4%)	3	25

5 of 162 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	LYS
1	A	66	LYS
1	A	76	GLU
1	A	194	ALA
1	A	319	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1250/1520 (82%)	1167 (93%)	83 (7%)	16	45
2	B	1009/1061 (95%)	956 (95%)	53 (5%)	22	51
3	C	233/274 (85%)	221 (95%)	12 (5%)	23	51
4	D	156/200 (78%)	150 (96%)	6 (4%)	33	59
5	E	196/197 (100%)	191 (97%)	5 (3%)	46	67
6	F	77/137 (56%)	72 (94%)	5 (6%)	17	45
7	G	152/160 (95%)	144 (95%)	8 (5%)	22	51
8	H	118/128 (92%)	114 (97%)	4 (3%)	37	61
9	I	107/116 (92%)	103 (96%)	4 (4%)	34	60
10	J	61/65 (94%)	54 (88%)	7 (12%)	5	25
11	K	99/102 (97%)	94 (95%)	5 (5%)	24	52
12	L	39/57 (68%)	34 (87%)	5 (13%)	4	22
All	All	3497/4017 (87%)	3300 (94%)	197 (6%)	21	49

5 of 197 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	282	ILE
2	B	836	GLU
10	J	9	SER
2	B	333	PHE
2	B	555	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	224	GLN
2	B	494	HIS
3	C	140	ASN
1	A	1390	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1195	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
13	R	7/9 (77%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1436/1733 (82%)	-0.31	0 100 100	62, 176, 266, 296	0
2	B	1172/1224 (95%)	-0.17	12 (1%) 82 74	73, 189, 282, 300	0
3	C	265/318 (83%)	-0.27	0 100 100	95, 184, 251, 288	0
4	D	178/221 (80%)	-0.25	1 (0%) 89 84	128, 206, 266, 293	0
5	E	214/215 (99%)	-0.26	0 100 100	105, 228, 280, 291	0
6	F	87/155 (56%)	-0.33	0 100 100	86, 151, 218, 234	0
7	G	171/179 (95%)	-0.28	0 100 100	120, 178, 253, 300	0
8	H	135/146 (92%)	-0.23	0 100 100	158, 232, 286, 294	0
9	I	114/122 (93%)	0.02	4 (3%) 44 35	137, 224, 283, 293	0
10	J	66/70 (94%)	-0.19	0 100 100	93, 186, 243, 270	0
11	K	115/120 (95%)	-0.20	0 100 100	92, 175, 245, 284	0
12	L	44/70 (62%)	-0.11	1 (2%) 60 51	142, 237, 271, 297	0
13	R	9/9 (100%)	-0.21	0 100 100	201, 226, 298, 299	0
14	S	38/53 (71%)	1.75	16 (42%) 0 0	300, 300, 300, 300	0
15	U	40/53 (75%)	1.24	9 (22%) 0 1	183, 300, 300, 300	0
All	All	4084/4688 (87%)	-0.21	43 (1%) 80 72	62, 189, 282, 300	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
15	U	38	DC	4.3
15	U	39	DC	4.3
14	S	3	DA	4.2
15	U	37	DT	4.2
15	U	34	DG	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
17	MG	A	1804	1/1	0.46	1.44	166,166,166,166	0
17	MG	A	1803	1/1	0.52	0.62	171,171,171,171	0
16	ZN	C	401	1/1	0.86	0.36	174,174,174,174	0
16	ZN	L	101	1/1	0.95	0.10	188,188,188,188	0
16	ZN	I	202	1/1	0.96	0.03	216,216,216,216	0
16	ZN	B	1301	1/1	0.98	0.16	166,166,166,166	0
16	ZN	A	1801	1/1	0.98	0.04	153,153,153,153	0
16	ZN	A	1802	1/1	0.98	0.09	166,166,166,166	0
16	ZN	J	101	1/1	0.99	0.28	172,172,172,172	0
16	ZN	I	201	1/1	0.99	0.15	173,173,173,173	0
16	ZN	C	402	1/1	1.00	0.09	148,148,148,148	0

6.5 Other polymers [i](#)

There are no such residues in this entry.