



Full wwPDB X-ray Structure Validation Report ⓘ

May 31, 2021 – 02:08 PM JST

PDB ID : 7C4S
Title : Sphingosine-1-phosphate receptor 3 with a natural ligand.
Authors : Iwata, S.; Maeda, S.; Luo, F.; Nango, E.; hirata, K.; Asada, H.
Deposited on : 2020-05-18
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.19
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.19


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	K	214	
1	L	214	
2	H	219	
2	J	219	
3	A	378	
3	B	378	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody Fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	214	Total	C	N	O	S	0	0	0
			1658	1033	275	341	9			
1	K	214	Total	C	N	O	S	0	0	0
			1658	1033	275	341	9			

- Molecule 2 is a protein called Antibody Fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	219	Total	C	N	O	S	0	0	0
			1639	1036	263	332	8			
2	J	219	Total	C	N	O	S	0	0	0
			1639	1036	263	332	8			


- Molecule 3 is a protein called Sphingosine 1-phosphate receptor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	282	Total	C	N	O	S	0	0	0
			2243	1494	366	363	20			
3	B	280	Total	C	N	O	S	0	0	0
			2223	1483	361	359	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	15	GLN	ASN	engineered mutation	UNP Q99500
B	15	GLN	ASN	engineered mutation	UNP Q99500

- Molecule 4 is (2S,3R,4E)-2-amino-3-hydroxyoctadec-4-en-1-yl dihydrogen phosphate (three-letter code: S1P) (formula: C₁₈H₃₈NO₅P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	18	1	5	1		
4	B	1	Total	C	N	O	P	0	0
			25	18	1	5	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antibody Fab fragment light chain




- Molecule 1: Antibody Fab fragment light chain




- Molecule 2: Antibody Fab fragment heavy chain




- Molecule 2: Antibody Fab fragment heavy chain




- Molecule 3: Sphingosine 1-phosphate receptor 3





● Molecule 3: Sphingosine 1-phosphate receptor 3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	79.48Å 247.08Å 98.61Å 90.00° 97.37° 90.00°	Depositor
Resolution (Å)	43.48 – 3.20 48.90 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.48-3.20) 100.0 (48.90-3.20)	Depositor EDS
R_{merge}	0.73	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.242 , 0.263 0.245 , 0.267	Depositor DCC
R_{free} test set	1520 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	89.6	Xtriage
Anisotropy	0.534	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	11110	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: S1P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	K	0.37	0/1694	0.60	0/2299
1	L	0.39	1/1694 (0.1%)	0.59	0/2299
2	H	0.36	0/1683	0.64	0/2302
2	J	0.35	0/1683	0.64	0/2302
3	A	0.38	0/2293	0.63	0/3107
3	B	0.37	0/2273	0.64	2/3081 (0.1%)
All	All	0.37	1/11320 (0.0%)	0.63	2/15390 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	202	THR	C-N	-5.33	1.21	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	32	LEU	CA-CB-CG	6.25	129.68	115.30
3	B	266	ASP	CB-CG-OD1	5.27	123.04	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	K	1658	0	1588	13	0
1	L	1658	0	1588	13	0
2	H	1639	0	1591	12	0
2	J	1639	0	1591	12	0
3	A	2243	0	2374	11	0
3	B	2223	0	2355	13	0
4	A	25	0	36	0	0
4	B	25	0	36	0	0
All	All	11110	0	11159	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:102:SER:HA	3:B:29:ALA:HA	1.77	0.67
2:J:6:GLN:O	2:J:111:GLN:NE2	2.37	0.58
1:L:133:VAL:HG22	1:L:178:THR:HG23	1.86	0.57
3:B:148:ALA:HB1	3:B:150:LYS:HD2	1.89	0.55
2:H:6:GLN:O	2:H:111:GLN:NE2	2.41	0.54
1:L:91:SER:HA	1:L:96:LEU:HD22	1.89	0.53
2:H:52:ASP:OD1	3:A:33:LYS:NZ	2.39	0.53
3:A:194:LYS:HD3	3:A:267:VAL:HA	1.91	0.52
2:H:6:GLN:NE2	2:H:94:TYR:O	2.43	0.52
3:A:242:LEU:HA	3:A:245:VAL:HG22	1.93	0.51
1:L:27:GLU:OE2	1:K:157:ASN:N	2.41	0.50
1:K:133:VAL:HG22	1:K:178:THR:HG23	1.93	0.49
1:L:19:VAL:HG12	1:L:75:LEU:HB2	1.94	0.49
1:L:90:GLN:NE2	1:L:93:ASN:O	2.42	0.49
1:K:19:VAL:HG12	1:K:75:LEU:HB2	1.94	0.49
1:K:32:PHE:O	1:K:91:SER:OG	2.31	0.49
1:K:91:SER:HA	1:K:96:LEU:HD22	1.94	0.49
1:L:32:PHE:O	1:L:91:SER:OG	2.30	0.49
2:H:118:SER:OG	2:H:119:SER:N	2.46	0.49
2:J:52:ASP:OD1	3:B:33:LYS:NZ	2.39	0.49
2:J:118:SER:OG	2:J:119:SER:N	2.46	0.49
1:L:103:LYS:HG2	1:L:105:GLU:HG3	1.95	0.48
3:B:23:GLN:HG3	3:B:28:LEU:HD23	1.94	0.48
2:J:30:THR:HB	2:J:54:GLU:HB3	1.95	0.47
3:B:222:VAL:HG11	3:B:242:LEU:HD21	1.95	0.47
1:L:30:GLY:H	1:L:68:ALA:HA	1.80	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.95	0.47
2:H:67:LYS:HE3	2:J:87:THR:HG21	1.98	0.46
1:K:103:LYS:HG2	1:K:105:GLU:HG3	1.98	0.46
3:A:130:LEU:HD22	3:A:207:ILE:HD12	1.98	0.46
3:B:186:SER:HA	3:B:194:LYS:HE2	1.98	0.46
1:K:90:GLN:NE2	1:K:93:ASN:O	2.44	0.45
2:J:6:GLN:NE2	2:J:94:TYR:O	2.49	0.45
2:J:38:LYS:HB2	2:J:48:ILE:HD11	1.98	0.45
3:B:87:LEU:HA	3:B:90:ILE:HG22	1.99	0.45
2:H:102:SER:OG	3:A:23:GLN:O	2.29	0.45
1:K:14:SER:OG	1:K:17:GLU:OE2	2.34	0.45
3:B:209:VAL:HA	3:B:212:VAL:HG12	1.99	0.44
3:B:144:ARG:HG2	3:B:146:TYR:H	1.81	0.43
3:A:194:LYS:HB3	3:A:267:VAL:HG23	2.00	0.43
1:L:13:MET:HE2	1:L:78:VAL:HG21	2.00	0.43
3:A:15:GLN:N	1:K:184:ASP:OD2	2.52	0.43
3:B:132:ILE:HG21	3:B:298:TYR:HE2	1.84	0.43
2:H:102:SER:HA	3:A:29:ALA:HA	2.01	0.43
2:H:85:SER:HB3	2:J:85:SER:HB3	2.00	0.43
2:J:125:PRO:HB3	2:J:151:TYR:HB3	2.00	0.43
1:K:30:GLY:H	1:K:68:ALA:HA	1.83	0.42
1:L:50:GLY:HA2	2:H:104:LEU:HD11	2.02	0.42
2:J:46:GLU:OE2	2:J:63:LYS:NZ	2.42	0.42
3:B:131:ALA:HA	3:B:134:ILE:HG22	2.01	0.42
2:H:30:THR:HB	2:H:54:GLU:HB3	2.01	0.42
2:H:125:PRO:HB3	2:H:151:TYR:HB3	2.00	0.42
1:L:14:SER:OG	1:L:17:GLU:OE2	2.36	0.42
1:L:157:ASN:N	1:K:27:GLU:OE2	2.45	0.41
3:B:285:VAL:HA	3:B:288:VAL:HG22	2.01	0.41
1:L:150:ILE:HD11	1:L:179:LEU:HD21	2.03	0.41
3:A:285:VAL:O	3:A:289:LEU:N	2.47	0.41
1:K:50:GLY:HA2	2:J:104:LEU:HD11	2.02	0.41
3:B:107:SER:HB3	3:B:110:VAL:HG23	2.02	0.41
3:A:107:SER:HB3	3:A:110:VAL:HG23	2.03	0.40
1:K:154:GLU:OE2	1:K:156:GLN:NE2	2.52	0.40
3:A:271:VAL:HG12	3:A:272:GLN:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	212/214 (99%)	204 (96%)	7 (3%)	1 (0%)	29	67
1	L	212/214 (99%)	205 (97%)	6 (3%)	1 (0%)	29	67
2	H	217/219 (99%)	202 (93%)	12 (6%)	3 (1%)	11	46
2	J	217/219 (99%)	202 (93%)	12 (6%)	3 (1%)	11	46
3	A	276/378 (73%)	255 (92%)	20 (7%)	1 (0%)	34	69
3	B	274/378 (72%)	258 (94%)	16 (6%)	0	100	100
All	All	1408/1622 (87%)	1326 (94%)	73 (5%)	9 (1%)	25	64

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	56	GLY
2	J	56	GLY
2	H	55	THR
2	H	155	SER
2	J	55	THR
2	J	155	SER
3	A	240	ALA
1	L	68	ALA
1	K	68	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	190/190 (100%)	190 (100%)	0	100	100
1	L	190/190 (100%)	190 (100%)	0	100	100
2	H	187/187 (100%)	186 (100%)	1 (0%)	88	95
2	J	187/187 (100%)	187 (100%)	0	100	100
3	A	245/329 (74%)	243 (99%)	2 (1%)	81	93
3	B	243/329 (74%)	241 (99%)	2 (1%)	81	93
All	All	1242/1412 (88%)	1237 (100%)	5 (0%)	91	95

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	203	VAL
3	A	32	LEU
3	A	311	ARG
3	B	150	LYS
3	B	223	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	S1P	A	1001	-	23,24,24	0.57	0	23,28,28	1.10	1 (4%)
4	S1P	B	401	-	23,24,24	0.57	0	23,28,28	1.10	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	S1P	A	1001	-	-	18/25/25/25	-
4	S1P	B	401	-	-	15/25/25/25	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	S1P	C3-C4-C5	-2.99	118.12	124.79
4	B	401	S1P	C3-C4-C5	-2.45	119.32	124.79

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1001	S1P	O1-C1-C2-N2
4	A	1001	S1P	O1-C1-C2-C3
4	A	1001	S1P	C1-O1-P22-O24
4	A	1001	S1P	C1-C2-C3-O3
4	A	1001	S1P	C1-C2-C3-C4
4	A	1001	S1P	C2-C3-C4-C5
4	A	1001	S1P	O3-C3-C4-C5
4	B	401	S1P	O1-C1-C2-N2
4	B	401	S1P	O1-C1-C2-C3
4	B	401	S1P	C1-C2-C3-O3
4	B	401	S1P	C1-C2-C3-C4

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	401	S1P	N2-C2-C3-C4
4	B	401	S1P	C2-C3-C4-C5
4	B	401	S1P	O3-C3-C4-C5
4	A	1001	S1P	C5-C6-C7-C8
4	A	1001	S1P	C10-C11-C12-C13
4	B	401	S1P	C5-C6-C7-C8
4	A	1001	S1P	C1-O1-P22-O25
4	A	1001	S1P	C6-C7-C8-C9
4	A	1001	S1P	C14-C15-C16-C17
4	A	1001	S1P	C1-O1-P22-O23
4	A	1001	S1P	C12-C13-C14-C15
4	B	401	S1P	N2-C2-C3-O3
4	B	401	S1P	C13-C14-C15-C16
4	A	1001	S1P	C7-C8-C9-C10
4	B	401	S1P	C12-C13-C14-C15
4	B	401	S1P	C1-O1-P22-O25
4	A	1001	S1P	N2-C2-C3-C4
4	B	401	S1P	C7-C8-C9-C10
4	B	401	S1P	C11-C12-C13-C14
4	A	1001	S1P	N2-C2-C3-O3
4	B	401	S1P	C4-C5-C6-C7
4	A	1001	S1P	C9-C10-C11-C12

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	K	214/214 (100%)	0.13	4 (1%) 66 53	55, 68, 84, 130	0
1	L	214/214 (100%)	0.17	4 (1%) 66 53	53, 66, 83, 122	0
2	H	219/219 (100%)	0.24	11 (5%) 28 16	56, 72, 89, 121	0
2	J	219/219 (100%)	0.29	10 (4%) 32 20	58, 72, 89, 125	0
3	A	282/378 (74%)	0.54	29 (10%) 6 4	65, 90, 146, 178	0
3	B	280/378 (74%)	0.57	31 (11%) 5 3	65, 93, 139, 170	0
All	All	1428/1622 (88%)	0.35	89 (6%) 20 11	53, 74, 127, 178	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	137	HIS	9.7
3	B	152	HIS	5.1
2	H	101	TYR	4.9
1	K	56	THR	4.6
3	A	139	THR	4.6
2	H	1	ASP	4.3
3	A	247	ILE	4.3
2	J	1	ASP	4.2
3	B	247	ILE	4.0
3	B	219	TYR	4.0
3	B	242	LEU	3.8
3	A	242	LEU	3.7
3	A	147	ASP	3.6
3	A	75	TYR	3.5
3	B	14	GLY	3.4
3	A	272	GLN	3.4
1	L	212	ASN	3.2
3	B	239	MET	3.2
3	A	306	ARG	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	220	PHE	3.2
3	A	14	GLY	3.1
3	B	74	MET	3.1
2	H	100	TYR	3.1
3	A	239	MET	3.1
3	A	77	PHE	3.0
1	K	163	TRP	3.0
3	A	135	GLU	3.0
3	A	140	MET	3.0
2	H	180	LEU	3.0
3	A	73	ARG	3.0
3	B	143	MET	2.9
1	L	56	THR	2.9
3	B	218	ILE	2.9
2	H	168	SER	2.9
3	B	65	TRP	2.8
1	L	163	TRP	2.8
3	B	245	VAL	2.8
2	J	101	TYR	2.8
3	B	135	GLU	2.7
2	H	102	SER	2.7
3	B	155	PHE	2.7
2	J	168	SER	2.7
3	B	243	ARG	2.7
3	A	307	ARG	2.6
3	B	241	LEU	2.6
2	J	138	THR	2.6
3	A	298	TYR	2.5
3	B	77	PHE	2.5
3	B	75	TYR	2.5
3	B	244	THR	2.5
3	A	243	ARG	2.5
3	A	74	MET	2.5
3	B	151	ARG	2.5
3	A	131	ALA	2.5
2	J	155	SER	2.5
3	A	25	VAL	2.4
3	A	150	LYS	2.4
3	B	64	ILE	2.4
2	H	175	LEU	2.4
3	B	272	GLN	2.4
3	A	153	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	151	ARG	2.3
3	A	155	PHE	2.3
2	H	176	LEU	2.3
3	B	308	ALA	2.3
1	K	212	ASN	2.3
3	A	273	ALA	2.3
3	B	76	PHE	2.2
2	J	174	ALA	2.2
3	B	273	ALA	2.2
2	H	104	LEU	2.2
2	J	170	HIS	2.2
3	B	132	ILE	2.2
2	H	138	THR	2.2
3	A	236	GLU	2.2
2	J	184	SER	2.2
3	A	256	TRP	2.1
1	K	187	GLU	2.1
3	A	149	ASN	2.1
1	L	57	GLY	2.1
3	B	147	ASP	2.1
2	H	139	GLY	2.1
2	J	169	VAL	2.1
3	B	306	ARG	2.0
3	A	76	PHE	2.0
2	J	176	LEU	2.0
3	A	237	ARG	2.0
3	B	113	LEU	2.0
3	B	146	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	S1P	A	1001	25/25	0.92	0.34	71,78,82,86	0
4	S1P	B	401	25/25	0.92	0.41	75,80,84,87	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.