



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:58 am BST

PDB ID : 4C5A  
Title : The X-ray crystal structures of D-alanyl-D-alanine ligase in complex ADP and D-cycloserine phosphate  
Authors : Batson, S.; Majce, V.; Lloyd, A.J.; Rea, D.; Fishwick, C.W.G.; Simmons, K.J.; Fulop, V.; Roper, D.I.  
Deposited on : 2013-09-10  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>7%</div> </div> </div>
1	B	330	<div> <div>18%</div> <div> <div></div> <div>78%</div> <div>13%</div> <div>8%</div> </div> </div>
2	C	8	<div> <div>63%</div> <div> <div></div> <div>63%</div> <div>38%</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-ALANINE–D-ALANINE LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	0	0
			2305	1464	382	446	13			
1	B	304	Total	C	N	O	S	0	0	0
			2290	1455	380	443	12			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	UNP C4ZRI7
A	-22	SER	-	expression tag	UNP C4ZRI7
A	-21	TYR	-	expression tag	UNP C4ZRI7
A	-20	TYR	-	expression tag	UNP C4ZRI7
A	-19	HIS	-	expression tag	UNP C4ZRI7
A	-18	HIS	-	expression tag	UNP C4ZRI7
A	-17	HIS	-	expression tag	UNP C4ZRI7
A	-16	HIS	-	expression tag	UNP C4ZRI7
A	-15	HIS	-	expression tag	UNP C4ZRI7
A	-14	HIS	-	expression tag	UNP C4ZRI7
A	-13	ASP	-	expression tag	UNP C4ZRI7
A	-12	TYR	-	expression tag	UNP C4ZRI7
A	-11	ILE	-	expression tag	UNP C4ZRI7
A	-10	PRO	-	expression tag	UNP C4ZRI7
A	-9	THR	-	expression tag	UNP C4ZRI7
A	-8	THR	-	expression tag	UNP C4ZRI7
A	-7	GLU	-	expression tag	UNP C4ZRI7
A	-6	ASN	-	expression tag	UNP C4ZRI7
A	-5	LEU	-	expression tag	UNP C4ZRI7
A	-4	TYR	-	expression tag	UNP C4ZRI7
A	-3	PHE	-	expression tag	UNP C4ZRI7
A	-2	ASN	-	expression tag	UNP C4ZRI7
A	-1	GLY	-	expression tag	UNP C4ZRI7
A	0	ALA	-	expression tag	UNP C4ZRI7
B	-23	MET	-	expression tag	UNP C4ZRI7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	SER	-	expression tag	UNP C4ZRI7
B	-21	TYR	-	expression tag	UNP C4ZRI7
B	-20	TYR	-	expression tag	UNP C4ZRI7
B	-19	HIS	-	expression tag	UNP C4ZRI7
B	-18	HIS	-	expression tag	UNP C4ZRI7
B	-17	HIS	-	expression tag	UNP C4ZRI7
B	-16	HIS	-	expression tag	UNP C4ZRI7
B	-15	HIS	-	expression tag	UNP C4ZRI7
B	-14	HIS	-	expression tag	UNP C4ZRI7
B	-13	ASP	-	expression tag	UNP C4ZRI7
B	-12	TYR	-	expression tag	UNP C4ZRI7
B	-11	ILE	-	expression tag	UNP C4ZRI7
B	-10	PRO	-	expression tag	UNP C4ZRI7
B	-9	THR	-	expression tag	UNP C4ZRI7
B	-8	THR	-	expression tag	UNP C4ZRI7
B	-7	GLU	-	expression tag	UNP C4ZRI7
B	-6	ASN	-	expression tag	UNP C4ZRI7
B	-5	LEU	-	expression tag	UNP C4ZRI7
B	-4	TYR	-	expression tag	UNP C4ZRI7
B	-3	PHE	-	expression tag	UNP C4ZRI7
B	-2	ASN	-	expression tag	UNP C4ZRI7
B	-1	GLY	-	expression tag	UNP C4ZRI7
B	0	ALA	-	expression tag	UNP C4ZRI7

- Molecule 2 is a protein called PEPTIDE.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	0	0	0
			66	43	10	13			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is [(4R)-4-azanyl-4,5-dihydro-1,2-oxazol-3-yl] dihydrogen phosphate (three-letter code: DS0) (formula: C<sub>3</sub>H<sub>7</sub>N<sub>2</sub>O<sub>5</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			11	3	2	5	1		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			11	3	2	5	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Mg	0	0
			2	2		
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	239	Total	O	0	0
			239	239		
7	B	101	Total	O	0	0
			101	101		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	7	Total	O	0	0
			7	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: D-ALANINE-D-ALANINE LIGASE



#### • Molecule 1: D-ALANINE-D-ALANINE LIGASE



#### • Molecule 2: PEPTIDE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.79 Å 97.51 Å 109.99 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.76 – 1.65 48.32 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.76-1.65) 99.7 (48.32-1.65)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 1.65 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.205 , 0.237 0.234 , 0.258	Depositor DCC
$R_{free}$ test set	2861 reflections (4.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.7136e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, DS0, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.78	0/2346	0.84	3/3175 (0.1%)
1	B	0.58	0/2331	0.66	1/3155 (0.0%)
2	C	0.68	0/67	0.54	0/89
All	All	0.68	0/4744	0.75	4/6419 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	LEU	CB-CG-CD2	5.63	120.57	111.00
1	A	255	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	A	248	LEU	CB-CG-CD1	5.48	120.31	111.00
1	B	98	LEU	CB-CG-CD2	5.29	120.00	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2305	0	2319	28	0
1	B	2290	0	2302	41	0
2	C	66	0	59	5	0
3	A	27	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	27	0	12	0	0
4	A	11	0	5	0	0
4	B	11	0	5	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	12	0	16	2	0
7	A	239	0	0	7	0
7	B	101	0	0	3	0
7	C	7	0	0	0	0
All	All	5100	0	4730	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:LYS:HE3	1:A:180:GLU:OE2	1.62	0.99
1:A:8:LEU:CD2	1:A:58:VAL:HG13	1.91	0.99
1:A:103:LEU:HD12	1:A:247:THR:CG2	1.94	0.96
1:A:103:LEU:HD12	1:A:247:THR:HG22	1.50	0.93
1:A:8:LEU:HD23	1:A:58:VAL:HG13	1.51	0.92
1:B:62:LEU:HD22	7:B:2002:HOH:O	1.76	0.85
1:A:8:LEU:HD23	1:A:58:VAL:CG1	2.07	0.84
1:A:120:ALA:O	1:A:124:LYS:HG2	1.84	0.78
1:A:8:LEU:HD21	1:A:58:VAL:HG13	1.68	0.74
1:B:5:ILE:HD12	1:B:57:LYS:HB2	1.71	0.73
1:A:2:THR:HG23	2:C:8:ALA:HB1	1.69	0.73
1:A:103:LEU:CD1	1:A:247:THR:CG2	2.66	0.73
7:A:2041:HOH:O	2:C:4:TYR:HB3	1.89	0.72
1:A:103:LEU:CD1	1:A:247:THR:HG22	2.21	0.71
1:B:101:LYS:CE	1:B:180:GLU:OE2	2.40	0.70
1:B:103:LEU:HD21	1:B:247:THR:HG22	1.74	0.69
1:A:5:ILE:HD12	1:A:57:LYS:HB2	1.76	0.68
1:A:101:LYS:CE	1:A:180:GLU:OE2	2.39	0.68
1:A:197:ILE:HD11	1:A:242:LEU:HB2	1.75	0.67
1:A:8:LEU:CD2	1:A:58:VAL:CG1	2.68	0.65
1:B:59:PHE:CE2	1:B:298:VAL:HG13	2.32	0.64
1:B:101:LYS:HE3	1:B:180:GLU:OE2	1.97	0.64
1:A:2:THR:HB	7:A:2006:HOH:O	1.98	0.63
1:B:5:ILE:CD1	1:B:57:LYS:HB2	2.29	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ILE:HD11	1:B:299:VAL:HG22	1.82	0.62
1:B:103:LEU:CD2	1:B:248:LEU:HD12	2.30	0.61
1:B:278:THR:OG1	1:B:279:SER:N	2.34	0.60
1:B:101:LYS:HE2	1:B:180:GLU:OE2	2.02	0.59
1:A:173:HIS:HD2	7:A:2074:HOH:O	1.85	0.59
1:A:251:LYS:HD3	7:A:2047:HOH:O	2.03	0.58
1:B:101:LYS:HG3	7:B:2042:HOH:O	2.03	0.58
1:B:101:LYS:HG2	7:B:2045:HOH:O	2.04	0.57
6:A:1307:GOL:H32	7:A:2239:HOH:O	2.03	0.57
1:B:72:LEU:CD1	1:B:76:LEU:HD11	2.35	0.57
1:B:104:TRP:NE1	1:B:248:LEU:HD13	2.20	0.56
1:B:142:ILE:HD12	1:B:182:TRP:HZ3	1.70	0.56
1:B:72:LEU:HD11	1:B:76:LEU:HD11	1.86	0.55
1:B:295:SER:O	1:B:299:VAL:HG23	2.07	0.55
1:B:26:VAL:HG13	1:B:298:VAL:HG11	1.89	0.55
1:A:8:LEU:HD22	1:A:8:LEU:N	2.23	0.54
1:B:104:TRP:HE1	1:B:248:LEU:HD13	1.73	0.52
1:A:101:LYS:NZ	1:A:111:VAL:CG1	2.73	0.52
1:B:47:VAL:O	1:B:50:LEU:HG	2.10	0.51
1:B:279:SER:C	1:B:281:SER:H	2.13	0.50
1:B:142:ILE:HD12	1:B:182:TRP:CZ3	2.47	0.50
1:B:279:SER:C	1:B:281:SER:N	2.65	0.50
1:B:72:LEU:CD1	1:B:76:LEU:CD1	2.90	0.49
1:B:103:LEU:CD2	1:B:247:THR:HG22	2.42	0.48
1:A:1:MET:HB2	2:C:8:ALA:O	2.14	0.47
1:B:72:LEU:HD13	1:B:76:LEU:CD1	2.44	0.47
1:B:253:TRP:CH2	1:B:275:PRO:HD2	2.50	0.47
1:B:30:LEU:CD2	1:B:35:ILE:HD13	2.44	0.47
7:A:2041:HOH:O	2:C:4:TYR:CB	2.57	0.47
1:B:30:LEU:HD21	1:B:298:VAL:HG12	1.97	0.46
1:B:119:ARG:NH2	1:B:175:GLU:OE2	2.39	0.45
1:A:101:LYS:HZ3	1:A:111:VAL:CG1	2.29	0.45
1:A:289:GLN:NE2	7:A:2229:HOH:O	2.45	0.44
1:B:75:MET:HE2	1:B:76:LEU:HD23	1.99	0.44
1:A:114:TRP:CD1	1:A:178:LEU:HD22	2.53	0.44
1:A:229:LEU:HG	1:A:233:GLN:HB3	1.99	0.44
1:A:168:ARG:NH2	6:A:1308:GOL:O2	2.52	0.42
1:B:72:LEU:HD13	1:B:76:LEU:HG	1.99	0.42
1:B:150:SER:HB3	1:B:216:TYR:HE2	1.84	0.42
1:A:1:MET:N	2:C:7:GLY:O	2.45	0.42
1:B:197:ILE:HD11	1:B:242:LEU:HB2	2.02	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:LYS:HB3	1:B:154:MET:HG2	2.01	0.42
1:B:30:LEU:HG	1:B:298:VAL:HG11	2.01	0.42
1:B:81:LEU:HA	1:B:82:PRO:HD3	1.91	0.41
1:B:284:PRO:HB3	1:B:294:PHE:HD2	1.85	0.41
1:B:288:ARG:HA	1:B:292:MET:O	2.22	0.40
1:A:5:ILE:CD1	1:A:57:LYS:HB2	2.48	0.40
1:B:78:LEU:HA	1:B:78:LEU:HD12	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/330 (92%)	302 (99%)	2 (1%)	0	100	100
1	B	302/330 (92%)	296 (98%)	4 (1%)	2 (1%)	22	6
2	C	6/8 (75%)	6 (100%)	0	0	100	100
All	All	612/668 (92%)	604 (99%)	6 (1%)	2 (0%)	41	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	278	THR
1	B	280	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/266 (92%)	237 (97%)	7 (3%)	42	16
1	B	242/266 (91%)	235 (97%)	7 (3%)	42	16
2	C	6/6 (100%)	6 (100%)	0	100	100
All	All	492/538 (91%)	478 (97%)	14 (3%)	43	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	LEU
1	A	95	MET
1	A	103	LEU
1	A	109	LEU
1	A	229	LEU
1	A	237	LEU
1	A	248	LEU
1	B	56	GLN
1	B	78	LEU
1	B	98	LEU
1	B	103	LEU
1	B	237	LEU
1	B	288	ARG
1	B	306	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	HIS
1	A	289	GLN
1	B	172	GLN
1	B	280	HIS
1	B	289	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DS0	B	311	5	10,11,11	4.28	2 (20%)	8,16,16	2.61	2 (25%)
6	GOL	A	1307	-	5,5,5	0.39	0	5,5,5	0.40	0
3	ADP	B	310	5	24,29,29	1.03	1 (4%)	29,45,45	1.12	2 (6%)
3	ADP	A	310	5	24,29,29	1.21	2 (8%)	29,45,45	1.39	3 (10%)
6	GOL	A	1308	-	5,5,5	0.32	0	5,5,5	0.89	0
4	DS0	A	311	5	10,11,11	3.44	4 (40%)	8,16,16	3.20	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DS0	B	311	5	-	0/3/15/15	0/1/1/1
6	GOL	A	1307	-	-	2/4/4/4	-
3	ADP	B	310	5	-	3/12/32/32	0/3/3/3
3	ADP	A	310	5	-	3/12/32/32	0/3/3/3
6	GOL	A	1308	-	-	2/4/4/4	-
4	DS0	A	311	5	-	0/3/15/15	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	311	DS0	O1-N2	-12.94	1.27	1.43
4	A	311	DS0	O1-N2	-9.12	1.32	1.43
4	A	311	DS0	P8-O7	3.46	1.64	1.59
4	A	311	DS0	C3-N2	3.41	1.32	1.29
4	B	311	DS0	C3-N2	3.25	1.32	1.29
3	A	310	ADP	C2-N3	2.90	1.36	1.32
4	A	311	DS0	O1-C5	-2.80	1.41	1.45
3	A	310	ADP	O4'-C1'	2.80	1.45	1.41
3	B	310	ADP	C5-C4	2.79	1.48	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	311	DS0	C5-O1-N2	7.63	115.61	108.07
4	B	311	DS0	C5-O1-N2	5.81	113.81	108.07
3	A	310	ADP	N3-C2-N1	-4.05	122.35	128.68
4	B	311	DS0	O7-P8-O9	-3.36	97.37	109.32
4	A	311	DS0	O1-C5-C4	-3.33	100.33	105.53
3	A	310	ADP	C2-N1-C6	3.03	123.94	118.75
3	B	310	ADP	N3-C2-N1	-2.75	124.39	128.68
4	A	311	DS0	O7-P8-O9	-2.47	100.55	109.32
3	B	310	ADP	N6-C6-N1	2.37	123.49	118.57
3	A	310	ADP	N6-C6-N1	2.12	122.97	118.57

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	310	ADP	PA-O3A-PB-O2B
6	A	1308	GOL	C1-C2-C3-O3
3	A	310	ADP	PA-O3A-PB-O3B
6	A	1308	GOL	O2-C2-C3-O3
3	B	310	ADP	PA-O3A-PB-O1B
3	B	310	ADP	PA-O3A-PB-O2B
3	B	310	ADP	PA-O3A-PB-O3B
6	A	1307	GOL	O1-C1-C2-C3
3	A	310	ADP	PA-O3A-PB-O1B
6	A	1307	GOL	O1-C1-C2-O2


There are no ring outliers.


2 monomers are involved in 2 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1307	GOL	1	0
6	A	1308	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	306/330 (92%)	0.33	5 (1%) 72 75	14, 20, 37, 59	0
1	B	304/330 (92%)	1.07	61 (20%) 1 1	16, 40, 62, 94	0
2	C	8/8 (100%)	3.79	5 (62%) 0 0	31, 37, 50, 51	0
All	All	618/668 (92%)	0.74	71 (11%) 4 4	14, 28, 58, 94	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	10.8
2	C	8	ALA	9.2
2	C	7	GLY	8.4
1	B	81	LEU	5.8
1	B	280	HIS	5.5
1	B	128	ASP	5.2
1	B	279	SER	5.2
1	B	226	PRO	4.9
1	B	306	ASP	4.7
1	B	35	ILE	4.7
1	B	50	LEU	4.7
1	B	54	GLY	4.5
1	B	39	PRO	4.5
1	B	5	ILE	4.5
1	B	55	PHE	4.4
1	B	28	ALA	4.3
1	B	126	LEU	4.2
1	B	12	THR	4.0
2	C	6	GLN	4.0
1	B	52	SER	4.0
1	B	127	SER	4.0
1	B	13	SER	3.7
1	B	125	GLY	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	75	MET	3.4
1	B	53	MET	3.4
1	B	82	PRO	3.4
1	B	298	VAL	3.3
1	A	128	ASP	3.3
1	B	36	ASP	3.3
1	B	42	PRO	3.3
1	B	219	ASP	3.3
1	B	282	LEU	3.3
2	C	1	GLU	3.3
1	B	122	PHE	3.2
1	B	129	LYS	3.1
1	B	79	MET	3.1
1	B	34	GLY	3.1
1	B	66	GLY	3.1
1	B	14	ALA	2.9
1	B	209	PHE	2.8
1	B	30	LEU	2.8
1	A	306	ASP	2.8
1	B	302	LEU	2.8
1	B	31	ARG	2.7
1	B	278	THR	2.7
1	B	6	ALA	2.7
1	B	78	LEU	2.7
1	B	56	GLN	2.7
1	A	2	THR	2.6
1	B	33	GLY	2.6
1	B	4	LYS	2.5
1	B	59	PHE	2.5
1	B	130	GLN	2.5
1	B	217	LEU	2.4
1	B	195	GLU	2.4
1	B	8	LEU	2.3
1	B	23	GLY	2.3
1	B	21	ASN	2.3
1	B	38	TYR	2.3
1	B	22	SER	2.3
1	B	114	TRP	2.2
1	B	158	VAL	2.2
1	B	26	VAL	2.2
1	B	17	GLU	2.2
1	B	157	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	292	MET	2.1
2	C	4	TYR	2.1
1	B	297	LEU	2.1
1	B	206	SER	2.0
1	B	295	SER	2.0
1	A	18	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	GOL	A	1308	6/6	0.74	0.13	41,42,45,46	0
6	GOL	A	1307	6/6	0.86	0.20	45,48,50,51	0
4	DS0	B	311	11/11	0.93	0.17	31,36,39,40	0
3	ADP	B	310	27/27	0.96	0.08	28,30,31,32	0
5	MG	B	331	1/1	0.97	0.05	28,28,28,28	0
4	DS0	A	311	11/11	0.97	0.11	14,17,23,28	0
3	ADP	A	310	27/27	0.98	0.11	13,15,17,18	0
5	MG	B	330	1/1	0.98	0.09	28,28,28,28	0
5	MG	A	331	1/1	0.98	0.12	14,14,14,14	0
5	MG	A	330	1/1	0.98	0.10	13,13,13,13	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ADP B 310:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP A 310:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.