



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 10:29 pm BST

PDB ID : 4C6G  
Title : Structural Investigations into the Stereochemistry and Activity of a Phenylalanine-2,3-Aminomutase from *Taxus chinensis*  
Authors : Wybenga, G.G.; Szymanski, W.; Wu, B.; Feringa, B.L.; Janssen, D.B.; Dijkstra, B.W.  
Deposited on : 2013-09-18  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	<div> <div>4%</div> <div>80% 8% • 12%</div> </div>
1	B	707	<div> <div>4%</div> <div>78% 10% • 11%</div> </div>
1	C	707	<div> <div>3%</div> <div>78% 10% • 12%</div> </div>
1	D	707	<div> <div>2%</div> <div>77% 11% • 11%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLALANINE AMMONIA-LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	624	Total	C	N	O	S	0	0	0
			4833	3074	826	911	22			
1	B	626	Total	C	N	O	S	0	0	0
			4848	3085	828	913	22			
1	C	625	Total	C	N	O	S	0	0	0
			4839	3078	825	914	22			
1	D	626	Total	C	N	O	S	0	0	0
			4847	3083	828	914	22			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q68G84
A	-18	GLY	-	expression tag	UNP Q68G84
A	-17	SER	-	expression tag	UNP Q68G84
A	-16	SER	-	expression tag	UNP Q68G84
A	-15	HIS	-	expression tag	UNP Q68G84
A	-14	HIS	-	expression tag	UNP Q68G84
A	-13	HIS	-	expression tag	UNP Q68G84
A	-12	HIS	-	expression tag	UNP Q68G84
A	-11	HIS	-	expression tag	UNP Q68G84
A	-10	HIS	-	expression tag	UNP Q68G84
A	-9	SER	-	expression tag	UNP Q68G84
A	-8	SER	-	expression tag	UNP Q68G84
A	-7	GLY	-	expression tag	UNP Q68G84
A	-6	LEU	-	expression tag	UNP Q68G84
A	-5	VAL	-	expression tag	UNP Q68G84
A	-4	PRO	-	expression tag	UNP Q68G84
A	-3	ARG	-	expression tag	UNP Q68G84
A	-2	GLY	-	expression tag	UNP Q68G84
A	-1	SER	-	expression tag	UNP Q68G84
A	0	HIS	-	expression tag	UNP Q68G84
A	231	ALA	ASN	engineered mutation	UNP Q68G84

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	expression tag	UNP Q68G84
B	-18	GLY	-	expression tag	UNP Q68G84
B	-17	SER	-	expression tag	UNP Q68G84
B	-16	SER	-	expression tag	UNP Q68G84
B	-15	HIS	-	expression tag	UNP Q68G84
B	-14	HIS	-	expression tag	UNP Q68G84
B	-13	HIS	-	expression tag	UNP Q68G84
B	-12	HIS	-	expression tag	UNP Q68G84
B	-11	HIS	-	expression tag	UNP Q68G84
B	-10	HIS	-	expression tag	UNP Q68G84
B	-9	SER	-	expression tag	UNP Q68G84
B	-8	SER	-	expression tag	UNP Q68G84
B	-7	GLY	-	expression tag	UNP Q68G84
B	-6	LEU	-	expression tag	UNP Q68G84
B	-5	VAL	-	expression tag	UNP Q68G84
B	-4	PRO	-	expression tag	UNP Q68G84
B	-3	ARG	-	expression tag	UNP Q68G84
B	-2	GLY	-	expression tag	UNP Q68G84
B	-1	SER	-	expression tag	UNP Q68G84
B	0	HIS	-	expression tag	UNP Q68G84
B	231	ALA	ASN	engineered mutation	UNP Q68G84
C	-19	MET	-	expression tag	UNP Q68G84
C	-18	GLY	-	expression tag	UNP Q68G84
C	-17	SER	-	expression tag	UNP Q68G84
C	-16	SER	-	expression tag	UNP Q68G84
C	-15	HIS	-	expression tag	UNP Q68G84
C	-14	HIS	-	expression tag	UNP Q68G84
C	-13	HIS	-	expression tag	UNP Q68G84
C	-12	HIS	-	expression tag	UNP Q68G84
C	-11	HIS	-	expression tag	UNP Q68G84
C	-10	HIS	-	expression tag	UNP Q68G84
C	-9	SER	-	expression tag	UNP Q68G84
C	-8	SER	-	expression tag	UNP Q68G84
C	-7	GLY	-	expression tag	UNP Q68G84
C	-6	LEU	-	expression tag	UNP Q68G84
C	-5	VAL	-	expression tag	UNP Q68G84
C	-4	PRO	-	expression tag	UNP Q68G84
C	-3	ARG	-	expression tag	UNP Q68G84
C	-2	GLY	-	expression tag	UNP Q68G84
C	-1	SER	-	expression tag	UNP Q68G84
C	0	HIS	-	expression tag	UNP Q68G84
C	231	ALA	ASN	engineered mutation	UNP Q68G84

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	expression tag	UNP Q68G84
D	-18	GLY	-	expression tag	UNP Q68G84
D	-17	SER	-	expression tag	UNP Q68G84
D	-16	SER	-	expression tag	UNP Q68G84
D	-15	HIS	-	expression tag	UNP Q68G84
D	-14	HIS	-	expression tag	UNP Q68G84
D	-13	HIS	-	expression tag	UNP Q68G84
D	-12	HIS	-	expression tag	UNP Q68G84
D	-11	HIS	-	expression tag	UNP Q68G84
D	-10	HIS	-	expression tag	UNP Q68G84
D	-9	SER	-	expression tag	UNP Q68G84
D	-8	SER	-	expression tag	UNP Q68G84
D	-7	GLY	-	expression tag	UNP Q68G84
D	-6	LEU	-	expression tag	UNP Q68G84
D	-5	VAL	-	expression tag	UNP Q68G84
D	-4	PRO	-	expression tag	UNP Q68G84
D	-3	ARG	-	expression tag	UNP Q68G84
D	-2	GLY	-	expression tag	UNP Q68G84
D	-1	SER	-	expression tag	UNP Q68G84
D	0	HIS	-	expression tag	UNP Q68G84
D	231	ALA	ASN	engineered mutation	UNP Q68G84


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	73	Total O 73 73	0	0
2	B	42	Total O 42 42	0	0
2	C	60	Total O 60 60	0	0
2	D	45	Total O 45 45	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: PHENYLALANINE AMMONIA-LYASE



#### • Molecule 1: PHENYLALANINE AMMONIA-LYASE



Chain C:  3% 78% 10% 12%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.96Å 146.57Å 99.33Å 90.00° 100.02° 90.00°	Depositor
Resolution (Å)	48.73 – 2.10 48.08 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.73-2.10) 99.7 (48.08-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.63 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.201 , 0.241 0.203 , 0.242	Depositor DCC
$R_{free}$ test set	8119 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	19587	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	5/4916 (0.1%)	0.68	8/6671 (0.1%)
1	B	0.60	4/4932 (0.1%)	0.65	4/6692 (0.1%)
1	C	0.64	7/4921 (0.1%)	0.76	2/6678 (0.0%)
1	D	0.63	6/4930 (0.1%)	0.65	4/6691 (0.1%)
All	All	0.62	22/19699 (0.1%)	0.68	18/26732 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	71	ARG	CZ-NH2	20.26	1.59	1.33
1	D	77	ALA	C-N	-19.91	0.88	1.34
1	A	77	ALA	C-N	-19.16	0.90	1.34
1	B	77	ALA	C-N	-18.35	0.91	1.34
1	C	71	ARG	CZ-NH1	-15.19	1.13	1.33

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	71	ARG	NE-CZ-NH2	-31.23	104.69	120.30
1	C	71	ARG	NE-CZ-NH1	14.93	127.77	120.30
1	A	71	ARG	NE-CZ-NH2	-14.50	113.05	120.30
1	B	77	ALA	O-C-N	-11.49	104.32	122.70
1	D	78	ASP	O-C-N	-11.29	104.63	122.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	71	ARG	Sidechain
1	D	77	ALA	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4833	0	4901	29	0
1	B	4848	0	4913	47	0
1	C	4839	0	4911	50	0
1	D	4847	0	4917	42	0
2	A	73	0	0	0	0
2	B	42	0	0	0	0
2	C	60	0	0	0	0
2	D	45	0	0	0	0
All	All	19587	0	19642	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:TYR:HB3	1:B:318:LYS:HG2	1.58	0.85
1:C:80:TYR:CB	1:D:318:LYS:HG2	2.09	0.81
1:C:304:LEU:HD11	1:C:619:ARG:HD3	1.64	0.80
1:A:80:TYR:CB	1:B:318:LYS:HG2	2.11	0.80
1:C:80:TYR:HB3	1:D:318:LYS:HG2	1.67	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	612/707 (87%)	596 (97%)	16 (3%)	0	100	100
1	B	614/707 (87%)	592 (96%)	22 (4%)	0	100	100
1	C	613/707 (87%)	598 (98%)	15 (2%)	0	100	100
1	D	614/707 (87%)	597 (97%)	17 (3%)	0	100	100
All	All	2453/2828 (87%)	2383 (97%)	70 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	527/592 (89%)	507 (96%)	20 (4%)	33	34
1	B	528/592 (89%)	514 (97%)	14 (3%)	44	48
1	C	528/592 (89%)	508 (96%)	20 (4%)	33	34
1	D	529/592 (89%)	509 (96%)	20 (4%)	33	34
All	All	2112/2368 (89%)	2038 (96%)	74 (4%)	36	38

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	624	LYS
1	C	275	LEU
1	D	575	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	59	ARG
1	C	135	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	557	GLN
1	A	567	HIS
1	B	557	GLN
1	C	557	GLN
1	D	557	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	2
1	A	2
1	D	2

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	78:ASP	C	79:ILE	N	1.10
1	B	78:ASP	C	79:ILE	N	1.05
1	D	78:ASP	C	79:ILE	N	1.03
1	B	77:ALA	C	78:ASP	N	0.91
1	A	77:ALA	C	78:ASP	N	0.90

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	624/707 (88%)	0.17	30 (4%) 30 36	27, 43, 66, 99	0
1	B	626/707 (88%)	0.15	25 (3%) 38 44	26, 45, 69, 91	0
1	C	625/707 (88%)	0.12	20 (3%) 47 54	27, 43, 68, 95	0
1	D	626/707 (88%)	0.18	16 (2%) 56 61	27, 43, 64, 81	0
All	All	2501/2828 (88%)	0.16	91 (3%) 42 49	26, 44, 67, 99	0

The worst 5 of 91 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	314	LEU	6.5
1	C	314	LEU	5.4
1	D	314	LEU	4.6
1	B	99	GLU	3.8
1	D	576	THR	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.