



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:34 am BST

PDB ID : 4C6R  
Title : Crystal structure of the TIR domain from the Arabidopsis Thaliana disease resistance protein RPS4  
Authors : Williams, S.J.; Sohn, K.H.; Wan, L.; Bernoux, M.; Ma, Y.; Segonzac, C.; Ve, T.; Sarris, P.; Ericsson, D.J.; Saucet, S.B.; Zhang, X.; Parker, J.; Dodds, P.N.; Jones, J.D.G.; Kobe, B.  
Deposited on : 2013-09-19  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>
1	B	171	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>
1	C	171	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>5%</div> </div> </div>
1	D	171	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>5%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DISEASE RESISTANCE PROTEIN RPS4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1317	844	230	239	4			
1	B	162	Total	C	N	O	S	0	0	0
			1317	844	230	239	4			
1	C	162	Total	C	N	O	S	0	0	0
			1317	844	230	239	4			
1	D	162	Total	C	N	O	S	0	1	0
			1322	847	230	241	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	SER	-	expression tag	UNP Q9XGM3
A	9	ASN	-	expression tag	UNP Q9XGM3
A	10	ALA	-	expression tag	UNP Q9XGM3
B	8	SER	-	expression tag	UNP Q9XGM3
B	9	ASN	-	expression tag	UNP Q9XGM3
B	10	ALA	-	expression tag	UNP Q9XGM3
C	8	SER	-	expression tag	UNP Q9XGM3
C	9	ASN	-	expression tag	UNP Q9XGM3
C	10	ALA	-	expression tag	UNP Q9XGM3
D	8	SER	-	expression tag	UNP Q9XGM3
D	9	ASN	-	expression tag	UNP Q9XGM3
D	10	ALA	-	expression tag	UNP Q9XGM3

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	61	Total	O	0	0
			61	61		
2	B	74	Total	O	0	0
			74	74		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	111	Total 111	O 111	0	0
2	D	97	Total 97	O 97	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DISEASE RESISTANCE PROTEIN RPS4

Chain A: 




- Molecule 1: DISEASE RESISTANCE PROTEIN RPS4

Chain B: 




- Molecule 1: DISEASE RESISTANCE PROTEIN RPS4

Chain C: 



- Molecule 1: DISEASE RESISTANCE PROTEIN RPS4

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.92Å 78.64Å 80.67Å 65.63° 78.64° 78.93°	Depositor
Resolution (Å)	19.60 – 2.05 19.60 – 2.05	Depositor EDS
% Data completeness (in resolution range)	94.1 (19.60-2.05) 94.1 (19.60-2.05)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.10 (at 2.06Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.186 , 0.230 0.194 , 0.234	Depositor DCC
$R_{free}$ test set	1879 reflections (4.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.8	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.048 for h,h-k,h-l 0.025 for -h,-l,-k 0.024 for -h,-h+l,-h+k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5616	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/1338	0.53	0/1797
1	B	0.38	0/1338	0.55	1/1797 (0.1%)
1	C	0.41	0/1338	0.58	0/1797
1	D	0.41	0/1346	0.54	0/1808
All	All	0.40	0/5360	0.55	1/7199 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	LEU	CA-CB-CG	5.77	128.56	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1317	0	1366	12	0
1	B	1317	0	1366	14	0
1	C	1317	0	1366	14	0
1	D	1322	0	1370	10	0
2	A	61	0	0	1	0
2	B	74	0	0	3	0
2	C	111	0	0	2	0
2	D	97	0	0	0	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5616	0	5468	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:137:LYS:HA	1:C:140:LYS:HE2	1.71	0.71
1:C:63:LYS:NZ	1:C:67:GLU:OE2	2.27	0.68
1:C:92:ILE:O	1:C:96:THR:HG23	1.95	0.66
1:C:26:ASP:OD2	2:C:2007:HOH:O	2.13	0.66
1:B:166:GLU:HG3	1:B:169:LYS:HE2	1.81	0.62
1:C:45:ASN:ND2	2:C:2002:HOH:O	2.32	0.62
1:B:146:ILE:HG12	1:B:149:ILE:HG12	1.86	0.58
1:B:51:TYR:CE1	1:B:52:GLU:HG3	2.39	0.56
1:C:51:TYR:CE2	1:C:52:GLU:HG3	2.42	0.55
1:A:89:LEU:HD11	1:A:146:ILE:HG13	1.89	0.54
1:A:71:VAL:HG11	1:A:92:ILE:HD12	1.91	0.53
1:A:66:GLU:HG3	1:A:91:LYS:HE3	1.91	0.52
1:D:134:GLU:CD	1:D:134:GLU:H	2.12	0.52
1:B:66:GLU:HG3	1:B:91:LYS:HE3	1.92	0.52
1:D:63:LYS:O	1:D:67:GLU:HG2	2.10	0.51
1:A:133:ASP:OD2	1:A:135:ARG:NH2	2.40	0.50
1:D:28:ARG:HG2	1:D:51:TYR:CZ	2.47	0.49
1:A:88:GLU:O	1:A:92:ILE:HG12	2.13	0.49
1:B:146:ILE:HG13	1:B:148:ASN:OD1	2.13	0.47
1:D:63:LYS:NZ	1:D:67:GLU:OE1	2.48	0.47
1:D:121:LYS:HD2	1:D:121:LYS:HA	1.72	0.47
1:A:51:TYR:CE1	1:A:52:GLU:HG3	2.50	0.47
1:B:20:ILE:O	1:B:49:ASP:HB2	2.16	0.45
1:C:16:HIS:HE1	1:C:64:ARG:O	1.99	0.45
1:D:93:LYS:HB2	1:D:146:ILE:HD11	1.99	0.45
1:B:138:LYS:O	2:B:2054:HOH:O	2.21	0.44
1:A:71:VAL:HG11	1:A:92:ILE:CD1	2.48	0.44
1:A:51:TYR:CD1	1:A:52:GLU:HG3	2.53	0.44
1:A:58:LEU:O	1:A:62:LEU:HG	2.18	0.44
1:B:71:VAL:HG23	1:B:101:LEU:HD11	1.99	0.43
2:B:2021:HOH:O	1:D:172:LYS:HE2	2.18	0.43
1:B:116:ARG:NH1	2:B:2046:HOH:O	2.51	0.43
1:B:156:LYS:HG3	1:B:157:LYS:HG3	2.01	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:ARG:HG2	1:B:51:TYR:CZ	2.53	0.43
1:B:69:LYS:HA	1:B:69:LYS:HD2	1.83	0.43
1:D:58:LEU:O	1:D:62:LEU:HG	2.19	0.43
1:A:20:ILE:O	1:A:49:ASP:HB2	2.19	0.42
1:B:166:GLU:O	1:B:169:LYS:HG2	2.20	0.42
1:C:96:THR:HG21	1:C:103:ALA:HB2	2.01	0.42
1:D:51:TYR:CD1	1:D:52:GLU:HG3	2.55	0.42
1:A:34:HIS:HD2	2:A:2008:HOH:O	2.02	0.42
1:B:51:TYR:CD1	1:B:52:GLU:HG3	2.54	0.42
1:C:165:ASN:O	1:C:169:LYS:HG3	2.20	0.41
1:C:66:GLU:HG3	1:C:91:LYS:CE	2.51	0.41
1:A:81:GLU:HG3	1:A:125:ARG:HD3	2.02	0.41
1:C:66:GLU:HG3	1:C:91:LYS:HE3	2.04	0.40
1:C:47:PHE:CE2	1:C:64:ARG:HG3	2.56	0.40
1:D:121:LYS:HG3	1:D:125:ARG:NE	2.36	0.40
1:C:163:LYS:HA	1:C:163:LYS:HD3	1.98	0.40
1:C:20:ILE:O	1:C:49:ASP:HB2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:2012:HOH:O	2:D:2082:HOH:O[1_455]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/171 (94%)	159 (99%)	1 (1%)	0	100	100
1	B	160/171 (94%)	159 (99%)	1 (1%)	0	100	100
1	C	160/171 (94%)	159 (99%)	1 (1%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	161/171 (94%)	160 (99%)	1 (1%)	0	100	100
All	All	641/684 (94%)	637 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/153 (95%)	146 (100%)	0	100	100
1	B	146/153 (95%)	145 (99%)	1 (1%)	84	84
1	C	146/153 (95%)	146 (100%)	0	100	100
1	D	147/153 (96%)	147 (100%)	0	100	100
All	All	585/612 (96%)	584 (100%)	1 (0%)	93	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS
1	B	16	HIS
1	B	165	ASN
1	C	16	HIS
1	C	45	ASN
1	D	16	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/171 (94%)	0.03	6 (3%)	41	45	11, 26, 47, 64	0
1	B	162/171 (94%)	0.17	8 (4%)	29	31	11, 25, 48, 62	0
1	C	162/171 (94%)	-0.04	3 (1%)	66	71	11, 21, 38, 62	0
1	D	162/171 (94%)	-0.14	3 (1%)	66	71	11, 20, 34, 53	0
All	All	648/684 (94%)	0.00	20 (3%)	49	53	11, 23, 44, 64	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	99	GLY	3.7
1	C	15	GLN	3.7
1	B	134	GLU	3.4
1	D	15	GLN	3.3
1	B	100	THR	2.7
1	A	134	GLU	2.6
1	A	59	ASP	2.6
1	A	98	GLU	2.6
1	B	59	ASP	2.4
1	D	59	ASP	2.4
1	D	176	THR	2.2
1	B	84	TRP	2.2
1	A	15	GLN	2.2
1	C	176	THR	2.2
1	B	15	GLN	2.1
1	B	149	ILE	2.1
1	B	148	ASN	2.1
1	A	132	GLY	2.0
1	C	99	GLY	2.0
1	A	99	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.