



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 05:15 am BST

PDB ID : 1C7G  
Title : TYROSINE PHENOL-LYASE FROM ERWINIA HERBICOLA  
Authors : Mikami, B.; Yamamoto, Y.; Katayama, T.; Suzuki, H.  
Deposited on : 2000-02-18  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	456	76% 21% .
1	B	456	70% 26% .
1	C	456	75% 21% .
1	D	456	73% 23% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLP	A	1000	-	X	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLP	B	1000	-	X	-	-
2	PLP	C	1000	-	X	-	-
2	PLP	D	1000	-	X	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 14905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYROSINE PHENOL-LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	B	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	C	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	D	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	77	Total	O	0	0
			77	77		
3	C	158	Total	O	0	0
			158	158		
3	D	87	Total	O	0	0
			87	87		





• Molecule 1: TYROSINE PHENOL-LYASE

Chain D: 73% 23% .



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	163.49Å 113.04Å 101.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.10	Depositor
% Data completeness (in resolution range)	78.6 (8.00-2.10)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
Refinement program	X-PLOR 3.85	Depositor
R, $R_{free}$	0.186 , 0.242	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14905	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/3678	0.60	1/4955 (0.0%)
1	B	0.35	0/3678	0.59	1/4955 (0.0%)
1	C	0.36	0/3678	0.62	1/4955 (0.0%)
1	D	0.35	0/3678	0.59	1/4955 (0.0%)
All	All	0.35	0/14712	0.60	4/19820 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ASP	N-CA-C	-7.04	91.98	111.00
1	C	214	ASP	N-CA-C	-6.72	92.86	111.00
1	D	214	ASP	N-CA-C	-6.03	94.71	111.00
1	B	214	ASP	N-CA-C	-5.65	95.75	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	223	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3605	0	3548	67	0
1	B	3605	0	3548	87	1
1	C	3605	0	3548	70	0
1	D	3605	0	3548	73	0
2	A	15	0	6	1	0
2	B	15	0	6	1	0
2	C	15	0	6	2	0
2	D	15	0	6	1	0
3	A	103	0	0	2	0
3	B	77	0	0	3	0
3	C	158	0	0	3	0
3	D	87	0	0	0	0
All	All	14905	0	14216	289	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 289 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:MET:HE3	1:C:368:ALA:HA	1.43	1.01
1:C:97:HIS:HA	1:C:294:LEU:HD13	1.59	0.83
1:B:362:PRO:HG2	1:B:401:GLU:HG2	1.61	0.83
1:A:374:THR:OG1	1:A:376:VAL:HG13	1.79	0.82
1:B:374:THR:OG1	1:B:376:VAL:HG13	1.79	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:ASP:OD1	1:B:22:ASP:OD1[2_575]	1.80	0.40

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/456 (100%)	437 (96%)	17 (4%)	0	100	100
1	B	454/456 (100%)	433 (95%)	20 (4%)	1 (0%)	47	49
1	C	454/456 (100%)	438 (96%)	16 (4%)	0	100	100
1	D	454/456 (100%)	433 (95%)	21 (5%)	0	100	100
All	All	1816/1824 (100%)	1741 (96%)	74 (4%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	391	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/380 (100%)	334 (88%)	46 (12%)	5	2
1	B	380/380 (100%)	333 (88%)	47 (12%)	4	2
1	C	380/380 (100%)	339 (89%)	41 (11%)	6	3
1	D	380/380 (100%)	338 (89%)	42 (11%)	6	3
All	All	1520/1520 (100%)	1344 (88%)	176 (12%)	5	3

5 of 176 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	388	ARG
1	C	48	LEU
1	D	318	ARG
1	B	391	GLU
1	B	446	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	ASN
1	B	130	GLN
1	C	130	GLN
1	D	130	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PLP	B	1000	1	15,15,16	3.64	5 (33%)	20,22,23	4.08	14 (70%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PLP	C	1000	1	15,15,16	3.64	6 (40%)	20,22,23	4.01	14 (70%)
2	PLP	A	1000	1	15,15,16	3.69	5 (33%)	20,22,23	4.09	14 (70%)
2	PLP	D	1000	1	15,15,16	3.44	6 (40%)	20,22,23	4.00	14 (70%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	1000	1	-	4/6/6/8	0/1/1/1
2	PLP	C	1000	1	-	4/6/6/8	0/1/1/1
2	PLP	A	1000	1	-	4/6/6/8	0/1/1/1
2	PLP	D	1000	1	-	4/6/6/8	0/1/1/1

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	PLP	C5-C4	12.50	1.54	1.40
2	C	1000	PLP	C5-C4	12.14	1.53	1.40
2	B	1000	PLP	C5-C4	12.04	1.53	1.40
2	D	1000	PLP	C5-C4	11.04	1.52	1.40
2	B	1000	PLP	C3-C2	4.45	1.45	1.40

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	PLP	O3P-P-O4P	-9.54	81.34	106.73
2	B	1000	PLP	O3P-P-O4P	-9.50	81.45	106.73
2	C	1000	PLP	O3P-P-O4P	-9.31	81.96	106.73
2	D	1000	PLP	O3P-P-O4P	-9.23	82.16	106.73
2	D	1000	PLP	O3P-P-O2P	-6.58	82.49	107.64

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1000	PLP	C4-C5-C5A-O4P
2	B	1000	PLP	C6-C5-C5A-O4P
2	B	1000	PLP	C5A-O4P-P-O2P

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	1000	PLP	C5A-O4P-P-O3P
2	C	1000	PLP	C4-C5-C5A-O4P

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1000	PLP	1	0
2	C	1000	PLP	2	0
2	A	1000	PLP	1	0
2	D	1000	PLP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.