



wwPDB EM Validation Summary Report ⓘ

Nov 19, 2022 – 04:23 PM EST

PDB ID : 3C91
EMDB ID : EMD-1733
Title : Thermoplasma acidophilum 20S proteasome with an open gate
Authors : Rabl, J.; Smith, D.M.; Yu, Y.; Chang, S.C.; Goldberg, A.L.; Cheng, Y.
Deposited on : 2008-02-14
Resolution : 6.80 Å(reported)
Based on initial model : 1PMA

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>26%</div> <div>42%</div> <div>47%</div> <div>7%</div> <div>..</div> </div>
1	B	233	<div> <div>28%</div> <div>45%</div> <div>45%</div> <div>7%</div> <div>..</div> </div>
1	C	233	<div> <div>27%</div> <div>43%</div> <div>45%</div> <div>8%</div> <div>..</div> </div>
1	D	233	<div> <div>30%</div> <div>42%</div> <div>47%</div> <div>8%</div> <div>..</div> </div>
1	E	233	<div> <div>32%</div> <div>42%</div> <div>47%</div> <div>8%</div> <div>..</div> </div>
1	F	233	<div> <div>35%</div> <div>42%</div> <div>47%</div> <div>7%</div> <div>..</div> </div>
1	G	233	<div> <div>31%</div> <div>41%</div> <div>49%</div> <div>7%</div> <div>..</div> </div>
1	O	233	<div> <div>30%</div> <div>43%</div> <div>46%</div> <div>7%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	P	233	
1	Q	233	
1	R	233	
1	S	233	
1	T	233	
1	U	233	
2	1	203	
2	2	203	
2	H	203	
2	I	203	
2	J	203	
2	K	203	
2	L	203	
2	M	203	
2	N	203	
2	V	203	
2	W	203	
2	X	203	
2	Y	203	
2	Z	203	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 46578 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	B	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	C	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	D	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	E	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	F	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	G	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	O	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	P	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	Q	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	R	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	S	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	T	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		
1	U	227	Total	C	N	O	S	0	0
			1769	1123	299	344	3		


- Molecule 2 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	Trace
2	I	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	J	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	K	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	L	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	M	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	N	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	V	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	W	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	X	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	Y	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	Z	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	1	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		
2	2	203	Total	C	N	O	S	0	0
			1558	985	264	298	11		




• Molecule 1: Proteasome subunit alpha






• Molecule 1: Proteasome subunit alpha






• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha







• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha





• Molecule 1: Proteasome subunit alpha







• Molecule 1: Proteasome subunit alpha




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta







• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta







• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D7	Depositor
Number of particles used	44794	Depositor
Resolution determination method	Not provided	
CTF correction method	PHASE AND AMPLITUDE	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	20.00	Depositor
Minimum defocus (nm)	-1500.00	Depositor
Maximum defocus (nm)	-3500.00	Depositor
Magnification	51159	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	9.018	Depositor
Minimum map value	-4.980	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	2.5	Depositor
Map size (\AA)	197.028, 197.028, 197.028	wwPDB
Map dimensions	144, 144, 144	wwPDB
Map angles ($^\circ$)	90, 90, 90	wwPDB
Pixel spacing (\AA)	1.36825, 1.36825, 1.36825	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1793	0.80	3/2416 (0.1%)
1	B	0.55	0/1793	0.80	3/2416 (0.1%)
1	C	0.55	0/1793	0.80	3/2416 (0.1%)
1	D	0.56	0/1793	0.80	3/2416 (0.1%)
1	E	0.55	0/1793	0.80	3/2416 (0.1%)
1	F	0.55	0/1793	0.80	3/2416 (0.1%)
1	G	0.55	0/1793	0.80	3/2416 (0.1%)
1	O	0.55	0/1793	0.80	3/2416 (0.1%)
1	P	0.55	0/1793	0.80	3/2416 (0.1%)
1	Q	0.56	0/1793	0.80	3/2416 (0.1%)
1	R	0.56	0/1793	0.80	3/2416 (0.1%)
1	S	0.55	0/1793	0.80	3/2416 (0.1%)
1	T	0.55	0/1793	0.80	3/2416 (0.1%)
1	U	0.56	0/1793	0.80	3/2416 (0.1%)
2	1	0.55	0/1577	0.75	1/2129 (0.0%)
2	2	0.54	0/1577	0.75	0/2129
2	H	0.57	0/1577	0.76	0/2129
2	I	0.57	0/1577	0.76	0/2129
2	J	0.54	0/1577	0.75	0/2129
2	K	0.55	0/1577	0.77	2/2129 (0.1%)
2	L	0.57	0/1577	0.77	0/2129
2	M	0.55	0/1577	0.75	1/2129 (0.0%)
2	N	0.54	0/1577	0.75	0/2129
2	V	0.57	0/1577	0.76	0/2129
2	W	0.57	0/1577	0.76	0/2129
2	X	0.54	0/1577	0.75	0/2129
2	Y	0.55	0/1577	0.77	2/2129 (0.1%)
2	Z	0.57	0/1577	0.77	0/2129
All	All	0.56	0/47180	0.78	48/63630 (0.1%)

There are no bond length outliers.

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	13	THR	C-N-CA	-11.65	92.58	121.70
1	S	13	THR	C-N-CA	-11.65	92.58	121.70
1	B	13	THR	C-N-CA	-11.63	92.62	121.70
1	P	13	THR	C-N-CA	-11.63	92.62	121.70
1	A	13	THR	C-N-CA	-11.63	92.63	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1769	0	1799	278	0
1	B	1769	0	1799	263	0
1	C	1769	0	1798	285	0
1	D	1769	0	1799	299	0
1	E	1769	0	1798	308	0
1	F	1769	0	1797	302	0
1	G	1769	0	1799	333	0
1	O	1769	0	1799	281	0
1	P	1769	0	1799	266	0
1	Q	1769	0	1798	280	0
1	R	1769	0	1799	299	0
1	S	1769	0	1798	308	0
1	T	1769	0	1797	306	0
1	U	1769	0	1799	333	0
2	1	1558	0	1609	130	0
2	2	1558	0	1609	123	0
2	H	1558	0	1609	144	0
2	I	1558	0	1609	96	0
2	J	1558	0	1609	109	0
2	K	1558	0	1609	124	0
2	L	1558	0	1609	134	0
2	M	1558	0	1609	131	0
2	N	1558	0	1609	123	0
2	V	1558	0	1609	145	0
2	W	1558	0	1609	101	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	X	1558	0	1609	98	0
2	Y	1558	0	1609	127	0
2	Z	1558	0	1609	130	0
All	All	46578	0	47704	4081	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 4081 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:177:GLU:HA	1:D:58:LEU:CG	1.26	1.65
1:T:12:ILE:CD1	1:T:23:GLN:CG	1.74	1.64
1:O:177:GLU:HA	1:P:58:LEU:CD1	1.21	1.64
1:S:12:ILE:CD1	1:S:23:GLN:CG	1.74	1.64
1:R:12:ILE:CD1	1:R:23:GLN:CG	1.74	1.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	B	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	C	225/233 (97%)	188 (84%)	27 (12%)	10 (4%)	2	22
1	D	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	E	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	F	225/233 (97%)	188 (84%)	27 (12%)	10 (4%)	2	22
1	G	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	P	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	Q	225/233 (97%)	188 (84%)	27 (12%)	10 (4%)	2	22
1	R	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	S	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
1	T	225/233 (97%)	188 (84%)	27 (12%)	10 (4%)	2	22
1	U	225/233 (97%)	189 (84%)	26 (12%)	10 (4%)	2	22
2	1	201/203 (99%)	176 (88%)	23 (11%)	2 (1%)	15	54
2	2	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	H	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	I	201/203 (99%)	183 (91%)	16 (8%)	2 (1%)	15	54
2	J	201/203 (99%)	180 (90%)	20 (10%)	1 (0%)	29	69
2	K	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	L	201/203 (99%)	181 (90%)	16 (8%)	4 (2%)	7	38
2	M	201/203 (99%)	176 (88%)	23 (11%)	2 (1%)	15	54
2	N	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	V	201/203 (99%)	185 (92%)	14 (7%)	2 (1%)	15	54
2	W	201/203 (99%)	183 (91%)	16 (8%)	2 (1%)	15	54
2	X	201/203 (99%)	180 (90%)	20 (10%)	1 (0%)	29	69
2	Y	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	15	54
2	Z	201/203 (99%)	181 (90%)	16 (8%)	4 (2%)	7	38
All	All	5964/6104 (98%)	5187 (87%)	607 (10%)	170 (3%)	7	29

5 of 170 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	VAL
1	A	128	GLY
1	A	182	GLU
1	A	200	SER
1	B	14	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	B	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	C	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	D	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	E	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	F	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	G	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	O	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	P	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	Q	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	R	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	S	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	T	188/193 (97%)	166 (88%)	22 (12%)	5	20
1	U	188/193 (97%)	166 (88%)	22 (12%)	5	20
2	1	170/170 (100%)	152 (89%)	18 (11%)	6	23
2	2	170/170 (100%)	146 (86%)	24 (14%)	3	16
2	H	170/170 (100%)	145 (85%)	25 (15%)	3	15
2	I	170/170 (100%)	148 (87%)	22 (13%)	4	18
2	J	170/170 (100%)	143 (84%)	27 (16%)	2	13
2	K	170/170 (100%)	143 (84%)	27 (16%)	2	13
2	L	170/170 (100%)	146 (86%)	24 (14%)	3	16
2	M	170/170 (100%)	152 (89%)	18 (11%)	6	23
2	N	170/170 (100%)	146 (86%)	24 (14%)	3	16
2	V	170/170 (100%)	145 (85%)	25 (15%)	3	15
2	W	170/170 (100%)	149 (88%)	21 (12%)	4	19
2	X	170/170 (100%)	143 (84%)	27 (16%)	2	13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	Y	170/170 (100%)	143 (84%)	27 (16%)	2	13
2	Z	170/170 (100%)	146 (86%)	24 (14%)	3	16
All	All	5012/5082 (99%)	4371 (87%)	641 (13%)	7	18

5 of 641 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	U	136	LEU
2	Y	178	ILE
2	V	7	THR
1	U	134	VAL
2	W	185	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 134 such sidechains are listed below:

Mol	Chain	Res	Type
2	W	164	GLN
2	X	164	GLN
2	1	164	GLN
2	I	164	GLN
2	I	73	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-1733. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 72



Y Index: 72




Z Index: 72


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 90



Y Index: 53




Z Index: 58


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 2.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 290 nm³; this corresponds to an approximate mass of 262 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.147 Å⁻¹


8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-1733 and PDB model 3C91. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 2.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.5).



























































9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 51% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (2.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5062	 0.1730
1	 0.5246	 0.1790
2	 0.5305	 0.1810
A	 0.4983	 0.1720
B	 0.4977	 0.1710
C	 0.4890	 0.1700
D	 0.4896	 0.1680
E	 0.4856	 0.1710
F	 0.4758	 0.1620
G	 0.4810	 0.1680
H	 0.5344	 0.1820
I	 0.5311	 0.1810
J	 0.5285	 0.1840
K	 0.5337	 0.1810
L	 0.5337	 0.1810
M	 0.5403	 0.1840
N	 0.5390	 0.1830
O	 0.4844	 0.1640
P	 0.4839	 0.1660
Q	 0.4821	 0.1660
R	 0.4913	 0.1660
S	 0.4775	 0.1650
T	 0.4787	 0.1590
U	 0.4971	 0.1670
V	 0.5213	 0.1770
W	 0.5219	 0.1720
X	 0.5226	 0.1740
Y	 0.5187	 0.1760
Z	 0.5187	 0.1770

