



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 02:41 am BST

PDB ID : 2CAG
Title : CATALASE COMPOUND II
Authors : Gouet, P.; Jouve, H.M.; Hajdu, J.
Deposited on : 1996-06-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	 75% 18% 5% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATALASE COMPOUND II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	475	Total	C	N	O	S	20	0	0
			3861	2447	684	715	15			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	OMT	MET	CONFLICT	UNP P42321

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	70	Total	O	0	0
			70	70		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CATALASE COMPOUND II



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	111.84Å 111.84Å 249.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.70 14.87 – 2.70	Depositor EDS
% Data completeness (in resolution range)	68.0 (15.00-2.70) 67.6 (14.87-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.69Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.179 , 0.241 0.163 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	43.8	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3974	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, OMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/3961	1.60	64/5359 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	371	ARG	NE-CZ-NH1	14.22	127.41	120.30
1	A	440	ARG	NE-CZ-NH1	10.19	125.39	120.30
1	A	413	ARG	NE-CZ-NH1	9.78	125.19	120.30
1	A	371	ARG	NE-CZ-NH2	-9.32	115.64	120.30
1	A	256	TRP	CD1-CG-CD2	8.92	113.44	106.30
1	A	156	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	A	282	TRP	CD1-CG-CD2	8.63	113.21	106.30
1	A	437	MET	CG-SD-CE	-8.36	86.83	100.20
1	A	342	ARG	NE-CZ-NH2	-8.33	116.14	120.30
1	A	162	TRP	CD1-CG-CD2	7.83	112.57	106.30
1	A	366	ARG	NE-CZ-NH1	7.75	124.17	120.30
1	A	256	TRP	CE2-CD2-CG	-7.51	101.29	107.30
1	A	210	TRP	CD1-CG-CD2	7.43	112.25	106.30
1	A	35	TRP	CE2-CD2-CG	-7.31	101.45	107.30
1	A	440	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	A	35	TRP	CD1-CG-CD2	7.24	112.09	106.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	TRP	CE2-CD2-CG	-7.20	101.54	107.30
1	A	162	TRP	CE2-CD2-CG	-7.17	101.56	107.30
1	A	122	TRP	CE2-CD2-CG	-7.08	101.63	107.30
1	A	189	ARG	NE-CZ-NH1	7.01	123.81	120.30
1	A	413	ARG	NE-CZ-NH2	-6.99	116.81	120.30
1	A	153	THR	CA-CB-CG2	6.99	122.18	112.40
1	A	216	ARG	NE-CZ-NH2	-6.85	116.87	120.30
1	A	135	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	A	410	TRP	CE2-CD2-CG	-6.81	101.85	107.30
1	A	88	MET	CA-CB-CG	6.79	124.84	113.30
1	A	410	TRP	CD1-CG-CD2	6.56	111.55	106.30
1	A	259	GLN	CA-CB-CG	6.49	127.69	113.40
1	A	122	TRP	CD1-CG-CD2	6.45	111.46	106.30
1	A	109	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	A	52	ARG	NE-CZ-NH2	-6.37	117.11	120.30
1	A	342	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	125	VAL	CG1-CB-CG2	-6.30	100.81	110.90
1	A	253	TYR	CB-CG-CD2	-6.26	117.24	121.00
1	A	153	THR	CA-CB-OG1	-6.24	95.90	109.00
1	A	273	TYR	CB-CG-CD2	-6.24	117.26	121.00
1	A	242	ARG	NE-CZ-NH1	6.21	123.41	120.30
1	A	210	TRP	CE2-CD2-CG	-6.18	102.36	107.30
1	A	156	ARG	NE-CZ-NH1	6.17	123.39	120.30
1	A	152	ARG	CA-CB-CG	6.16	126.96	113.40
1	A	423	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	A	52	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	208	ARG	NE-CZ-NH1	5.97	123.28	120.30
1	A	210	TRP	CG-CD2-CE3	5.87	139.18	133.90
1	A	131	VAL	N-CA-CB	-5.80	98.75	111.50
1	A	287	TYR	CB-CG-CD1	-5.77	117.54	121.00
1	A	467	TYR	CB-CG-CD2	-5.76	117.54	121.00
1	A	333	ARG	NE-CZ-NH1	5.71	123.16	120.30
1	A	256	TRP	CG-CD1-NE1	-5.61	104.49	110.10
1	A	294	TYR	CB-CG-CD2	-5.58	117.65	121.00
1	A	216	ARG	NE-CZ-NH1	5.57	123.09	120.30
1	A	88	MET	CG-SD-CE	-5.55	91.32	100.20
1	A	122	TRP	CG-CD2-CE3	5.49	138.84	133.90
1	A	216	ARG	CA-CB-CG	-5.48	101.34	113.40
1	A	35	TRP	CG-CD2-CE3	5.48	138.83	133.90
1	A	208	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	A	122	TRP	CB-CG-CD1	-5.33	120.07	127.00
1	A	153	THR	N-CA-CB	-5.30	100.22	110.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	271	VAL	CG1-CB-CG2	5.26	119.31	110.90
1	A	455	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	182	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	A	33	ASP	CB-CG-OD1	5.21	122.98	118.30
1	A	271	VAL	N-CA-CB	-5.16	100.14	111.50
1	A	147	VAL	N-CA-CB	-5.00	100.49	111.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	359	CYS	Peptide
1	A	382	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3861	0	3678	40	0
2	A	43	0	30	0	0
3	A	70	0	0	1	0
All	All	3974	0	3708	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:348:ASN:HD21	1:A:368:GLY:HA3	1.11	1.10
1:A:348:ASN:ND2	1:A:368:GLY:HA3	1.89	0.88
1:A:331:GLN:O	1:A:334:LEU:HB2	1.94	0.67
1:A:153:THR:HG22	1:A:155:MET:H	1.60	0.65
1:A:132:PHE:HB3	1:A:278:LEU:HD13	1.82	0.60
1:A:60:ALA:HA	1:A:301:PRO:HD3	1.85	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464:HIS:HD2	1:A:466:GLU:H	1.51	0.57
1:A:310:GLN:HB3	1:A:352:ILE:HD13	1.87	0.56
1:A:239:SER:HB2	3:A:511:HOH:O	2.05	0.56
1:A:423:ARG:HD3	1:A:463:VAL:O	2.06	0.55
1:A:146:ILE:HD12	1:A:147:VAL:HG23	1.90	0.54
1:A:53:OMT:HE3	1:A:144:ASN:HB2	1.90	0.53
1:A:189:ARG:HD3	1:A:243:ASP:OD2	2.09	0.53
1:A:208:ARG:NH2	1:A:400:PRO:O	2.42	0.52
1:A:398:GLU:HB2	1:A:399:PRO:HD2	1.91	0.51
1:A:278:LEU:HD12	1:A:329:MET:HG3	1.92	0.51
1:A:198:THR:OG1	1:A:212:LYS:HE2	2.12	0.50
1:A:210:TRP:CE3	1:A:210:TRP:HA	2.46	0.49
1:A:65:THR:HB	1:A:294:TYR:CE1	2.48	0.49
1:A:56:LYS:HE3	1:A:101:ALA:O	2.14	0.47
1:A:248:ILE:HD12	1:A:300:ASN:ND2	2.30	0.46
1:A:437:MET:O	1:A:441:ILE:HG13	2.15	0.46
1:A:192:HIS:CD2	1:A:221:ILE:HD11	2.51	0.46
1:A:329:MET:O	1:A:333:ARG:HG3	2.16	0.46
1:A:429:LEU:HD23	1:A:437:MET:CE	2.45	0.46
1:A:66:VAL:HG21	1:A:79:PHE:HA	1.98	0.45
1:A:215:PHE:CZ	1:A:258:LEU:HD13	2.51	0.45
1:A:330:LEU:O	1:A:334:LEU:HD13	2.16	0.44
1:A:67:THR:OG1	1:A:293:GLY:HA2	2.18	0.44
1:A:23:ALA:O	1:A:27:GLY:HA3	2.18	0.43
1:A:328:LYS:O	1:A:331:GLN:HB2	2.18	0.43
1:A:246:GLU:O	1:A:250:ARG:HB2	2.19	0.43
1:A:380:ILE:HG23	1:A:382:TYR:CE1	2.53	0.43
1:A:128:ASN:HA	1:A:192:HIS:O	2.19	0.42
1:A:366:ARG:HB3	1:A:367:ASP:H	1.75	0.42
1:A:210:TRP:HE3	1:A:210:TRP:HA	1.85	0.42
1:A:436:ARG:HH11	1:A:436:ARG:HD3	1.69	0.41
1:A:371:ARG:HD3	1:A:371:ARG:HA	1.80	0.41
1:A:456:GLN:HG3	1:A:460:PHE:CE2	2.56	0.41
1:A:254:PRO:HG2	1:A:297:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/484 (98%)	438 (93%)	31 (7%)	3 (1%)	25	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	360	PRO
1	A	359	CYS
1	A	366	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/414 (98%)	373 (92%)	34 (8%)	11	25

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	37	LEU
1	A	52	ARG
1	A	87	GLU
1	A	88	MET
1	A	113	LEU
1	A	131	VAL
1	A	153	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	172	LEU
1	A	182	ARG
1	A	191	VAL
1	A	206	ASN
1	A	227	ASP
1	A	237	ARG
1	A	239	SER
1	A	244	LEU
1	A	255	ARG
1	A	259	GLN
1	A	262	ILE
1	A	267	GLU
1	A	271	VAL
1	A	278	LEU
1	A	283	PRO
1	A	297	LEU
1	A	300	ASN
1	A	318	ILE
1	A	330	LEU
1	A	359	CYS
1	A	360	PRO
1	A	363	ASN
1	A	366	ARG
1	A	429	LEU
1	A	435	GLN
1	A	467	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	144	ASN
1	A	145	HIS
1	A	218	GLN
1	A	300	ASN
1	A	317	ASN
1	A	348	ASN
1	A	350	HIS
1	A	363	ASN
1	A	464	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OMT	A	53	1	8,9,10	2.85	3 (37%)	6,12,14	2.91	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OMT	A	53	1	-	4/7/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	OMT	CG-SD	-6.19	1.70	1.78
1	A	53	OMT	CE-SD	-4.35	1.59	1.75
1	A	53	OMT	CB-CG	-2.26	1.50	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	OMT	OD2-SD-CG	4.83	111.72	108.34
1	A	53	OMT	OD2-SD-OD1	-4.37	107.75	117.09
1	A	53	OMT	OD1-SD-CG	-2.03	106.92	108.34

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	53	OMT	CB-CG-SD-OD2
1	A	53	OMT	CB-CG-SD-CE
1	A	53	OMT	CB-CG-SD-OD1
1	A	53	OMT	CA-CB-CG-SD

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	53	OMT	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	485	1,3	27,50,50	2.04	7 (25%)	17,82,82	6.76	15 (88%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	485	1,3	-	0/6/54/54	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	485	HEM	C3B-CAB	-5.05	1.37	1.47
2	A	485	HEM	C3B-C2B	-4.74	1.33	1.40
2	A	485	HEM	C3C-CAC	-4.59	1.38	1.47
2	A	485	HEM	C3C-C2C	-3.46	1.35	1.40
2	A	485	HEM	CBC-CAC	2.47	1.45	1.29
2	A	485	HEM	CBB-CAB	2.20	1.43	1.29
2	A	485	HEM	C1A-CHA	-2.18	1.34	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	485	HEM	CAA-CBA-CGA	13.62	135.52	112.67
2	A	485	HEM	C1D-C2D-C3D	-11.93	98.69	107.00
2	A	485	HEM	C4A-C3A-C2A	-10.27	99.85	107.00
2	A	485	HEM	C4C-C3C-C2C	-10.00	99.91	106.90
2	A	485	HEM	CBD-CAD-C3D	7.43	126.17	112.48
2	A	485	HEM	CMD-C2D-C3D	6.20	136.64	124.94
2	A	485	HEM	CMA-C3A-C4A	5.11	136.31	128.46
2	A	485	HEM	CAD-C3D-C2D	-5.09	112.61	127.25
2	A	485	HEM	C3C-C4C-NC	4.83	120.06	110.94
2	A	485	HEM	C3B-C4B-NB	-4.82	102.98	109.21
2	A	485	HEM	CMC-C2C-C3C	-4.04	117.11	124.68
2	A	485	HEM	CMB-C2B-C3B	-3.33	118.45	124.68
2	A	485	HEM	CAD-CBD-CGD	2.79	117.36	112.67
2	A	485	HEM	CBA-CAA-C2A	2.71	117.49	112.49
2	A	485	HEM	CMD-C2D-C1D	-2.47	124.67	128.46

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/484 (97%)	-1.29	1 (0%) 95 96	11, 26, 50, 84	6 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	PRO	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OMT	A	53	10/11	0.99	0.06	9,19,30,31	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEM	A	485	43/43	0.99	0.07	2,18,29,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.