



# Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 01:50 am BST

PDB ID : 2CDF  
Title : Structure and binding kinetics of three different human CD1d-alpha- Galactosylceramide-specific T cell receptors (TCR 5E)  
Authors : Gadola, S.D.; Koch, M.; Marles-Wright, J.; Lissin, N.M.; Sheperd, D.; Matulis, G.; Harlos, K.; Villiger, P.M.; Stuart, D.I.; Jakobsen, B.K.; Cerundolo, V.; Jones, E.Y.  
Deposited on : 2006-01-23  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	<div> <div>9%</div> <div>81%</div> <div>16%</div> <div>.</div> </div>
2	B	244	<div> <div>8%</div> <div>77%</div> <div>20%</div> <div>..</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TCR 5E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1481	921	253	300	7			

- Molecule 2 is a protein called TCR 5E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1955	1229	335	383	8			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	127	Total	O	0	0
			127	127		
3	B	157	Total	O	0	0
			157	157		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TCR 5E



- Molecule 2: TCR 5E



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.45Å 64.45Å 184.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.25 28.56 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.7 (30.00-2.25) 98.7 (28.56-2.25)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.56 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.188 , 0.269 0.205 , 0.285	Depositor DCC
$R_{free}$ test set	1098 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.2	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 36.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3720	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	1/1510 (0.1%)	0.69	1/2045 (0.0%)
2	B	0.60	0/2008	0.72	2/2729 (0.1%)
All	All	0.62	1/3518 (0.0%)	0.70	3/4774 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	90	CYS	CB-SG	-6.65	1.71	1.82

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	178	LEU	CA-CB-CG	6.29	129.77	115.30
2	B	194	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	A	126	LEU	CA-CB-CG	5.45	127.83	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1481	0	1413	20	0
2	B	1955	0	1844	33	0
3	A	127	0	0	4	0
3	B	157	0	0	9	0
All	All	3720	0	3257	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:40:PRO:HA	3:B:2026:HOH:O	1.79	0.80
1:A:42:GLU:HG2	3:B:2025:HOH:O	1.82	0.78
2:B:14:ILE:HD11	2:B:114:VAL:HG22	1.70	0.74
2:B:44:LEU:HD21	2:B:91:LEU:HD12	1.69	0.74
1:A:109:GLN:OE1	3:A:2076:HOH:O	2.10	0.69
2:B:3:ALA:HB1	2:B:6:TYR:OH	1.92	0.69
1:A:39:GLU:OE2	3:A:2039:HOH:O	2.11	0.68
1:A:182:LYS:HB3	1:A:183:SER:CA	2.26	0.65
1:A:65:GLN:HG2	3:A:2059:HOH:O	1.97	0.65
2:B:163:ASN:HD21	2:B:207:ASN:HD22	1.45	0.64
2:B:36:TYR:HB3	2:B:44:LEU:HD22	1.80	0.62
1:A:182:LYS:HB3	1:A:183:SER:HA	1.83	0.61
2:B:84:PRO:HG3	2:B:116:GLU:HG3	1.83	0.59
2:B:121:VAL:O	2:B:228:ARG:NH2	2.38	0.57
2:B:230:LYS:HG2	2:B:232:VAL:HG13	1.87	0.56
1:A:145:GLN:HE21	1:A:145:GLN:H	1.54	0.55
2:B:136:ILE:HG23	2:B:199:ALA:HB1	1.87	0.55
2:B:208:HIS:HD2	3:B:2093:HOH:O	1.89	0.54
1:A:3:ASN:HB3	3:A:2003:HOH:O	2.07	0.54
2:B:228:ARG:NH1	3:B:2144:HOH:O	2.40	0.53
2:B:98:ARG:HB2	2:B:103:LYS:HD2	1.91	0.52
2:B:71:ARG:HA	3:B:2054:HOH:O	2.09	0.52
2:B:163:ASN:HD21	2:B:207:ASN:ND2	2.09	0.51
2:B:178:LEU:HD22	2:B:190:ALA:HB3	1.92	0.50
2:B:224:TRP:HB2	2:B:230:LYS:HD2	1.94	0.50
1:A:151:SER:HA	1:A:158:ILE:HD12	1.93	0.50
1:A:182:LYS:HB3	1:A:183:SER:CB	2.42	0.49
1:A:15:GLN:NE2	1:A:113:TRP:HB2	2.28	0.48
2:B:25:SER:HB2	2:B:73:GLU:OE1	2.14	0.48
1:A:31:SER:OG	1:A:95:ARG:HG3	2.15	0.47
2:B:139:THR:HG23	2:B:141:LYS:HB2	1.97	0.46
2:B:135:GLU:O	2:B:139:THR:HG22	2.14	0.46
2:B:84:PRO:CG	2:B:116:GLU:HG3	2.45	0.46
1:A:152:LYS:HB2	1:A:152:LYS:NZ	2.31	0.46
1:A:175:SER:OG	2:B:194:ARG:HD3	2.16	0.45
1:A:66:PHE:CE2	1:A:70:ARG:HG2	2.52	0.45
1:A:91:ALA:HB1	1:A:102:LEU:HG	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:181:GLN:HG2	3:B:2119:HOH:O	2.17	0.44
1:A:181:ASN:O	1:A:182:LYS:HG2	2.17	0.44
2:B:39:ASP:HB3	2:B:40:PRO:HD2	2.00	0.44
2:B:222:ASP:O	2:B:230:LYS:NZ	2.42	0.43
2:B:3:ALA:HB3	2:B:27:THR:OG1	2.18	0.43
2:B:208:HIS:CD2	3:B:2093:HOH:O	2.69	0.42
2:B:61:LEU:HG	3:B:2041:HOH:O	2.20	0.42
2:B:12:LEU:HG	2:B:14:ILE:HG23	2.01	0.42
2:B:205:PRO:HA	2:B:242:GLY:O	2.20	0.42
1:A:15:GLN:HE22	1:A:113:TRP:HB2	1.85	0.41
1:A:69:ALA:O	1:A:70:ARG:HB2	2.21	0.41
1:A:182:LYS:CB	1:A:183:SER:HA	2.48	0.41
2:B:42:MET:HE1	3:B:2032:HOH:O	2.21	0.41
2:B:181:GLN:O	2:B:187:SER:HB2	2.21	0.40
2:B:38:GLN:HG3	2:B:89:GLN:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/191 (99%)	182 (96%)	6 (3%)	1 (0%)	29	29
2	B	242/244 (99%)	233 (96%)	7 (3%)	2 (1%)	19	17
All	All	431/435 (99%)	415 (96%)	13 (3%)	3 (1%)	22	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	244	ALA
1	A	97	SER
2	B	4	ASP



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/167 (100%)	151 (90%)	16 (10%)	8	6
2	B	216/216 (100%)	199 (92%)	17 (8%)	12	10
All	All	383/383 (100%)	350 (91%)	33 (9%)	10	9

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	27	SER
1	A	52	THR
1	A	57	LYS
1	A	58	LYS
1	A	66	PHE
1	A	87	LEU
1	A	94	ASP
1	A	98	THR
1	A	102	LEU
1	A	126	LEU
1	A	133	ASP
1	A	144	SER
1	A	145	GLN
1	A	152	LYS
1	A	162	CYS
2	B	2	GLU
2	B	14	ILE
2	B	16	THR
2	B	18	LYS
2	B	42	MET
2	B	58	LYS
2	B	73	GLU
2	B	81	SER
2	B	88	SER
2	B	118	LEU
2	B	139	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	140	GLN
2	B	176	GLN
2	B	178	LEU
2	B	184	LEU
2	B	185	ASN
2	B	194	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	4	GLN
1	A	15	GLN
1	A	109	GLN
1	A	145	GLN
1	A	189	ASN
2	B	138	HIS
2	B	140	GLN
2	B	168	HIS
2	B	207	ASN
2	B	208	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	191/191 (100%)	0.59	18 (9%) 8 9	10, 28, 51, 62	0
2	B	244/244 (100%)	0.55	19 (7%) 13 14	5, 27, 45, 68	0
All	All	435/435 (100%)	0.57	37 (8%) 10 12	5, 28, 48, 68	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	97	SER	7.6
2	B	3	ALA	5.7
1	A	100	GLY	4.9
1	A	184	ASP	4.6
2	B	183	ALA	4.4
2	B	227	ASP	4.0
2	B	2	GLU	3.9
2	B	42	MET	3.8
2	B	245	ASP	3.8
1	A	152	LYS	3.5
2	B	244	ALA	3.1
2	B	99	ASP	3.0
1	A	133	ASP	2.9
2	B	80	GLU	2.8
2	B	225	THR	2.7
1	A	185	PHE	2.6
1	A	95	ARG	2.5
1	A	183	SER	2.5
1	A	98	THR	2.4
1	A	181	ASN	2.4
1	A	53	GLY	2.3
1	A	3	ASN	2.3
1	A	91	ALA	2.3
2	B	185	ASN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	226	GLN	2.2
1	A	104	PHE	2.2
2	B	83	ARG	2.2
1	A	119	PRO	2.2
2	B	100	GLY	2.2
1	A	102	LEU	2.2
1	A	186	ALA	2.2
2	B	147	LEU	2.2
2	B	191	LEU	2.1
2	B	41	GLY	2.1
2	B	70	ILE	2.1
1	A	90	CYS	2.1
2	B	242	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.