



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 7, 2020 – 03:27 PM BST

PDB ID : 7CDV  
Title : STRUCTURE OF A NOVEL VIRULENCE REGULATION FACTOR SghR  
Authors : Ye, F.Z.; Wang, C.; Yan, X.F.; Zhang, L.H.; Gao, Y.G.  
Deposited on : 2020-06-20  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.14.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>66%</span> <span>8%</span> <span>25%</span> </div> </div>
1	B	370	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>66%</span> <span>8%</span> <span>25%</span> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LacI-type transcription factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	0	0
			2156	1364	381	402	9			
1	B	276	Total	C	N	O	S	0	0	0
			2141	1354	378	400	9			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP A0A2I4PGE9
A	-18	GLY	-	expression tag	UNP A0A2I4PGE9
A	-17	SER	-	expression tag	UNP A0A2I4PGE9
A	-16	SER	-	expression tag	UNP A0A2I4PGE9
A	-15	HIS	-	expression tag	UNP A0A2I4PGE9
A	-14	HIS	-	expression tag	UNP A0A2I4PGE9
A	-13	HIS	-	expression tag	UNP A0A2I4PGE9
A	-12	HIS	-	expression tag	UNP A0A2I4PGE9
A	-11	HIS	-	expression tag	UNP A0A2I4PGE9
A	-10	HIS	-	expression tag	UNP A0A2I4PGE9
A	-9	SER	-	expression tag	UNP A0A2I4PGE9
A	-8	SER	-	expression tag	UNP A0A2I4PGE9
A	-7	GLY	-	expression tag	UNP A0A2I4PGE9
A	-6	LEU	-	expression tag	UNP A0A2I4PGE9
A	-5	VAL	-	expression tag	UNP A0A2I4PGE9
A	-4	PRO	-	expression tag	UNP A0A2I4PGE9
A	-3	ARG	-	expression tag	UNP A0A2I4PGE9
A	-2	GLY	-	expression tag	UNP A0A2I4PGE9
A	-1	SER	-	expression tag	UNP A0A2I4PGE9
A	0	HIS	-	expression tag	UNP A0A2I4PGE9
B	-19	MET	-	expression tag	UNP A0A2I4PGE9
B	-18	GLY	-	expression tag	UNP A0A2I4PGE9
B	-17	SER	-	expression tag	UNP A0A2I4PGE9
B	-16	SER	-	expression tag	UNP A0A2I4PGE9
B	-15	HIS	-	expression tag	UNP A0A2I4PGE9

*Continued on next page...*



*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0A2I4PGE9
B	-13	HIS	-	expression tag	UNP A0A2I4PGE9
B	-12	HIS	-	expression tag	UNP A0A2I4PGE9
B	-11	HIS	-	expression tag	UNP A0A2I4PGE9
B	-10	HIS	-	expression tag	UNP A0A2I4PGE9
B	-9	SER	-	expression tag	UNP A0A2I4PGE9
B	-8	SER	-	expression tag	UNP A0A2I4PGE9
B	-7	GLY	-	expression tag	UNP A0A2I4PGE9
B	-6	LEU	-	expression tag	UNP A0A2I4PGE9
B	-5	VAL	-	expression tag	UNP A0A2I4PGE9
B	-4	PRO	-	expression tag	UNP A0A2I4PGE9
B	-3	ARG	-	expression tag	UNP A0A2I4PGE9
B	-2	GLY	-	expression tag	UNP A0A2I4PGE9
B	-1	SER	-	expression tag	UNP A0A2I4PGE9
B	0	HIS	-	expression tag	UNP A0A2I4PGE9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	192	Total O 192 192	0	0
2	B	193	Total O 193 193	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 
- 
- 66% 8% 25%
- MET GLY SER SER HIS HIS HIS HIS HIS HIS SER SER GLY LEU VAL PRO ARG GLY SER HIS MET ASN ASP THR GLY ASP GLU ALA LYS ALA THR THR THR GLY GLU ARG ARG PRO THR LEU LYS THR THR ILE ALA TYR MET THR GLY LEU GLY ILE THR THR VAL SER ARG ALA LEU
- LYS ASP ALA PRO ASP ILE GLY ALA THR LYS GLU ARG VAL ARG LEU ILE ALA ALA GLN ILE GLY TYR GLN PRO ASN ARG ALA GLY VAL ARG LEU
- R73 R74 R97 T102 P116 L131 D148 P149 M169 D179 P204 P205 S206 F210 H211 R215 P232
- D224 E229 I237 E238 T239 I244 I245 D255 D259 G260 I261 L272 E277 V281 D286 I287 V290 I301 Q302 H306 I312 R317 R340 R349 PRO

- Chain B:
- 
- 66% 8% 25%
- | Amino Acid | Frequency (%) |
|------------|---------------|
| Met        | 0.1           |
| Gly        | 0.1           |
| Ser        | 0.1           |
| Ser        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| His        | 0.1           |
| Ser        | 0.1           |
| Ser        | 0.1           |
| Gly        | 0.1           |
| Leu        | 0.1           |
| Val        | 0.1           |
| Pro        | 0.1           |
| Arg        | 0.1           |
| Gly        | 0.1           |
| Ser        | 0.1           |
| His        | 0.1           |
| Met        | 0.1           |
| Asn        | 0.1           |
| Asp        | 0.1           |
| Thr        | 0.1           |
| Gly        | 0.1           |
| Asn        | 0.1           |
| Ser        | 0.1           |
| Gly        | 0.1           |
| Arg        | 0.1           |
| Asp        | 0.1           |
| Glu        | 0.1           |
| Ala        | 0.1           |
| Ala        | 0.1           |
| Lys        | 0.1           |
| Ala        | 0.1           |
| Thr        | 0.1           |
| Thr        | 0.1           |
| Gly        | 0.1           |
| Glu        | 0.1           |
| Arg        | 0.1           |
| Pro        | 0.1           |
| Thr        | 0.1           |
| Leu        | 0.1           |
| Lys        | 0.1           |
| Thr        | 0.1           |
| Ile        | 0.1           |
| Ala        | 0.1           |
| Tyr        | 0.1           |
| Met        | 0.1           |
| Thr        | 0.1           |
| Gly        | 0.1           |
| Leu        | 0.1           |
| Ile        | 0.1           |
| Thr        | 0.1           |
| Val        | 0.1           |
| Ser        | 0.1           |
| Arg        | 0.1           |
| Ala        | 0.1           |
| Leu        | 0.1           |
- | Amino Acid | Frequency (%) |
|------------|---------------|
| Lys        | 0.1           |
| Asp        | 0.1           |
| Ala        | 0.1           |
| Pro        | 0.1           |
| Asp        | 0.1           |
| Ile        | 0.1           |
| Gly        | 0.1           |
| Glu        | 0.1           |
| Thr        | 0.1           |
| Lys        | 0.1           |
| Glu        | 0.1           |
| Arg        | 0.1           |
| Val        | 0.1           |
| Arg        | 0.1           |
| Leu        | 0.1           |
| Ile        | 0.1           |
| Ala        | 0.1           |
| Gln        | 0.1           |
| Gln        | 0.1           |
| Ile        | 0.1           |
| Gly        | 0.1           |
| Tyr        | 0.1           |
| Gln        | 0.1           |
| Pro        | 0.1           |
| Asn        | 0.1           |
| Arg        | 0.1           |
| Ala        | 0.1           |
| Gly        | 0.1           |
| Val        | 0.1           |
| Arg        | 0.1           |
| Leu        | 0.1           |
| Lys        | 0.1           |
| Tyr        | 0.1           |
| Gly        | 0.1           |
| Thr        | 0.1           |
| Thr        | 0.1           |
| Glu        | 0.1           |
| Arg        | 0.1           |
| Pro        | 0.1           |
| Thr        | 0.1           |
| Leu        | 0.1           |
| Ile        | 0.1           |
| Ala        | 0.1           |
| Tyr        | 0.1           |
| Met        | 0.1           |
| Thr        | 0.1           |
| Gly        | 0.1           |
| Leu        | 0.1           |
| Ile        | 0.1           |
| Thr        | 0.1           |
| Val        | 0.1           |
| Ser        | 0.1           |
| Arg        | 0.1           |
| Ala        | 0.1           |
| Leu        | 0.1           |
- | Amino Acid | Frequency (%) |
|------------|---------------|
| I237       | 0.1           |
| E238       | 0.1           |
| T239       | 0.1           |
| I244       | 0.1           |
| K245       | 0.1           |
| K249       | 0.1           |
| M252       | 0.1           |
| Q253       | 0.1           |
| S254       | 0.1           |
| D255       | 0.1           |
| D259       | 0.1           |
| G260       | 0.1           |
| I261       | 0.1           |
| E277       | 0.1           |
| V281       | 0.1           |
| D286       | 0.1           |
| I287       | 0.1           |
| V290       | 0.1           |
| I301       | 0.1           |
| Q302       | 0.1           |
| H306       | 0.1           |
| I312       | 0.1           |
| R340       | 0.1           |
| R349       | 0.1           |
| Pro        | 0.1           |

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.53Å 119.44Å 121.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.66 – 2.10 42.66 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.66-2.10) 99.6 (42.66-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.30 (at 2.10Å)	Xtriage
Refinement program	PHENIX DEV_1932	Depositor
R, $R_{free}$	0.202 , 0.240 0.203 , 0.238	Depositor DCC
$R_{free}$ test set	1655 reflections (5.32%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.8	Xtriage
Anisotropy	0.440	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4682	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.00 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9719e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/2201	0.42	1/2981 (0.0%)
1	B	0.27	0/2186	0.42	0/2963
All	All	0.25	0/4387	0.42	1/5944 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	245	ARG	NE-CZ-NH1	5.05	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2156	0	2160	22	0
1	B	2141	0	2131	18	0
2	A	192	0	0	3	0
2	B	193	0	0	3	0
All	All	4682	0	4291	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:223:ARG:NH2	1:B:224:ASP:OD1	2.14	0.81
1:A:179:ASP:OD2	1:A:340:ARG:NH2	2.25	0.70
1:B:277:GLU:OE2	1:B:302:GLN:NE2	2.29	0.65
1:A:245:ARG:HD2	2:A:550:HOH:O	1.95	0.64
1:A:223:ARG:NH2	2:A:402:HOH:O	2.28	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/370 (74%)	265 (96%)	10 (4%)	0	100	100
1	B	274/370 (74%)	265 (97%)	9 (3%)	0	100	100
All	All	549/740 (74%)	530 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/308 (76%)	229 (98%)	4 (2%)	60	67
1	B	230/308 (75%)	222 (96%)	8 (4%)	36	38
All	All	463/616 (75%)	451 (97%)	12 (3%)	46	50



5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	210	PHE
1	B	215	ARG
1	B	255	ASP
1	B	131	LEU
1	B	253	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	277/370 (74%)	-0.21	3 (1%) 80 84	10, 24, 48, 66	0
1	B	276/370 (74%)	-0.16	5 (1%) 68 72	10, 24, 46, 94	0
All	All	553/740 (74%)	-0.18	8 (1%) 75 78	10, 24, 46, 94	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	74	THR	11.9
1	A	74	THR	4.5
1	B	234	ALA	3.4
1	A	301	ILE	2.7
1	B	301	ILE	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.