



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2021 – 02:09 PM JST

PDB ID : 7CE2  
Title : The Crystal structure of TeNT Hc complexed with neutralizing antibody  
Authors : Wang, X.; Wang, Y.; Wu, C.; Yu, J.; Liao, H.  
Deposited on : 2020-06-21  
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

**i**

## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	458	<div><div>5%</div><div>85%</div><div>12%</div><div>••</div></div>	
2	Z	223	<div><div>3%</div><div>88%</div><div>9%</div><div>•</div></div>	
3	B	214	<div><div>3%</div><div>90%</div><div>8%</div><div>••</div></div>	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetanus toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	0	0
			3619	2317	602	690	10			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	864	MET	-	initiating methionine	UNP Q9LA13
A	1316	HIS	-	expression tag	UNP Q9LA13
A	1317	HIS	-	expression tag	UNP Q9LA13
A	1318	HIS	-	expression tag	UNP Q9LA13
A	1319	HIS	-	expression tag	UNP Q9LA13
A	1320	HIS	-	expression tag	UNP Q9LA13
A	1321	HIS	-	expression tag	UNP Q9LA13

- Molecule 2 is a protein called neutralizing antibody heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Z	216	Total	C	N	O	S	0	0	0
			1635	1034	274	322	5			

- Molecule 3 is a protein called neutralizing antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	212	Total	C	N	O	S	0	1	0
			1632	1017	278	332	5			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	159	Total	O	0	0
			159	159		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Z	76	Total 76	O 76	0	0
4	B	58	Total 58	O 58	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tetanus toxin



- Molecule 2: neutralizing antibody heavy chain



- Molecule 3: neutralizing antibody light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.69Å 136.99Å 137.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.07 – 2.01 45.05 – 2.01	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.07-2.01) 99.9 (45.05-2.01)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 2.01Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.180 , 0.218 0.181 , 0.219	Depositor DCC
$R_{free}$ test set	3700 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.4	Xtriage
Anisotropy	0.585	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7179	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	1/3701 (0.0%)	0.63	3/5024 (0.1%)
2	Z	0.39	0/1677	0.59	0/2289
3	B	0.38	0/1670	0.59	0/2266
All	All	0.41	1/7048 (0.0%)	0.61	3/9579 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	967	GLU	CD-OE1	-5.04	1.20	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	985	LEU	CB-CG-CD2	-6.06	100.69	111.00
1	A	883	LYS	CB-CG-CD	-5.83	96.45	111.60
1	A	883	LYS	CD-CE-NZ	-5.71	98.56	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	939	MET	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3619	0	3550	46	0
2	Z	1635	0	1603	12	0
3	B	1632	0	1583	15	0
4	A	159	0	0	2	0
4	B	58	0	0	1	0
4	Z	76	0	0	0	0
All	All	7179	0	6736	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:871:VAL:HG23	1:A:883:LYS:HE2	1.44	0.99
1:A:985:LEU:HD12	1:A:986:SER:N	1.77	0.99
1:A:985:LEU:CD1	1:A:986:SER:OG	2.14	0.95
1:A:1274:GLN:NE2	1:A:1278:ASP:O	2.01	0.94
2:Z:161:SER:H	2:Z:202:ASN:HD21	1.26	0.81
1:A:871:VAL:CG2	1:A:883:LYS:HE2	2.14	0.77
1:A:1002:TRP:HE1	1:A:1038:ASN:HD21	1.33	0.77
1:A:880:ILE:HA	1:A:883:LYS:NZ	2.05	0.71
3:B:148:TRP:CD2	3:B:179:LEU:HD12	2.25	0.71
2:Z:86:THR:HG22	2:Z:88:ALA:H	1.57	0.69
1:A:1184:ASN:C	1:A:1186:ILE:HD11	2.12	0.69
3:B:190:LYS:HD3	3:B:191:VAL:N	2.09	0.67
1:A:985:LEU:HD13	1:A:986:SER:OG	1.95	0.67
1:A:985:LEU:HD12	1:A:986:SER:OG	1.94	0.67
1:A:871:VAL:HG21	1:A:883:LYS:HZ3	1.60	0.67
1:A:869:CYS:HB3	1:A:883:LYS:O	1.95	0.66
3:B:191:VAL:HG22	3:B:210:ASN:OD1	1.95	0.65
3:B:190:LYS:HD3	3:B:191:VAL:H	1.61	0.64
2:Z:161:SER:H	2:Z:202:ASN:ND2	1.96	0.64
1:A:1171:SER:O	4:A:1401:HOH:O	2.14	0.64
1:A:1183:ASN:O	1:A:1184:ASN:HB3	2.01	0.61

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:944:ASN:ND2	1:A:1066:GLU:OE1	2.34	0.61
1:A:891:ILE:HD13	1:A:914:LEU:HD22	1.82	0.60
3:B:148:TRP:CE3	3:B:179:LEU:HD12	2.36	0.60
1:A:985:LEU:HD12	1:A:986:SER:CB	2.33	0.59
1:A:985:LEU:CD1	1:A:986:SER:CB	2.83	0.57
1:A:947:PHE:HA	1:A:1040:ARG:HG3	1.88	0.56
1:A:985:LEU:HD12	1:A:986:SER:CA	2.35	0.55
1:A:1097:ASN:OD1	1:A:1100:GLU:HG3	2.06	0.54
1:A:1176:ILE:HD11	1:A:1201:SER:HB2	1.87	0.54
1:A:880:ILE:HA	1:A:883:LYS:HZ3	1.73	0.54
1:A:871:VAL:HG21	1:A:883:LYS:NZ	2.24	0.52
1:A:944:ASN:O	1:A:948:ASN:HB2	2.09	0.52
1:A:1223:ARG:NH1	4:A:1406:HOH:O	2.39	0.51
1:A:967:GLU:HB3	1:A:1162:LEU:HD21	1.93	0.50
3:B:148:TRP:CG	3:B:179:LEU:HD12	2.45	0.50
2:Z:156:THR:OG1	2:Z:204:ASN:HB2	2.12	0.49
3:B:115:VAL:HG21	3:B:196:VAL:HG21	1.93	0.49
1:A:871:VAL:CG2	1:A:883:LYS:CE	2.87	0.49
3:B:124:GLN:NE2	4:B:305:HOH:O	2.45	0.49
1:A:982:LYS:HD2	1:A:1075:ASP:OD2	2.13	0.48
1:A:879:VAL:HG12	1:A:883:LYS:CE	2.43	0.48
2:Z:87:ALA:HA	2:Z:116:VAL:HG13	1.96	0.48
1:A:1135:TYR:HB3	1:A:1138:LYS:HG3	1.96	0.47
2:Z:12:VAL:O	2:Z:116:VAL:HA	2.15	0.47
3:B:185:ASP:O	3:B:188:LYS:HB2	2.14	0.47
3:B:125:LEU:O	3:B:183:LYS:HD3	2.15	0.46
1:A:879:VAL:C	1:A:883:LYS:HZ2	2.19	0.46
1:A:870:TRP:CD2	1:A:871:VAL:N	2.84	0.46
2:Z:86:THR:O	2:Z:116:VAL:HG11	2.16	0.46
2:Z:11:LEU:HD13	2:Z:115:THR:HB	1.98	0.45
3:B:190:LYS:CD	3:B:191:VAL:HG23	2.47	0.45
3:B:197:THR:HG22	3:B:204:PRO:HB3	1.99	0.45
3:B:183:LYS:O	3:B:187:GLU:HG3	2.17	0.44
1:A:985:LEU:H	1:A:985:LEU:HG	1.47	0.44
1:A:870:TRP:CG	1:A:871:VAL:N	2.86	0.43
1:A:1021:ASP:HB2	1:A:1165:TYR:CZ	2.53	0.43
2:Z:99:LYS:HB3	2:Z:102:THR:OG1	2.19	0.43
2:Z:17:THR:HG22	2:Z:83:ARG:HA	2.01	0.43
3:B:134:CYS:HB2	3:B:148:TRP:CH2	2.54	0.43
1:A:879:VAL:HG12	1:A:883:LYS:HE3	2.01	0.43
1:A:981:LYS:O	1:A:987:ILE:O	2.37	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:904:SER:HA	1:A:938:ALA:HB2	2.00	0.42
1:A:875:GLU:HG3	1:A:879:VAL:HB	2.01	0.42
2:Z:110:GLN:H	2:Z:110:GLN:CD	2.21	0.42
1:A:941:ILE:HD12	1:A:941:ILE:N	2.35	0.41
1:A:1170:TYR:CE1	1:A:1308:THR:HA	2.56	0.41
1:A:1023:PHE:CE2	1:A:1136:SER:HB2	2.55	0.41
1:A:890:ASP:HB2	1:A:899:ILE:HD13	2.03	0.40
1:A:1051:VAL:O	1:A:1053:MET:HG2	2.21	0.40
1:A:1145:ILE:HD11	1:A:1207:HIS:CG	2.57	0.40
2:Z:4:LEU:HD22	2:Z:24:VAL:HG22	2.03	0.40
3:B:146:VAL:HG22	3:B:196:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/458 (96%)	421 (95%)	20 (4%)	1 (0%)	47	44
2	Z	212/223 (95%)	208 (98%)	4 (2%)	0	100	100
3	B	211/214 (99%)	205 (97%)	6 (3%)	0	100	100
All	All	865/895 (97%)	834 (96%)	30 (4%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	872	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/416 (97%)	398 (98%)	6 (2%)	65	69
2	Z	189/195 (97%)	188 (100%)	1 (0%)	88	92
3	B	185/186 (100%)	181 (98%)	4 (2%)	52	55
All	All	778/797 (98%)	767 (99%)	11 (1%)	67	72

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	939	MET
1	A	940	ASP
1	A	947	PHE
1	A	1019	LEU
1	A	1065	ARG
1	A	1073	LYS
2	Z	221	CYS
3	B	61	ASP
3	B	105	GLU
3	B	188	LYS
3	B	190	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	998	ASN
1	A	1038	ASN
2	Z	202	ASN
3	B	124	GLN
3	B	137	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	446/458 (97%)	0.53	25 (5%) 24 23	31, 49, 91, 143	0
2	Z	216/223 (96%)	0.22	6 (2%) 53 51	34, 49, 72, 121	0
3	B	212/214 (99%)	0.16	7 (3%) 46 45	34, 56, 89, 95	0
All	All	874/895 (97%)	0.36	38 (4%) 35 34	31, 51, 88, 143	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	869	CYS	12.9
1	A	947	PHE	10.8
1	A	943	TYR	8.1
1	A	902	PHE	5.9
1	A	946	MET	4.4
1	A	938	ALA	4.3
1	A	985	LEU	4.3
1	A	939	MET	4.1
1	A	941	ILE	3.6
1	A	944	ASN	3.6
1	A	883	LYS	3.5
1	A	987	ILE	2.9
3	B	153	ALA	2.8
3	B	188	LYS	2.8
1	A	870	TRP	2.6
1	A	1133	VAL	2.6
1	A	940	ASP	2.5
2	Z	83	ARG	2.5
1	A	942	GLU	2.5
1	A	1074	LEU	2.5
3	B	150	VAL	2.4
1	A	1065	ARG	2.4
2	Z	133	SER	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	B	190	LYS	2.4
1	A	1063	ALA	2.3
2	Z	2	VAL	2.3
1	A	988	GLY	2.3
2	Z	26	GLY	2.2
3	B	193	ALA	2.2
3	B	205	VAL	2.2
2	Z	1	GLN	2.2
2	Z	221	CYS	2.2
1	A	1061	LEU	2.1
1	A	1250	THR	2.1
1	A	970	ASP	2.1
1	A	886	ILE	2.1
1	A	1134	ALA	2.0
3	B	169	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.