



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 06:49 am BST

PDB ID : 3CHW  
Title : Complex of Dictyostelium discoideum Actin with Profilin and the Last Poly-Pro of Human VASP  
Authors : Baek, K.; Dominguez, R.  
Deposited on : 2008-03-10  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• •</div> </div> </div>
2	P	139	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>22%</div> <div>•</div> </div> </div>
3	V	16	<div> <div>31%</div> <div> <div>50%</div> <div>25%</div> <div>6%</div> <div>19%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major actin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	6	0
			2855	1813	480	542	20			

- Molecule 2 is a protein called Profilin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	139	Total	C	N	O	S	0	7	0
			1083	684	182	210	7			

- Molecule 3 is a protein called Vasodilator-stimulated phosphoprotein 16-residue peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	V	13	Total	C	N	O	0	0	0
			82	54	14	14			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	213	Total	O	0	0
			213	213		
6	P	102	Total	O	0	0
			102	102		
6	V	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major actin



- Molecule 2: Profilin-1



- Molecule 3: Vasodilator-stimulated phosphoprotein 16-residue peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.04Å 76.18Å 180.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.87 – 2.30 38.09 – 2.30	Depositor EDS
% Data completeness (in resolution range)	89.6 (38.87-2.30) 89.6 (38.09-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	15.68 (at 2.31Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.157 , 0.227 0.170 , 0.229	Depositor DCC
$R_{free}$ test set	1101 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.0	Xtriage
Anisotropy	0.309	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4369	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HIC, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.55	8/2919 (0.3%)	0.80	1/3947 (0.0%)
2	P	1.49	1/1123 (0.1%)	0.91	4/1516 (0.3%)
3	V	1.23	0/87	0.78	0/123
All	All	1.53	9/4129 (0.2%)	0.83	5/5586 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	294	TYR	CD1-CE1	-5.91	1.30	1.39
1	A	188	TYR	CD2-CE2	-5.53	1.31	1.39
1	A	188	TYR	CD1-CE1	-5.45	1.31	1.39
1	A	168	GLY	C-O	-5.37	1.15	1.23
2	P	84	SER	CB-OG	-5.36	1.35	1.42
1	A	53	TYR	CD2-CE2	-5.27	1.31	1.39
1	A	294	TYR	CD2-CE2	-5.17	1.31	1.39
1	A	344	SER	CB-OG	-5.07	1.35	1.42
1	A	362	TYR	CD1-CE1	-5.03	1.31	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	56[A]	SER	N-CA-CB	-5.35	102.47	110.50
2	P	56[B]	SER	N-CA-CB	-5.35	102.47	110.50
2	P	132[A]	SER	N-CA-CB	5.34	118.52	110.50
2	P	132[B]	SER	N-CA-CB	5.34	118.52	110.50
1	A	62	ARG	NE-CZ-NH1	5.22	122.91	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2855	0	2841	48	0
2	P	1083	0	1093	16	0
3	V	82	0	81	9	0
4	A	1	0	0	0	0
5	A	31	0	12	2	0
6	A	213	0	0	5	0
6	P	102	0	0	2	0
6	V	2	0	0	0	0
All	All	4369	0	4027	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:V:213:GLN:N	3:V:213:GLN:HE21	1.57	1.01
1:A:196:ARG:HH11	1:A:196:ARG:CG	1.73	1.01
1:A:196:ARG:HH11	1:A:196:ARG:HG3	1.27	0.98
3:V:213:GLN:H	3:V:213:GLN:HE21	1.02	0.97
1:A:196:ARG:HG3	1:A:196:ARG:NH1	1.85	0.86
3:V:213:GLN:N	3:V:213:GLN:NE2	2.30	0.79
1:A:172:PRO:HA	1:A:175:ILE:HD12	1.65	0.78
1:A:244:ASP:OD1	1:A:246:GLN:HB3	1.85	0.77
3:V:213:GLN:H	3:V:213:GLN:NE2	1.83	0.75
1:A:257:CYS:HB3	1:A:258:PRO:HD3	1.73	0.71
1:A:351:THR:O	1:A:354:GLN:HG2	1.91	0.71
1:A:196:ARG:HD2	1:A:198:TYR:HE2	1.54	0.70
1:A:194:THR:HA	1:A:198:TYR:O	1.92	0.69
1:A:196:ARG:NH1	6:A:565:HOH:O	2.23	0.66
1:A:214:GLU:HG2	5:A:402:ATP:C5	2.29	0.66
1:A:314:ASN:ND2	6:A:580:HOH:O	2.27	0.66
1:A:352:PHE:HE2	1:A:356:TRP:CZ2	2.14	0.66
1:A:273:GLY:O	1:A:277:THR:HG23	1.96	0.65
2:P:20:ALA:HB2	2:P:39:PHE:HB3	1.81	0.62

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:THR:N	1:A:205:GLU:OE1	2.28	0.61
1:A:196:ARG:HD2	1:A:198:TYR:CE2	2.36	0.60
1:A:115:ASN:O	1:A:119:MET:HG3	2.02	0.60
1:A:196:ARG:HH11	1:A:196:ARG:HG2	1.66	0.58
2:P:13:ASP:OD2	2:P:126:LYS:NZ	2.36	0.58
1:A:300:SER:HB2	1:A:338[A]:SER:OG	2.03	0.58
2:P:37:LYS:HE3	6:P:186:HOH:O	2.06	0.56
1:A:20:GLY:HA3	1:A:340:TRP:HZ2	1.70	0.55
1:A:336:LYS:HB2	6:A:440:HOH:O	2.05	0.55
2:P:137:SER:O	2:P:138[A]:GLN:HB2	2.07	0.55
1:A:16:MET:CE	1:A:30:VAL:HG12	2.39	0.53
1:A:31:PHE:HB2	1:A:32:PRO:HD2	1.91	0.53
1:A:244:ASP:CG	1:A:244:ASP:O	2.46	0.52
1:A:257:CYS:CB	1:A:258:PRO:HD3	2.38	0.52
1:A:20:GLY:HA3	1:A:340:TRP:CZ2	2.45	0.52
2:P:112:LEU:HD12	2:P:127:CYS:HB2	1.92	0.52
1:A:250:ILE:HD12	1:A:253:GLU:HB2	1.92	0.51
1:A:143:TYR:HE1	6:A:595:HOH:O	1.93	0.51
1:A:200:PHE:HD1	1:A:205:GLU:HG2	1.77	0.50
1:A:244:ASP:O	1:A:244:ASP:OD2	2.29	0.49
1:A:352:PHE:HE2	1:A:356:TRP:CH2	2.30	0.49
1:A:151:ILE:HD12	1:A:164:PRO:HG3	1.95	0.49
1:A:16:MET:HE3	1:A:30:VAL:HG12	1.94	0.48
1:A:334:GLU:N	1:A:334:GLU:OE1	2.40	0.48
2:P:34:VAL:HG22	2:P:37:LYS:CG	2.44	0.48
3:V:209:LEU:CD2	3:V:210:PRO:HD2	2.44	0.47
2:P:1:ALA:HA	3:V:206:ALA:HB3	1.97	0.47
1:A:157:ASP:HB2	5:A:402:ATP:H5'1	1.98	0.46
2:P:18:ASP:OD1	2:P:34:VAL:HG13	2.16	0.46
1:A:180:LEU:HD22	1:A:269:MET:SD	2.56	0.45
1:A:18:LYS:N	1:A:18:LYS:CD	2.80	0.45
1:A:354:GLN:HG3	1:A:355:MET:HG3	1.99	0.44
2:P:76:SER:HA	2:P:79[B]:GLN:OE1	2.17	0.44
1:A:250:ILE:CG1	1:A:254:ARG:HG3	2.48	0.44
1:A:200:PHE:CD1	1:A:205:GLU:HG2	2.54	0.43
2:P:42:ILE:HD12	2:P:47:VAL:HG23	2.00	0.43
2:P:63:LEU:C	2:P:63:LEU:HD12	2.38	0.43
1:A:206:ARG:NH1	6:A:606:HOH:O	2.52	0.43
2:P:114:GLY:HA3	2:P:123:ILE:HD11	2.00	0.43
1:A:272:ALA:HB1	1:A:276:GLU:HB3	2.01	0.43
2:P:34:VAL:HA	2:P:35:PRO:HD3	1.77	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:57:SER:HB2	6:P:164:HOH:O	2.19	0.43
1:A:106:THR:HB	1:A:137:GLN:HG3	2.01	0.43
3:V:209:LEU:HD23	3:V:210:PRO:HD2	2.01	0.42
1:A:200:PHE:C	1:A:201:THR:HG23	2.41	0.41
3:V:209:LEU:HB3	3:V:210:PRO:HD2	2.02	0.41
2:P:27:SER:HA	2:P:28:PRO:HD2	1.91	0.41
1:A:70:PRO:HB3	1:A:81:ASP:HB2	2.03	0.41
1:A:138:ALA:HB1	1:A:152:VAL:HB	2.03	0.41
1:A:242:LEU:HA	1:A:242:LEU:HD23	1.82	0.41
1:A:257:CYS:HB3	1:A:258:PRO:CD	2.45	0.40
2:P:139:TYR:CD2	3:V:207:PRO:HG3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/375 (96%)	351 (97%)	10 (3%)	1 (0%)	41	50
2	P	144/139 (104%)	140 (97%)	4 (3%)	0	100	100
3	V	11/16 (69%)	11 (100%)	0	0	100	100
All	All	517/530 (98%)	502 (97%)	14 (3%)	1 (0%)	47	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	180	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/314 (99%)	289 (93%)	21 (7%)	16	21
2	P	120/113 (106%)	111 (92%)	9 (8%)	13	17
3	V	8/8 (100%)	7 (88%)	1 (12%)	4	5
All	All	438/435 (101%)	407 (93%)	31 (7%)	14	19

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	VAL
1	A	6	GLN
1	A	25	ASP
1	A	57	GLU
1	A	119	MET
1	A	180	LEU
1	A	196	ARG
1	A	215	LYS
1	A	221	LEU
1	A	228	GLN
1	A	232	SER
1	A	244	ASP
1	A	246	GLN
1	A	249	THR
1	A	250	ILE
1	A	277	THR
1	A	300	SER
1	A	317	LEU
1	A	326	LYS
1	A	336	LYS
1	A	373	LYS
2	P	30	VAL
2	P	34	VAL
2	P	53	LYS
2	P	58	PHE
2	P	71	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	P	82	GLU
2	P	122	LEU
2	P	138[A]	GLN
2	P	138[B]	GLN
3	V	213	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	275	HIS
2	P	61	ASN
3	V	213	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	HIC	A	73	1	8,11,12	1.66	1 (12%)	6,14,16	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	0/5/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	HIC	CD2-NE2	-4.38	1.31	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ATP	A	402	4	26,33,33	1.03	2 (7%)	31,52,52	1.35	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ATP	A	402	4	-	2/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	402	ATP	C2-N3	2.31	1.35	1.32
5	A	402	ATP	C5-C4	2.22	1.46	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	402	ATP	N3-C2-N1	-3.30	123.52	128.68
5	A	402	ATP	N6-C6-N1	2.38	123.51	118.57
5	A	402	ATP	O2B-PB-O1B	2.25	123.34	112.24
5	A	402	ATP	PA-O3A-PB	-2.05	125.78	132.83
5	A	402	ATP	C2-N1-C6	2.01	122.20	118.75
5	A	402	ATP	C4-C5-N7	-2.00	107.31	109.40

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	402	ATP	PB-O3B-PG-O2G
5	A	402	ATP	PB-O3B-PG-O1G

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	402	ATP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/375 (96%)	-0.05	10 (2%) 53 60	7, 20, 46, 69	0
2	P	139/139 (100%)	-0.25	1 (0%) 87 91	9, 19, 31, 51	0
3	V	13/16 (81%)	1.98	5 (38%) 0 0	30, 35, 48, 52	0
All	All	512/530 (96%)	-0.05	16 (3%) 49 56	7, 20, 44, 69	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	V	201	GLY	8.5
1	A	245	GLY	6.1
3	V	212	ALA	4.5
1	A	199	SER	3.3
1	A	243	PRO	3.3
1	A	40	HIS	3.3
1	A	200	PHE	3.2
1	A	203	THR	2.7
3	V	202	GLY	2.7
1	A	247	VAL	2.6
2	P	1	ALA	2.6
3	V	213	GLN	2.5
1	A	244	ASP	2.5
3	V	211	ALA	2.3
1	A	248	ILE	2.1
1	A	5	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column



labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	HIC	A	73	11/12	0.95	0.12	11,22,40,41	0

### 6.3 Carbohydrates


There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ATP	A	402	31/31	0.99	0.12	2,9,14,16	0
4	CA	A	401	1/1	0.99	0.02	5,5,5,5	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.