



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 01:47 AM BST

PDB ID : 6CHF
Title : Crystal structure of a Fc fragment LALA mutant (L234A, L235A) of human IgG1 (crystal form 1)
Authors : Gohain, N.; Tolbert, W.D.; Pazgier, M.
Deposited on : 2018-02-22
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 3% 80% 17%
1	B	214	 9% 71% 22% 5%
2	C	8	 25% 63% 13%
2	D	8	 63% 38%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein DKFZp686C11235.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	211	Total 1688	C 1072	N 285	O 324	S 7	0	1	0
1	B	208	Total 1668	C 1062	N 280	O 319	S 7	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	ALA	LEU	engineered mutation	UNP Q6MZV7
A	235	ALA	LEU	engineered mutation	UNP Q6MZV7
B	234	ALA	LEU	engineered mutation	UNP Q6MZV7
B	235	ALA	LEU	engineered mutation	UNP Q6MZV7

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	Total 99	C 56	N 4	O 39	0	0	0
2	D	8	Total 99	C 56	N 4	O 39	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total 27	O 27	0	0
3	B	20	Total 20	O 20	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein DKFZp686C11235



- Molecule 1: Uncharacterized protein DKFZp686C11235



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3
MAN4
MAG5
MAN6
MAG7
FUC8

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.91Å 73.81Å 136.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.05 – 2.40 46.05 – 2.39	Depositor EDS
% Data completeness (in resolution range)	89.3 (46.05-2.40) 89.1 (46.05-2.39)	Depositor EDS
R_{merge}	0.32	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.216 , 0.273 0.216 , 0.273	Depositor DCC
R_{free} test set	921 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtrriage
Anisotropy	0.176	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3601	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3283e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1734	0.61	0/2361
1	B	0.55	0/1715	0.72	2/2336 (0.1%)
All	All	0.54	0/3449	0.67	2/4697 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	297	ASN	N-CA-CB	-7.23	97.59	110.60
1	B	296	TYR	C-N-CA	-6.45	105.57	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1650	23	0
1	B	1668	0	1635	54	0
2	C	99	0	85	1	0
2	D	99	0	85	2	0
3	A	27	0	0	0	0
3	B	20	0	0	1	0
All	All	3601	0	3455	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:TYR:OH	1:B:322:LYS:HD2	1.66	0.94
1:B:320:LYS:HE3	1:B:335:THR:HG23	1.57	0.84
1:B:265:ASP:HA	1:B:299:THR:HG22	1.63	0.80
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.64	0.80
1:B:278:TYR:CZ	1:B:322:LYS:HD2	2.17	0.79
1:B:328:LEU:HD12	1:B:329:PRO:HD2	1.66	0.78
1:B:278:TYR:OH	1:B:322:LYS:CD	2.33	0.76
1:A:266:VAL:HB	1:A:300:TYR:HB2	1.72	0.70
1:B:322:LYS:HA	1:B:333:GLU:HG3	1.77	0.67
1:B:320:LYS:HD3	1:B:333:GLU:HB3	1.74	0.67
1:B:248:LYS:CE	1:B:380:GLU:OE1	2.44	0.66
1:B:248:LYS:NZ	1:B:380:GLU:OE1	2.28	0.65
1:B:276:ASN:HB2	1:B:322:LYS:HG2	1.80	0.63
1:B:296:TYR:O	1:B:298:SER:N	2.32	0.62
1:A:358:MET:O	1:A:414:LYS:NZ	2.28	0.62
1:B:278:TYR:HA	1:B:282:VAL:O	1.98	0.62
1:B:268:HIS:CE1	1:B:298:SER:HB2	2.38	0.59
1:A:414:LYS:O	1:A:418:GLN:HG2	2.03	0.58
1:B:245:PRO:HD3	1:B:259:VAL:HG22	1.85	0.58
1:B:276:ASN:HB2	1:B:322:LYS:CD	2.34	0.58
1:B:242:LEU:HD22	1:B:336:ILE:HG12	1.86	0.57
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.85	0.57
1:B:297:ASN:O	1:B:299:THR:N	2.37	0.57
1:B:278:TYR:HE2	1:B:283:GLU:HG3	1.70	0.57
1:B:328:LEU:HD12	1:B:329:PRO:CD	2.34	0.56
1:A:297:ASN:N	1:A:297:ASN:OD1	2.37	0.56
1:B:270:ASP:OD1	1:B:326:LYS:HE3	2.06	0.56
1:A:365:LEU:HD12	1:A:410:LEU:HD23	1.87	0.56
1:B:325:ASN:OD1	1:B:326:LYS:N	2.39	0.56
1:B:267:SER:O	1:B:271:PRO:HB3	2.06	0.56
1:B:296:TYR:O	1:B:297:ASN:C	2.38	0.56
1:B:293:GLU:O	1:B:300:TYR:HA	2.06	0.55
1:A:264:VAL:HG12	1:A:301:ARG:HG3	1.90	0.54
1:B:325:ASN:OD1	1:B:327:ALA:N	2.38	0.54
1:B:248:LYS:HE2	1:B:380:GLU:OE1	2.07	0.54
1:B:263:VAL:HB	1:B:302:VAL:HG13	1.89	0.53
1:B:240:VAL:HG22	1:B:263:VAL:HG22	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:MET:HE3	1:A:254:SER:H	1.74	0.53
2:D:1:NAG:O3	2:D:2:NAG:O5	2.28	0.51
1:B:265:ASP:HA	1:B:299:THR:CG2	2.38	0.51
1:B:294:GLU:HG3	1:B:300:TYR:CE2	2.45	0.51
1:B:276:ASN:HB2	1:B:322:LYS:CG	2.42	0.50
1:B:367:CYS:HB2	1:B:381:TRP:CZ2	2.47	0.50
1:B:278:TYR:CE1	1:B:322:LYS:HD2	2.48	0.47
1:A:328:LEU:HD13	1:A:332:ILE:HG13	1.95	0.47
1:A:320:LYS:HE3	1:A:333:GLU:OE1	2.13	0.47
1:B:268:HIS:HE1	1:B:298:SER:O	1.96	0.47
1:B:278:TYR:CE2	1:B:283:GLU:HG3	2.49	0.47
1:A:259:VAL:HG13	1:A:306:LEU:HB3	1.97	0.46
1:B:252:MET:CE	1:B:428:MET:HG2	2.46	0.46
1:A:234:ALA:N	1:A:267:SER:HB3	2.29	0.46
1:A:234:ALA:N	1:A:265:ASP:O	2.49	0.46
1:A:359:THR:HG23	1:A:360:LYS:HG2	1.97	0.46
1:B:267:SER:OG	1:B:269:GLU:HB2	2.16	0.46
1:B:270:ASP:HA	1:B:271:PRO:HD3	1.65	0.46
1:B:294:GLU:HG3	1:B:300:TYR:CZ	2.51	0.45
1:A:297:ASN:HD22	2:C:1:NAG:C7	2.28	0.45
1:A:298:SER:OG	1:A:298:SER:O	2.31	0.45
1:A:259:VAL:CG1	1:A:306:LEU:HB3	2.47	0.44
1:B:248:LYS:NZ	1:B:380:GLU:CD	2.70	0.44
1:B:274:LYS:HB3	1:B:324:SER:HB2	1.99	0.44
1:A:388:GLU:CD	1:A:416:ARG:HH22	2.21	0.44
1:B:243:PHE:CE1	2:D:3:BMA:H62	2.53	0.44
1:A:353:PRO:HD3	1:A:365:LEU:HD23	2.00	0.44
1:A:290:LYS:HB3	1:A:291:PRO:HD2	1.99	0.44
1:B:320:LYS:CE	1:B:335:THR:HG23	2.40	0.44
1:A:278:TYR:CE2	1:A:283:GLU:HG3	2.53	0.44
1:A:356:GLU:O	1:A:359:THR:HG22	2.17	0.44
1:B:261:CYS:HB2	1:B:277:TRP:CH2	2.53	0.43
1:B:248:LYS:HZ1	1:B:380:GLU:CD	2.21	0.43
1:A:350:THR:HB	1:A:441:LEU:HD22	2.01	0.42
1:B:295:GLN:NE2	1:B:301:ARG:HD2	2.34	0.42
1:B:296:TYR:CD1	1:B:296:TYR:N	2.88	0.41
1:B:274:LYS:HE2	1:B:276:ASN:OD1	2.20	0.41
1:B:240:VAL:HG21	1:B:323:VAL:HG21	2.03	0.41
1:B:361:ASN:ND2	3:B:603:HOH:O	2.53	0.41
1:B:318:GLU:HG2	1:B:337:SER:HB3	2.03	0.40
1:B:357:GLU:OE1	1:B:364:SER:OG	2.38	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/214 (98%)	201 (96%)	7 (3%)	2 (1%)	15	23
1	B	206/214 (96%)	192 (93%)	9 (4%)	5 (2%)	6	6
All	All	416/428 (97%)	393 (94%)	16 (4%)	7 (2%)	9	11

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	291	PRO
1	A	292	ARG
1	B	271	PRO
1	B	300	TYR
1	B	269	GLU
1	B	390	ASN
1	B	273	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/196 (100%)	192 (98%)	3 (2%)	65	80
1	B	195/196 (100%)	189 (97%)	6 (3%)	40	60
All	All	390/392 (100%)	381 (98%)	9 (2%)	50	70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	297	ASN
1	A	334	LYS
1	A	355	ARG
1	B	248	LYS
1	B	294	GLU
1	B	295	GLN
1	B	320	LYS
1	B	324	SER
1	B	354	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	295	GLN
1	B	421	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	C	1	1,2	14,14,15	0.72	1 (7%)	17,19,21	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	2	2	14,14,15	0.64	1 (7%)	17,19,21	0.51	0
2	BMA	C	3	2	11,11,12	1.14	1 (9%)	15,15,17	0.96	0
2	MAN	C	4	2	11,11,12	1.18	1 (9%)	15,15,17	1.06	1 (6%)
2	NAG	C	5	2	14,14,15	0.57	0	17,19,21	0.48	0
2	MAN	C	6	2	11,11,12	0.72	0	15,15,17	1.41	2 (13%)
2	NAG	C	7	2	14,14,15	0.43	0	17,19,21	0.49	0
2	FUC	C	8	2	10,10,11	0.89	0	14,14,16	1.13	2 (14%)
2	NAG	D	1	1,2	14,14,15	1.57	2 (14%)	17,19,21	1.54	2 (11%)
2	NAG	D	2	2	14,14,15	0.88	2 (14%)	17,19,21	0.65	0
2	BMA	D	3	2	11,11,12	1.22	2 (18%)	15,15,17	2.04	4 (26%)
2	MAN	D	4	2	11,11,12	1.26	2 (18%)	15,15,17	2.01	5 (33%)
2	NAG	D	5	2	14,14,15	0.64	0	17,19,21	1.68	3 (17%)
2	MAN	D	6	2	11,11,12	1.44	3 (27%)	15,15,17	1.26	1 (6%)
2	NAG	D	7	2	14,14,15	1.62	1 (7%)	17,19,21	1.23	1 (5%)
2	FUC	D	8	2	10,10,11	2.21	3 (30%)	14,14,16	1.45	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	0/6/23/26	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	NAG	C	7	2	-	2/6/23/26	0/1/1/1
2	FUC	C	8	2	-	-	0/1/1/1
2	NAG	D	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	1/2/19/22	0/1/1/1
2	MAN	D	4	2	-	2/2/19/22	0/1/1/1
2	NAG	D	5	2	-	2/6/23/26	0/1/1/1
2	MAN	D	6	2	-	2/2/19/22	0/1/1/1
2	NAG	D	7	2	-	2/6/23/26	0/1/1/1
2	FUC	D	8	2	-	-	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	7	NAG	O5-C1	-5.43	1.35	1.43
2	D	8	FUC	O5-C1	4.35	1.50	1.43
2	D	1	NAG	C1-C2	3.73	1.57	1.52
2	D	8	FUC	C1-C2	3.72	1.60	1.52
2	D	1	NAG	C8-C7	3.57	1.58	1.50
2	D	8	FUC	O5-C5	3.46	1.51	1.43
2	C	3	BMA	O5-C1	-2.79	1.39	1.43
2	C	1	NAG	O5-C1	-2.60	1.39	1.43
2	D	6	MAN	C1-C2	2.55	1.58	1.52
2	D	4	MAN	C4-C5	2.53	1.58	1.53
2	D	2	NAG	O5-C1	2.37	1.47	1.43
2	C	2	NAG	O5-C1	-2.34	1.40	1.43
2	D	6	MAN	O5-C1	-2.29	1.40	1.43
2	D	3	BMA	C4-C5	2.29	1.57	1.53
2	D	3	BMA	O5-C1	-2.23	1.40	1.43
2	D	6	MAN	C2-C3	2.22	1.55	1.52
2	D	4	MAN	O5-C5	2.15	1.47	1.43
2	C	4	MAN	C2-C3	2.10	1.55	1.52
2	D	2	NAG	C1-C2	2.03	1.55	1.52

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4	MAN	C1-O5-C5	5.40	119.51	112.19
2	D	5	NAG	C1-O5-C5	4.46	118.24	112.19
2	D	1	NAG	C2-N2-C7	4.19	128.87	122.90
2	D	3	BMA	O5-C1-C2	-4.09	104.46	110.77
2	D	5	NAG	C3-C4-C5	4.00	117.37	110.24
2	D	7	NAG	C4-C3-C2	3.95	116.81	111.02
2	C	6	MAN	C1-O5-C5	3.90	117.47	112.19
2	D	3	BMA	C3-C4-C5	3.84	117.09	110.24
2	C	6	MAN	O2-C2-C3	-3.30	103.52	110.14
2	D	3	BMA	C1-C2-C3	-3.27	105.64	109.67
2	D	6	MAN	O2-C2-C3	-3.01	104.11	110.14
2	D	8	FUC	O2-C2-C1	2.96	115.21	109.15
2	D	4	MAN	O2-C2-C3	-2.94	104.24	110.14
2	D	4	MAN	C3-C4-C5	2.82	115.28	110.24
2	C	4	MAN	C1-O5-C5	2.66	115.80	112.19
2	D	4	MAN	C1-C2-C3	-2.47	106.63	109.67
2	D	8	FUC	C1-C2-C3	2.43	112.65	109.67
2	C	8	FUC	C1-O5-C5	2.39	118.20	112.78
2	D	5	NAG	O5-C5-C4	2.36	116.57	110.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	BMA	C2-C3-C4	2.31	114.89	110.89
2	D	8	FUC	C1-O5-C5	2.29	117.96	112.78
2	C	8	FUC	C1-C2-C3	2.16	112.32	109.67
2	D	4	MAN	O5-C5-C4	2.11	115.97	110.83
2	D	1	NAG	O7-C7-C8	-2.07	118.22	122.06
2	D	8	FUC	O5-C5-C4	2.00	113.11	109.52

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	4	MAN	C4-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	C	7	NAG	O5-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	D	4	MAN	O5-C5-C6-O6
2	C	7	NAG	C4-C5-C6-O6
2	D	6	MAN	C4-C5-C6-O6
2	D	7	NAG	O5-C5-C6-O6
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	D	6	MAN	O5-C5-C6-O6
2	D	7	NAG	C4-C5-C6-O6
2	D	5	NAG	C1-C2-N2-C7
2	D	3	BMA	O5-C5-C6-O6
2	D	5	NAG	C3-C2-N2-C7


There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
2	D	1	NAG	1	0
2	D	2	NAG	1	0
2	D	3	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/214 (98%)	-0.04	6 (2%) 53 51	26, 42, 72, 117	0
1	B	208/214 (97%)	0.33	19 (9%) 9 8	28, 53, 108, 150	0
All	All	419/428 (97%)	0.15	25 (5%) 21 20	26, 47, 99, 150	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	ALA	12.5
1	A	235	ALA	10.3
1	B	323	VAL	6.4
1	B	273	VAL	5.2
1	B	324	SER	4.7
1	B	267	SER	3.7
1	B	266	VAL	3.6
1	B	292	ARG	3.5
1	B	296	TYR	3.4
1	B	294	GLU	3.2
1	B	269	GLU	3.1
1	A	333	GLU	2.9
1	B	271	PRO	2.9
1	B	272	GLU	2.7
1	B	270	ASP	2.7
1	B	318	GLU	2.5
1	B	298	SER	2.4
1	A	329	PRO	2.4
1	B	328	LEU	2.3
1	B	327	ALA	2.3
1	A	328	LEU	2.2
1	B	334	LYS	2.1
1	B	380	GLU	2.1
1	B	252	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	380	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	5	14/15	0.63	0.20	126,140,145,148	0
2	MAN	D	4	11/12	0.70	0.22	152,159,165,166	0
2	NAG	D	2	14/15	0.71	0.20	118,126,128,129	0
2	NAG	D	7	14/15	0.74	0.18	100,104,110,110	0
2	MAN	D	6	11/12	0.81	0.18	103,107,110,110	0
2	FUC	D	8	10/11	0.82	0.28	140,146,149,153	0
2	BMA	D	3	11/12	0.84	0.12	114,130,136,144	0
2	MAN	C	4	11/12	0.88	0.13	88,95,98,100	0
2	NAG	D	1	14/15	0.89	0.15	113,114,125,132	0
2	NAG	C	5	14/15	0.89	0.13	102,103,106,107	0
2	FUC	C	8	10/11	0.92	0.10	70,71,72,72	0
2	NAG	C	1	14/15	0.93	0.11	55,61,65,69	0
2	NAG	C	7	14/15	0.94	0.14	71,90,101,102	0
2	MAN	C	6	11/12	0.94	0.12	69,71,76,78	0
2	NAG	C	2	14/15	0.95	0.16	56,61,66,66	0
2	BMA	C	3	11/12	0.95	0.12	64,67,73,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.