



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2021 – 11:36 AM JST

PDB ID : 7CKH
Title : Crystal structure of TMSiPheRS
Authors : Sun, J.P.; Wang, J.Y.; Zhu, Z.L.; He, Q.T.; Xiao, P.
Deposited on : 2020-07-17
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.18
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 89% 9% . </div> </div>
1	B	314	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 11% 89% 9% .. </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	2	0
			2494	1591	428	462	13			
1	B	310	Total	C	N	O	S	0	2	0
			2497	1592	431	462	12			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	HIS	TYR	engineered mutation	UNP Q57834
A	63	GLY	ILE	engineered mutation	UNP Q57834
A	65	VAL	LEU	engineered mutation	UNP Q57834
A	70	GLN	HIS	engineered mutation	UNP Q57834
A	158	GLY	ASP	engineered mutation	UNP Q57834
A	159	GLY	ILE	engineered mutation	UNP Q57834
A	164	GLY	VAL	engineered mutation	UNP Q57834
A	307	LEU	-	expression tag	UNP Q57834
A	308	GLU	-	expression tag	UNP Q57834
A	309	HIS	-	expression tag	UNP Q57834
A	310	HIS	-	expression tag	UNP Q57834
A	311	HIS	-	expression tag	UNP Q57834
A	312	HIS	-	expression tag	UNP Q57834
A	313	HIS	-	expression tag	UNP Q57834
A	314	HIS	-	expression tag	UNP Q57834
B	32	HIS	TYR	engineered mutation	UNP Q57834
B	63	GLY	ILE	engineered mutation	UNP Q57834
B	65	VAL	LEU	engineered mutation	UNP Q57834
B	70	GLN	HIS	engineered mutation	UNP Q57834
B	158	GLY	ASP	engineered mutation	UNP Q57834
B	159	GLY	ILE	engineered mutation	UNP Q57834
B	164	GLY	VAL	engineered mutation	UNP Q57834
B	307	LEU	-	expression tag	UNP Q57834
B	308	GLU	-	expression tag	UNP Q57834
B	309	HIS	-	expression tag	UNP Q57834

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	310	HIS	-	expression tag	UNP Q57834
B	311	HIS	-	expression tag	UNP Q57834
B	312	HIS	-	expression tag	UNP Q57834
B	313	HIS	-	expression tag	UNP Q57834
B	314	HIS	-	expression tag	UNP Q57834

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	213	Total 213	O 213	0	0
2	B	168	Total 168	O 168	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.18Å 91.20Å 107.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.75 – 1.79 49.75 – 1.79	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.75-1.79) 99.4 (49.75-1.79)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.12rc0_2787	Depositor
R, R_{free}	0.206 , 0.245 0.206 , 0.245	Depositor DCC
R_{free} test set	3198 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.779	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5372	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2543	0.52	0/3411
1	B	0.34	0/2546	0.50	1/3415 (0.0%)
All	All	0.35	0/5089	0.51	1/6826 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	253	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2562	16	0
1	B	2497	0	2563	16	0
2	A	213	0	0	3	1
2	B	168	0	0	0	1
All	All	5372	0	5125	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ALA:HA	1:A:305[B]:ARG:HD2	1.62	0.80
1:A:305[A]:ARG:NH1	2:A:402:HOH:O	2.20	0.74
1:B:129:ARG:NH2	1:B:182:GLU:OE1	2.21	0.73
1:B:265:LEU:HD21	1:B:276:LEU:HD13	1.76	0.66
1:B:257:ARG:H	1:B:264:ASP:H	1.52	0.58
1:A:245:ALA:HA	1:A:249:LEU:HG	1.86	0.57
1:B:181:ARG:HB3	1:B:187:LYS:HE3	1.87	0.55
1:B:252:PRO:HA	1:B:268:ASN:HA	1.89	0.54
1:A:90:LYS:HE2	1:A:102:TYR:CD1	2.45	0.52
1:A:141:ASP:O	2:A:401:HOH:O	2.18	0.50
1:B:263:GLY:HA2	1:B:264:ASP:HB2	1.93	0.49
1:B:80:ASP:OD2	1:B:81:GLU:N	2.45	0.49
1:A:124:LYS:HB3	1:A:124:LYS:HE2	1.69	0.47
1:B:117:ASN:OD1	1:B:120:ARG:NH2	2.47	0.47
1:B:250:GLU:O	1:B:253:LEU:HB2	2.15	0.47
1:A:42:HIS:HA	1:A:214:ALA:HA	1.97	0.46
1:B:268:ASN:OD1	1:B:268:ASN:N	2.49	0.46
1:A:53:ILE:O	1:A:56:GLN:HG2	2.16	0.46
1:B:304:LYS:O	1:B:308:GLU:HG3	2.17	0.45
1:A:169:GLY:O	1:A:192:HIS:HA	2.16	0.45
1:A:95:ALA:CA	1:A:305[B]:ARG:HD2	2.43	0.45
1:A:9:ARG:NH2	2:A:410:HOH:O	2.50	0.44
1:A:13:GLU:OE1	1:A:192:HIS:ND1	2.39	0.44
1:A:129:ARG:HH12	1:A:182:GLU:CD	2.21	0.44
1:A:1:MET:HE2	1:A:5:GLU:HB3	1.99	0.44
1:B:9[B]:ARG:CZ	1:B:9[B]:ARG:HB3	2.47	0.43
1:A:205:MET:HG3	1:A:212:PHE:HB3	2.02	0.42
1:A:268:ASN:HB2	1:A:272:GLU:OE1	2.20	0.42
1:B:245:ALA:HA	1:B:249:LEU:HG	2.02	0.41
1:B:257:ARG:NH1	1:B:261:PHE:O	2.53	0.41
1:B:254:THR:HG23	1:B:266:THR:HB	2.02	0.41
1:B:42:HIS:HA	1:B:214:ALA:HA	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:490:HOH:O	2:B:536:HOH:O[2_455]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/314 (99%)	308 (99%)	2 (1%)	0	100	100
1	B	310/314 (99%)	309 (100%)	1 (0%)	0	100	100
All	All	620/628 (99%)	617 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/272 (99%)	262 (97%)	8 (3%)	41	27
1	B	270/272 (99%)	262 (97%)	8 (3%)	41	27
All	All	540/544 (99%)	524 (97%)	16 (3%)	43	27

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	ASP
1	A	84	LYS
1	A	139	ARG
1	A	187	LYS
1	A	275	SER
1	A	300	GLU
1	A	305[A]	ARG
1	A	305[B]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	81	GLU
1	B	87	ASP
1	B	112	LYS
1	B	120	ARG
1	B	253	LEU
1	B	254	THR
1	B	256	LYS
1	B	311	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/314 (98%)	0.42	4 (1%) 77 74	31, 43, 63, 70	0
1	B	310/314 (98%)	0.64	34 (10%) 5 4	32, 44, 67, 77	0
All	All	620/628 (98%)	0.53	38 (6%) 21 16	31, 43, 65, 77	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	265	LEU	7.7
1	B	267	VAL	4.9
1	B	253	LEU	4.7
1	B	282	LEU	4.5
1	B	262	GLY	4.3
1	B	279	ASN	3.7
1	B	285	MET	3.6
1	B	266	THR	3.5
1	B	255	ILE	3.5
1	B	284	PRO	3.3
1	B	311	HIS	3.1
1	B	258	PRO	3.0
1	B	233	ALA	3.0
1	B	229	ALA	3.0
1	B	257	ARG	2.8
1	B	256	LYS	2.8
1	B	230	TYR	2.7
1	B	199	LEU	2.7
1	B	278	LYS	2.6
1	B	291	VAL	2.6
1	B	283	HIS	2.6
1	B	252	PRO	2.6
1	B	261	PHE	2.5
1	B	264	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	231	CYS	2.5
1	B	281	GLU	2.5
1	B	275	SER	2.5
1	A	309	HIS	2.4
1	B	254	THR	2.4
1	A	265	LEU	2.3
1	A	310	HIS	2.3
1	B	287	LEU	2.2
1	B	286	ASP	2.2
1	B	228	LYS	2.2
1	B	235	VAL	2.2
1	B	263	GLY	2.2
1	A	161	TYR	2.1
1	B	277	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.