



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 07:19 pm BST

PDB ID : 3CNS  
Title : Crystal structure of fms1 in complex with S-Bz-MeSpermidine  
Authors : Huang, Q.; Hao, H.  
Deposited on : 2008-03-26  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	<div> <div>4%</div> <div>71%</div> <div>21%</div> <div>• 5%</div> </div>
1	B	516	<div> <div>5%</div> <div>64%</div> <div>28%</div> <div>• 6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SP8	A	518	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called fms1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	492	Total	C	N	O	S	0	0	0
			3935	2489	684	740	22			
1	B	487	Total	C	N	O	S	0	0	0
			3896	2462	676	736	22			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	509	LEU	-	EXPRESSION TAG	UNP P50264
A	510	GLU	-	EXPRESSION TAG	UNP P50264
A	511	HIS	-	EXPRESSION TAG	UNP P50264
A	512	HIS	-	EXPRESSION TAG	UNP P50264
A	513	HIS	-	EXPRESSION TAG	UNP P50264
A	514	HIS	-	EXPRESSION TAG	UNP P50264
A	515	HIS	-	EXPRESSION TAG	UNP P50264
A	516	HIS	-	EXPRESSION TAG	UNP P50264
B	509	LEU	-	EXPRESSION TAG	UNP P50264
B	510	GLU	-	EXPRESSION TAG	UNP P50264
B	511	HIS	-	EXPRESSION TAG	UNP P50264
B	512	HIS	-	EXPRESSION TAG	UNP P50264
B	513	HIS	-	EXPRESSION TAG	UNP P50264
B	514	HIS	-	EXPRESSION TAG	UNP P50264
B	515	HIS	-	EXPRESSION TAG	UNP P50264
B	516	HIS	-	EXPRESSION TAG	UNP P50264

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0
			53	27	9	15	2	
2	B	1	Total	C	N	O	P	0
			53	27	9	15	2	

- Molecule 3 is N-{(1S)-3-[(4-aminobutyl)amino]-1-methylpropyl}benzamide (three-letter code: SP8) (formula: C<sub>15</sub>H<sub>25</sub>N<sub>3</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			19	15	3	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			19	15	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	63	Total	O	0	0
			63	63		
4	B	52	Total	O	0	0
			52	52		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
- 
- 4%
- 71%
- 21%
- 5%
- 4487
- 4508
- 4509
- 4510
- 4511
- 4512
- 4513
- 4514
- 4515
- 4516
- 4517
- 4518
- 4519
- 4520
- 4521
- 4522
- 4523
- 4524
- 4525
- 4526
- 4527
- 4528
- 4529
- 4530
- 4531
- 4532
- 4533
- 4534
- 4535
- 4536
- 4537
- 4538
- 4539
- 4540
- 4541
- 4542
- 4543
- 4544
- 4545
- 4546
- 4547
- 4548
- 4549
- 4550
- 4551
- 4552
- 4553
- 4554
- 4555
- 4556
- 4557
- 4558
- 4559
- 4560
- 4561
- 4562
- 4563
- 4564
- 4565
- 4566
- 4567
- 4568
- 4569
- 4570
- 4571
- 4572
- 4573
- 4574
- 4575
- 4576
- 4577
- 4578
- 4579
- 4580
- 4581
- 4582
- 4583
- 4584
- 4585
- 4586
- 4587
- 4588
- 4589
- 4590
- 4591
- 4592
- 4593
- 4594
- 4595
- 4596
- 4597
- 4598
- 4599
- 4600
- 4601
- 4602
- 4603
- 4604
- 4605
- 4606
- 4607
- 4608
- 4609
- 4610
- 4611
- 4612
- 4613
- 4614
- 4615
- 4616
- 4617
- 4618
- 4619
- 4620
- 4621
- 4622
- 4623
- 4624
- 4625
- 4626
- 4627
- 4628
- 4629
- 4630
- 4631
- 4632
- 4633
- 4634
- 4635
- 4636
- 4637
- 4638
- 4639
- 4640
- 4641
- 4642
- 4643
- 4644
- 4645
- 4646
- 4647
- 4648
- 4649
- 4650
- 4651
- 4652
- 4653
- 4654
- 4655
- 4656
- 4657
- 4658
- 4659
- 4660
- 4661
- 4662
- 4663
- 4664
- 4665
- 4666
- 4667
- 4668
- 4669
- 4670
- 4671
- 4672
- 4673
- 4674
- 4675
- 4676
- 4677
- 4678
- 4679
- 4680
- 4681
- 4682
- 4683
- 4684
- 4685
- 4686
- 4687
- 4688
- 4689
- 4690
- 4691
- 4692
- 4693
- 4694
- 4695
- 4696
- 4697
- 4698
- 4699
- 4700
- 4701
- 4702
- 4703
- 4704
- 4705
- 4706
- 4707
- 4708
- 4709
- 4710
- 4711
- 4712
- 4713
- 4714
- 4715
- 4716
- 4717
- 4718
- 4719
- 4720
- 4721
- 4722
- 4723
- 4724
- 4725
- 4726
- 4727
- 4728
- 4729
- 4730
- 4731
- 4732
- 4733
- 4734
- 4735
- 4736
- 4737
- 4738
- 4739
- 4740
- 4741
- 4742
- 4743
- 4744
- 4745
- 4746
- 4747
- 4748
- 4749
- 4750
- 4751
- 4752
- 4753
- 4754
- 4755
- 4756
- 4757
- 4758
- 4759
- 4760
- 4761
- 4762
- 4763
- 4764
- 4765
- 4766
- 4767
- 4768
- 4769
- 4770
- 4771
- 4772
- 4773
- 4774
- 4775
- 4776
- 4777
- 4778
- 4779
- 4780
- 4781
- 4782
- 4783
- 4784
- 4785
- 4786
- 4787
- 4788
- 4789
- 4790
- 4791
- 4792
- 4793
- 4794
- 4795
- 4796
- 4797
- 4798
- 4799
- 4800
- 4801
- 4802
- 4803
- 4804
- 4805
- 4806
- 4807
- 4808
- 4809
- 4810
- 4811
- 4812
- 4813
- 4814
- 4815
- 4816
- 4817
- 4818
- 4819
- 4820
- 4821
- 4822
- 4823
- 4824
- 4825
- 4826
- 4827
- 4828
- 4829
- 4830
- 4831
- 4832
- 4833
- 4834
- 4835
- 4836
- 4837
- 4838
- 4839
- 4840
- 4841
- 4842
- 4843
- 4844
- 4845
- 4846
- 4847
- 4848
- 4849
- 4850
- 4851
- 4852
- 4853
- 4854
- 4855
- 4856
- 4857
- 4858
- 4859
- 4860
- 4861
- 4862
- 4863
- 4864
- 4865
- 4866
- 4867
- 4868
- 4869
- 4870
- 4871
- 4872
- 4873
- 4874
- 4875
- 4876
- 4877
- 4878
- 4879
- 4880
- 4881
- 4882
- 4883
- 4884
- 4885
- 4886
- 4887
- 4888
- 4889
- 

- [illegible]



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.28Å 215.13Å 117.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.40 39.66 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.00-2.40) 98.0 (39.66-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.85 (at 1.89Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.223 , 0.272 0.219 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.8	Xtriage
Anisotropy	0.708	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SP8, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/4016	0.62	0/5429
1	B	0.36	0/3973	0.59	0/5370
All	All	0.37	0/7989	0.61	0/10799

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3935	0	3862	112	0
1	B	3896	0	3811	128	0
2	A	53	0	31	3	0
2	B	53	0	31	2	0
3	A	19	0	25	11	0
3	B	19	0	25	4	0
4	A	63	0	0	1	0
4	B	52	0	0	1	0
All	All	8090	0	7785	244	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 244 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:LEU:O	3:A:518:SP8:HAB	1.50	1.07
1:A:67:HIS:HE2	3:A:518:SP8:HNAL	0.98	0.94
1:B:382:ASN:HD22	1:B:382:ASN:H	1.10	0.93
1:A:132:GLN:HG2	1:A:134:LEU:HD13	1.53	0.90
1:A:385:GLU:O	1:A:388:ARG:HG2	1.70	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/516 (94%)	453 (93%)	26 (5%)	7 (1%)	11	15
1	B	477/516 (92%)	436 (91%)	32 (7%)	9 (2%)	8	10
All	All	963/1032 (93%)	889 (92%)	58 (6%)	16 (2%)	9	11

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	200	ASN
1	A	265	GLU
1	B	200	ASN
1	A	104	ARG
1	B	262	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/457 (95%)	417 (96%)	16 (4%)	34	53
1	B	429/457 (94%)	410 (96%)	19 (4%)	28	45
All	All	862/914 (94%)	827 (96%)	35 (4%)	30	48

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	388	ARG
1	B	173	LEU
1	B	391	LYS
1	B	72	ASN
1	B	118	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	72	ASN
1	B	255	GLN
1	B	434	ASN
1	B	109	HIS
1	B	195	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	FAD	A	517	-	51,58,58	1.62	7 (13%)	60,89,89	1.58	9 (15%)
3	SP8	A	518	-	19,19,19	0.47	0	20,22,22	0.74	0
2	FAD	B	517	-	51,58,58	1.63	6 (11%)	60,89,89	1.56	9 (15%)
3	SP8	B	518	-	19,19,19	0.48	0	20,22,22	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	517	-	-	1/30/50/50	0/6/6/6
3	SP8	A	518	-	-	2/16/16/16	0/1/1/1
2	FAD	B	517	-	-	2/30/50/50	0/6/6/6
3	SP8	B	518	-	-	2/16/16/16	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	517	FAD	C10-N1	5.61	1.40	1.33
2	B	517	FAD	C10-N1	5.55	1.40	1.33
2	B	517	FAD	C4X-N5	5.03	1.40	1.33
2	A	517	FAD	C4X-N5	4.92	1.40	1.33
2	A	517	FAD	C4-N3	4.25	1.40	1.33

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	517	FAD	C4-N3-C2	5.83	120.06	115.14
2	B	517	FAD	C4-N3-C2	5.78	120.02	115.14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	517	FAD	N3A-C2A-N1A	-4.57	121.54	128.68
2	B	517	FAD	N3A-C2A-N1A	-4.46	121.71	128.68
2	A	517	FAD	O4B-C1B-C2B	-3.99	101.10	106.93

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	518	SP8	CAB-CAI-NAH-CAG
3	B	518	SP8	CAJ-CAK-NAL-CAM
3	B	518	SP8	CAJ-CAI-NAH-CAG
2	A	517	FAD	O4B-C4B-C5B-O5B
2	B	517	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	517	FAD	3	0
3	A	518	SP8	11	0
2	B	517	FAD	2	0
3	B	518	SP8	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	492/516 (95%)	-0.10	19 (3%) 39 38	27, 43, 69, 86	0
1	B	487/516 (94%)	-0.06	25 (5%) 28 26	28, 45, 83, 109	0
All	All	979/1032 (94%)	-0.08	44 (4%) 33 31	27, 44, 77, 109	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	425	ALA	7.1
1	B	424	ILE	6.9
1	A	428	ASN	5.6
1	A	420	PRO	4.8
1	B	426	ASN	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SP8	B	518	19/19	0.69	0.30	58,65,70,70	0
3	SP8	A	518	19/19	0.71	0.30	58,67,71,71	0
2	FAD	B	517	53/53	0.96	0.14	25,34,39,40	0
2	FAD	A	517	53/53	0.96	0.16	24,29,33,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around SP8 A 518:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD B 517:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.