



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 06:57 pm BST

PDB ID : 4CT6
Title : CNOT9-CNOT1 complex
Authors : Basquin, J.; Ozgur, S.; Conti, E.
Deposited on : 2014-03-12
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	<div> <div>11%</div> <div> <div></div> <div>85%</div> <div>11%</div> <div>..</div> </div> </div>
2	B	270	<div> <div>%</div> <div> <div></div> <div>94%</div> <div>..</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4378 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	0	0
			1833	1148	336	331	18			

- Molecule 2 is a protein called CELL DIFFERENTIATION PROTEIN RCD1 HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	269	Total	C	N	O	S	0	0	0
			2128	1365	363	390	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	ALA	-	expression tag	UNP Q92600
B	17	ALA	-	expression tag	UNP Q92600


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total	O	0	0
			211	211		
3	B	206	Total	O	0	0
			206	206		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1



• Molecule 2: CELL DIFFERENTIATION PROTEIN RCD1 HOMOLOG



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	153.67Å 66.14Å 72.44Å 90.00° 100.35° 90.00°	Depositor
Resolution (Å)	47.90 – 2.10 47.90 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.6 (47.90-2.10) 99.3 (47.90-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.10Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.194 , 0.236 0.197 , 0.239	Depositor DCC
R_{free} test set	2088 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4378	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1868	0.57	1/2530 (0.0%)
2	B	0.44	0/2170	0.55	0/2953
All	All	0.44	0/4038	0.56	1/5483 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1478	ARG	C-N-CA	5.71	135.99	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1833	0	1832	20	0
2	B	2128	0	2185	12	0
3	A	211	0	0	11	0
3	B	206	0	0	3	0
All	All	4378	0	4017	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:264:GLN:HE21	2:B:264:GLN:HA	1.07	1.12
2:B:21:LYS:O	3:B:2003:HOH:O	1.71	1.08
2:B:264:GLN:NE2	2:B:264:GLN:HA	1.83	0.91
1:A:1478:ARG:N	1:A:1479:THR:O	2.07	0.88
1:A:1561:VAL:O	3:A:2122:HOH:O	2.07	0.72
1:A:1366:SER:OG	3:A:2059:HOH:O	2.10	0.69
1:A:1365:TYR:O	3:A:2057:HOH:O	2.11	0.68
1:A:1415:ILE:O	3:A:2106:HOH:O	2.14	0.66
2:B:43:SER:O	2:B:46:ARG:NH1	2.32	0.62
1:A:1363:ASN:OD1	3:A:2054:HOH:O	2.16	0.60
2:B:77:ILE:HD11	2:B:122:PHE:CD1	2.36	0.60
2:B:26:ILE:O	3:B:2011:HOH:O	2.17	0.58
1:A:1368:ALA:N	3:A:2059:HOH:O	2.39	0.55
1:A:1478:ARG:CB	1:A:1479:THR:HG22	2.37	0.54
1:A:1574:GLU:OE2	3:A:2197:HOH:O	2.19	0.52
1:A:1433:GLU:OE1	3:A:2128:HOH:O	2.19	0.51
2:B:278:GLN:NE2	3:B:2167:HOH:O	2.43	0.51
2:B:77:ILE:HD11	2:B:122:PHE:HD1	1.76	0.51
1:A:1398:GLU:OE2	3:A:2062:HOH:O	2.18	0.50
1:A:1564:VAL:HG12	3:A:2124:HOH:O	2.12	0.50
1:A:1560:LYS:HB2	1:A:1563:GLY:O	2.12	0.49
1:A:1555:GLU:OE2	1:A:1558:ARG:NH1	2.47	0.48
1:A:1377:ASN:HB3	1:A:1380:ILE:HG13	1.96	0.47
1:A:1383:PHE:CZ	1:A:1495:LEU:HD11	2.50	0.47
1:A:1382:LEU:HD22	1:A:1488:MET:HA	1.95	0.46
2:B:77:ILE:HD11	2:B:122:PHE:CE1	2.52	0.45
1:A:1519:MET:HE2	1:A:1523:LEU:HG	2.00	0.43
2:B:163:ILE:O	2:B:163:ILE:HG12	2.18	0.42
1:A:1537:GLY:HA2	3:A:2162:HOH:O	2.18	0.42
2:B:31:SER:HA	2:B:32:PRO:HD3	1.90	0.42
1:A:1443:HIS:CG	1:A:1576:ALA:HB2	2.55	0.42
2:B:103:HIS:HA	2:B:104:PRO:HD3	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/243 (97%)	228 (97%)	6 (3%)	1 (0%)	34	32
2	B	267/270 (99%)	264 (99%)	3 (1%)	0	100	100
All	All	502/513 (98%)	492 (98%)	9 (2%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1479	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/209 (93%)	189 (97%)	5 (3%)	46	50
2	B	238/242 (98%)	235 (99%)	3 (1%)	69	75
All	All	432/451 (96%)	424 (98%)	8 (2%)	57	63

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1364	VAL
1	A	1366	SER
1	A	1479	THR
1	A	1571	VAL
1	A	1588	LEU
2	B	176	GLU
2	B	264	GLN
2	B	278	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1391	GLN
2	B	264	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/243 (97%)	0.83	26 (10%) 5 7	32, 55, 98, 124	0
2	B	269/270 (99%)	0.05	4 (1%) 73 77	31, 44, 75, 100	0
All	All	506/513 (98%)	0.42	30 (5%) 22 27	31, 49, 83, 124	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1479	THR	8.1
1	A	1562	GLY	6.3
1	A	1563	GLY	6.2
1	A	1477	LEU	6.2
1	A	1480	ALA	5.3
1	A	1483	GLN	5.3
2	B	17	ALA	5.1
1	A	1482	PRO	3.8
1	A	1486	GLU	3.8
1	A	1389	LEU	3.6
1	A	1476	ALA	3.5
1	A	1481	SER	3.5
1	A	1485	ARG	3.5
1	A	1587	ASP	3.5
1	A	1478	ARG	3.3
1	A	1565	ASP	3.2
1	A	1567	LYS	3.1
1	A	1392	CYS	3.1
1	A	1484	GLN	2.7
1	A	1474	ALA	2.7
1	A	1588	LEU	2.6
1	A	1387	PRO	2.5
2	B	18	ASP	2.4
1	A	1487	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	266	LEU	2.3
1	A	1393	VAL	2.2
1	A	1420	CYS	2.2
1	A	1352	PRO	2.1
1	A	1353	PRO	2.1
2	B	16	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.